




VISIT...

LANZAROTE
Caliente.COM





1. *Coryphantha runyonii*.

2. *Dolichothele sphaerica*.

THE CACTACEAE

DESCRIPTIONS AND ILLUSTRATIONS OF PLANTS OF THE CACTUS FAMILY

BY

N. L. BRITTON AND J. N. ROSE

Volume IV



THE CARNEGIE INSTITUTION OF WASHINGTON
WASHINGTON, DECEMBER 24, 1923

CARNEGIE INSTITUTION OF WASHINGTON
PUBLICATION No. 248 VOLUME IV

Pages 1-80, text only, were distributed
under date of October 9, 1923

PRESS OF GIBSON BROTHERS, INC
WASHINGTON

CONTENTS.

Tribe 3. Cereeae— <i>continued from Vol. III</i>	PAGE
Subtribe 6, Coryphanthanae	3
Key to Genera	3
Ancistrocactus	3
Thelocactus	6
Neolloydia	14
Mamillopsis	19
Cochemiea	21
Coryphantha	23
Neobesseyia	51
Escobaria	53
Bartschella	57
Pelecyphora	59
Phellosperma	60
Dolichothele	61
Solisia	64
Neomammillaria	65
Subtribe 7, Epiphyllanae	177
Key to Genera	177
Zygocactus	177
Epiphyllanthus	180
Schlumbergera	182
Epiphyllum	185
Disocactus	201
Chiapasias	203
Eccremocactus	204
Nopalxochia	204
Wittia	206
Subtribe 8, Rhipsalidanae	208
Key to Genera	208
Erythrorhipsalis	208
Rhipsalidopsis	209
Pfeiffera	210
Acanthorhipsalis	211
Pseudorhipsalis	213
Lepismium	215
Hatiora	216
Rhipsalis	219
Appendix	249
Index	291

ILLUSTRATIONS.

	PLATES	FACING PAGE
Plate 1.	(1) Plant of <i>Coryphantha runyonii</i> from Texas. (2) Plant of <i>Dolichothele sphaerica</i> from Texas Frontispiece.	
Plate 2.	(1) Fruiting plant of <i>Coryphantha neo-mexicana</i> . (1 <i>a</i>) Fruit of same. (2) Top of flowering plant of <i>Ancistrocactus scheeri</i> . (3) Flowering plant of <i>Cochemia poselgeri</i> . (3 <i>a</i>) Fruit of same. (3 <i>b</i>) Seed of same. (4) Flowering plant of <i>Coryphantha cornifera</i>	4
Plate 3.	(1) <i>Coryphantha nickelsae</i> from Monterey, Mexico. (2) <i>Neobesseyia similis</i> from Texas	34
Plate 4.	(1) <i>Coryphantha aggregata</i> from Arizona	46
Plate 5.	(1) A flowering plant of <i>Coryphantha cubensis</i> . (1 <i>a</i>) Fruit of same. (1 <i>b</i>) Tubercle of same. (2) Flowering plant of <i>Neomammillaria confusa</i> . (3) Flowering plant of <i>Neomammillaria geminispina</i> . (4) Top of flowering plant of <i>Coryphantha durangensis</i> . (5) A flowering plant of <i>Coryphantha arizonica</i> . (6) Flowering plant of <i>Coryphantha bumamma</i> . (7) Flowering plant of <i>Coryphantha chlorantha</i>	48
Plate 6.	(1) <i>Escobaria runyonii</i> from Texas. (2) <i>Neomammillaria hemisphaerica</i> from Texas	56
Plate 7.	(1) Flowering plant of <i>Escobaria dasyacantha</i> . (2) Fruit of <i>Dolichothele sphaerica</i> . (2 <i>a</i>) Seed of same. (3) Flowering plant of <i>Neomammillaria arida</i> . (4) Flowering plant of <i>Escobaria bella</i> . (4 <i>a</i>) Tubercle of same. (5) Flowering plant of <i>Neomammillaria crocidata</i> . (6) Flowering plant of <i>Bartschella schumannii</i> . (7) A flowering plant of <i>Neomammillaria carnea</i>	58
Plate 8.	(1) Fruiting plant of <i>Neomammillaria gaumeri</i> . (2) Flowering plant of <i>Neomammillaria heyderi</i> . (2 <i>a</i>) Fruit of same. (3) Flowering and fruiting plant of <i>Neomammillaria hemisphaerica</i> . (4) Flowering plant of <i>Neomammillaria compressa</i> . (5) Flowering plant of <i>Neomammillaria geminispina</i> . (6) Flowering plant of <i>Neomammillaria hemisphaerica</i>	72
Plate 9.	(1) Flowering and fruiting plant of <i>Neomammillaria applanata</i> . (2) Top of fruiting plant of <i>Neomammillaria karwinskiana</i> . (3) Top of flowering plant of <i>Neomammillaria aureiceps</i> . (4) Flowering plant of <i>Neomammillaria macracantha</i> . (5) Fruiting plant of <i>Neomammillaria mystax</i>	76
Plate 10.	(1) <i>Coryphantha sulcata</i> from Sabinal, Texas. (2) <i>Neomammillaria runyonii</i> from Monterey, Mexico	80
Plate 11.	(1) Flowering plant of <i>Neomammillaria magnimamma</i> . (2) Flowering plant of <i>Neomammillaria karwinskiana</i> . (3) Fruiting plant of <i>Neomammillaria gigantea</i> . (4) Fruiting plant of <i>Neobesseyia missouriensis</i>	96
Plate 12.	(1) Flowering plant of <i>Neomammillaria pseudoperbella</i> . (2) Top of flowering plant of <i>Neomammillaria spinosissima</i> . (3) Flowering plant of <i>Neomammillaria dealbata</i> . (4) Flowering plant of <i>Neomammillaria amoena</i> . (5) Flowering plant of <i>Neomammillaria polyedra</i> . (6) Flowering plant of <i>Neomammillaria celsiana</i>	110
Plate 13.	(1) <i>Neomammillaria wilcoxii</i> from Calabasas, Arizona. (2) <i>Neomammillaria gaumeri</i> from Yucatan, Mexico	154
Plate 14.	(1) Flowering plant of <i>Neomammillaria kunzeana</i> . (2) Flowering plant of <i>Neomammillaria bocasana</i> . (3) Flowering plant of <i>Neomammillaria decipiens</i> . (4) Top of flowering plant of <i>Neomammillaria armillata</i> . (5) Flowering plant of <i>Neomammillaria multiceps</i> . (6) Flowering plant of <i>Neomammillaria multiceps</i> . (7) Flowering plant of <i>Neomammillaria palmeri</i> . (8) Flowering plant of <i>Neomammillaria wildii</i>	156
Plate 15.	(1) Flowering plant of <i>Neomammillaria bombycina</i> . (2) Flowering plant of <i>Neomammillaria armillata</i> . (3) Top of flowering plant of <i>Neomammillaria armillata</i> . (4) Flowering plant of <i>Neomammillaria armillata</i> . (5) Flowering plant of <i>Neomammillaria goodridgei</i>	158
Plate 16.	(1) Flowering plant of <i>Epiphyllum darrahii</i> . (2) Top of flowering plant of <i>Epiphyllum pittieri</i>	190
Plate 17.	(1) End of branch of <i>Epiphyllum macropterum</i> . (2) Base of branch of <i>Epiphyllum macropterum</i>	192
Plate 18.	<i>Epiphyllum pittieri</i> from Costa Rica	194
Plate 19.	<i>Epiphyllum hookeri</i> from Tobago, West Indies	198
Plate 20.	<i>Eccremocactus bradei</i> from Costa Rica	204
Plate 21.	(1) Fruiting branch of <i>Rhipsalis grandiflora</i> . (2) Flowering branch of <i>Rhipsalis lindbergiana</i> . (3) Fruiting branch of <i>Rhipsalis shaferi</i> . (4) Flowering and fruiting branch of <i>Rhipsalis lindbergiana</i> . (5) Flowering plant of <i>Erythrorhipsalis pilocarpa</i> . (6) Flowering branch of <i>Rhipsalis grandiflora</i>	208
Plate 22.	(1) Flowering branch of <i>Pfeiffera ianthothele</i> . (2) Flowering and fruiting branch of <i>Lepismium cruciforme</i> . (3) Top of fruiting branch of <i>Pfeiffera ianthothele</i> . (4) Flowering and fruiting branch of <i>Rhipsalis jamaicensis</i> . (5) Flowering branch of <i>Pseudorhipsalis alata</i> . (6) Flowering and fruiting branch of <i>Pseudorhipsalis himantoclada</i> . (7) Flowering branch of <i>Pfeiffera ianthothele</i>	210
Plate 23.	(1) Flowering branch of <i>Hattoria cylindrica</i> . (2) Fruiting branch of <i>Rhipsalis heteroclada</i> . (3) Fruiting branch of <i>Rhipsalis cribrata</i> . (4) Flowering branch of <i>Hattoria salicornioides</i>	218

Plate 24.	(1) Flowering branch of <i>Rhipsalis leucorhaphis</i> . (2) Fruiting branch of <i>Rhipsalis leucorhaphis</i> . (3) Fruiting branch of <i>Rhipsalis megalantha</i> . (4) Fruiting branch of <i>Rhipsalis neves-armondii</i> . (5) Fruiting branch of <i>Rhipsalis neves-armondii</i> . (6) Fruiting branch of <i>Rhipsalis pittieri</i> . (7) Flowering and fruiting branch of <i>Rhipsalis shaferi</i> . (8) Fruiting branch of <i>Rhipsalis aculeata</i> . (9) Fruiting branch of <i>Rhipsalis mesembryanthemoides</i> . (10) Flowering branch of <i>Rhipsalis mesembryanthemoides</i>	220
Plate 25.	(1) Fruiting branch of <i>Rhipsalis heteroclada</i> . (2) Fruiting branch of <i>Rhipsalis heteroclada</i> . (3) Fruiting branch of <i>Rhipsalis capilliformis</i> . (4) Fruiting branch of <i>Rhipsalis virgata</i>	222
Plate 26.	(1) Flowering branch of <i>Rhipsalis cribrata</i> . (2) Fruiting branch of <i>Rhipsalis capilliformis</i> . (3) Flowering branch of <i>Rhipsalis capilliformis</i> . (4) Flowering branch of <i>Rhipsalis capilliformis</i> . (5) Flowering branch of <i>Rhipsalis teres</i>	224
Plate 27.	(1) Flowering and fruiting branch of <i>Rhipsalis cassutha</i> . (2) Fruiting branch of <i>Rhipsalis burchellii</i> . (3) Flowering branch of <i>Rhipsalis cereuscula</i>	226
Plate 28.	(1) Flowering branch of <i>Rhipsalis neves-armondii</i> . (2) Flowering branch of <i>Rhipsalis paradoxa</i> . (3) Fruiting branch of <i>Rhipsalis pulvinigera</i> . (4) Flowering branch of <i>Rhipsalis tucumanensis</i> . (5) Fruiting branch of <i>Rhipsalis tucumanensis</i>	228
Plate 29.	(1) Flowering branch of <i>Rhipsalis floccosa</i> . (2) Flowering branch of <i>Rhipsalis floccosa</i> . (3) Fruiting branch of <i>Rhipsalis puniceo-discus</i> . (4) Flowering branch of <i>Rhipsalis gibberula</i> . (5) Branch of <i>Rhipsalis dissimilis</i> . (6) Flowering and fruiting branch of <i>Rhipsalis dissimilis</i>	230
Plate 30.	(1) Fruiting branch of <i>Rhipsalis gonocarpa</i> . (2) Flowering branches of <i>Rhipsalis warmingiana</i> . (3) Fruiting branch of <i>Rhipsalis tonduzii</i> . (4) Fruiting branch of <i>Rhipsalis trigona</i> . (5) Flowering branch of <i>Rhipsalis pentaptera</i> . (6) Fruiting branch of <i>Rhipsalis pentaptera</i>	232
Plate 31.	(1) A flowering branch of <i>Rhipsalis grandiflora</i> . (2) Flowering branch of <i>Rhipsalis pulchra</i> . (3) Flowering branch of <i>Rhipsalis grandiflora</i>	234
Plate 32.	(1) A fruiting branch of <i>Rhipsalis heteroclada</i> . (2) Fruiting branch of <i>Disocactus biformis</i> . (3) Flowering and fruiting branch of <i>Rhipsalis prismatica</i> . (4) Flowering branch of <i>Rhipsalis coriacea</i> . (5) Fruiting branch of <i>Rhipsalis coriacea</i> . (6) Fruiting branch of <i>Rhipsalis dissimilis</i> . (7) Branch of <i>Rhipsalis dissimilis</i>	236
Plate 33.	(1) Flowering branch of <i>Rhipsalis houlettiana</i> . (2) Flowering branch of <i>Rhipsalis houlettiana</i> . (3) Flower of <i>Rhipsalis houlettiana</i> . (4) Fruiting branch of <i>Rhipsalis houlettiana</i>	238
Plate 34.	(1) Flowering branch of <i>Rhipsalis houlettiana</i> . (2) Flower of <i>Rhipsalis houlettiana</i> . (3) Fruiting branch of <i>Rhipsalis warmingiana</i> . (4) Fruiting branch of <i>Rhipsalis warmingiana</i>	240
Plate 35.	(1) Fruiting branch of <i>Rhipsalis oblonga</i> . (2) Fruiting branch of <i>Rhipsalis elliptica</i> . (3) Flowering branch of <i>Rhipsalis crispata</i>	242
Plate 36.	(1) Flowering branch of <i>Rhipsalis pachyptera</i> . (2) Flowering branch of <i>Rhipsalis rhombea</i>	244
Plate 37.	(1) A flowering branch of <i>Rhipsalis russellii</i> . (2) Cluster of flowers of <i>Rhipsalis russellii</i> . (3) Enlarged flower of <i>Rhipsalis russellii</i> . (4) Fruiting branch of <i>Rhipsalis russellii</i> . (5) Fruiting branch of <i>Rhipsalis crispimarginata</i> . (6) Fruiting branch of <i>Rhipsalis pachyptera</i>	246

TEXT-FIGURES.

Fig.		PAGE.	Fig.		PAGE.
1.	<i>Ancistrocactus megarhizus</i>	4	22.	<i>Cochemiea halei</i>	21
2.	<i>Ancistrocactus scheeri</i>	5	23.	<i>Cochemiea poselgeri</i>	21
3.	<i>Ancistrocactus brevihamatus</i>	5	24.	Fruit and seed of <i>Cochemiea pondii</i>	23
4.	<i>Thelocactus rinconensis</i>	7	24a.	Fruit and seed of <i>Cochemiea pondii</i>	23
5.	<i>Thelocactus phymatothele</i>	7	25.	Tubercles of <i>Coryphantha macromeris</i>	25
6.	<i>Thelocactus buekii</i>	9	26.	<i>Coryphantha recurvata</i>	27
7.	<i>Thelocactus leucacanthus</i>	9	27.	<i>Coryphantha recurvata</i>	27
8.	<i>Thelocactus nidulans</i>	10	28.	<i>Coryphantha muehlenpfordtii</i>	29
9.	<i>Thelocactus fossulatus</i>	10	29.	<i>Coryphantha bumamma</i>	29
10.	<i>Thelocactus tulensis</i>	11	30.	<i>Coryphantha erecta</i>	32
11.	<i>Thelocactus bicolor</i>	11	31.	<i>Coryphantha erecta</i>	32
12.	<i>Neolloydia pilispina</i>	14	31a.	<i>Coryphantha pectinata</i>	35
13.	<i>Neolloydia horripila</i>	14	31b.	<i>Coryphantha echinus</i>	35
14.	<i>Neolloydia clavata</i>	15	32.	<i>Coryphantha nickelsae</i>	36
15.	<i>Neolloydia conoidea</i>	15	33.	<i>Coryphantha compacta</i>	36
16.	<i>Neolloydia ceratites</i>	17	34.	<i>Coryphantha radians</i>	37
17.	<i>Neolloydia conoidea</i>	17	35.	<i>Coryphantha sulcolanata</i>	37
18.	<i>Neolloydia texensis</i>	18	36.	<i>Coryphantha retusa</i>	38
19.	<i>Mamilloopsis senilis</i>	19	37.	<i>Coryphantha salm-dyckiana</i>	38
20.	<i>Mamilloopsis senilis</i>	19	38.	<i>Coryphantha pallida</i>	40
21.	<i>Mamilloopsis senilis</i>	20	39.	<i>Coryphantha pycnacantha</i>	40

TEXT-FIGURES—continued.

FIG.		PAGE.	FIG.		PAGE.
40.	<i>Coryphantha durangensis</i>	43	103.	<i>Neomammillaria collinsii</i>	204
41.	<i>Coryphantha durangensis</i>	43	104.	<i>Neomammillaria conzattii</i>	105
42.	<i>Coryphantha chlorantha</i>	44	105.	<i>Neomammillaria lanata</i>	105
43.	<i>Coryphantha neo-mexicana</i>	44	106.	<i>Neomammillaria kewensis</i>	106
44.	<i>Coryphantha deserti</i>	46	107.	<i>Neomammillaria subpolyedra</i>	106
45.	<i>Coryphantha deserti</i>	46	108.	<i>Neomammillaria tetracantha</i>	108
46.	Flower of <i>Coryphantha deserti</i>	46	109.	<i>Neomammillaria elegans</i>	108
47.	<i>Coryphantha aggregata</i>	47	110.	<i>Neomammillaria pseudoperbella</i>	109
48.	<i>Mammillaria recurispina</i>	47	111.	<i>Neomammillaria dealbata</i>	109
49.	Fruit of <i>Coryphantha sulcata</i>	49	112.	<i>Neomammillaria haageana</i>	110
50.	<i>Neobesseyia wissmannii</i>	52	113.	<i>Neomammillaria mundtii</i>	110
51.	<i>Escobaria tuberculosa</i>	54	114.	<i>Neomammillaria donatii</i>	111
52.	<i>Escobaria dasyacantha</i>	54	115.	<i>Neomammillaria collina</i>	111
53.	<i>Escobaria runyonii</i>	56	116.	<i>Neomammillaria celsiana</i>	112
54.	<i>Escobaria sneedii</i>	56	117.	<i>Neomammillaria aureiceps</i>	113
55.	<i>Bartschella schumannii</i>	58	118.	<i>Neomammillaria aureiceps</i>	113
56.	<i>Pelecyphora aselliformis</i>	58	119.	<i>Neomammillaria yucatanensis</i>	114
57.	Seed of <i>Phellosperma tetrancistra</i>	60	120.	<i>Neomammillaria ruestii</i>	114
58.	<i>Phellosperma tetrancistra</i>	61	121.	<i>Neomammillaria pringlei</i>	115
59.	<i>Dolichothele longimamma</i>	61	121a.	<i>Neomammillaria cerralboa</i>	116
60.	<i>Dolichothele sphaerica</i>	62	122.	<i>Neomammillaria phaeacantha</i>	116
61.	<i>Dolichothele longimamma</i>	62	123.	<i>Neomammillaria graessneriana</i>	116
62.	<i>Dolichothele uberiformis</i>	64	124.	<i>Neomammillaria spinosissima</i>	117
63.	<i>Solisia pectinata</i>	64	125.	<i>Neomammillaria spinosissima</i>	117
64.	<i>Neomammillaria mammillaris</i>	70	126.	<i>Neomammillaria densispina</i>	119
65.	<i>Neomammillaria macdougallii</i>	70	127.	<i>Neomammillaria nunezii</i>	119
66.	<i>Neomammillaria nivosa</i>	72	128.	<i>Neomammillaria nunezii</i>	120
67.	<i>Neomammillaria macdougallii</i>	75	129.	<i>Neomammillaria rhodantha</i>	120
68.	<i>Neomammillaria phymatothele</i>	75	130.	<i>Neomammillaria amoena</i>	121
69.	<i>Neomammillaria phymatothele</i>	77	131.	<i>Neomammillaria plumosa</i>	121
70.	<i>Neomammillaria phymatothele</i>	77	132.	<i>Neomammillaria prolifera</i>	125
71.	<i>Neomammillaria magnimamma</i>	78	133.	<i>Neomammillaria multiceps</i>	125
72.	<i>Neomammillaria macracantha</i>	78	134.	<i>Neomammillaria multiceps</i>	126
72a.	<i>Neomammillaria macracantha</i>	80	135.	<i>Neomammillaria campotricha</i>	127
72b.	<i>Neomammillaria johnstonii</i>	81	136.	<i>Neomammillaria schiedeana</i>	127
73.	<i>Neomammillaria melanocentra</i>	82	137.	<i>Neomammillaria lenta</i>	128
74.	<i>Neomammillaria seitziana</i>	82	138.	<i>Neomammillaria eriacantha</i>	128
75.	<i>Neomammillaria sartorii</i>	83	139.	<i>Neomammillaria denudata</i>	129
76.	<i>Neomammillaria ortegae</i>	84	140.	<i>Neomammillaria lenta</i>	129
77.	<i>Neomammillaria meiacantha</i>	85	141.	<i>Neomammillaria candida</i>	130
78.	<i>Neomammillaria scrippsiana</i>	85	142.	<i>Neomammillaria candida</i>	130
79.	<i>Neomammillaria sempervivi</i>	86	143.	<i>Neomammillaria vetula</i>	131
80.	<i>Neomammillaria polythele</i>	86	144.	<i>Neomammillaria discolor</i>	131
81.	<i>Neomammillaria carnea</i>	89	145.	<i>Neomammillaria fragilis</i>	133
82.	<i>Neomammillaria lloydii</i>	89	146.	<i>Neomammillaria elongata</i>	133
83.	<i>Neomammillaria zuccariniana</i>	90	147.	<i>Neomammillaria oliviae</i>	135
84.	<i>Neomammillaria compressa</i>	90	148.	<i>Neomammillaria echinaria</i>	136
85.	<i>Neomammillaria compressa</i>	91	149.	<i>Neomammillaria rekoi</i>	136
86.	<i>Neomammillaria compressa</i>	91	150.	<i>Neomammillaria pottsii</i>	137
87.	Spine-cluster of <i>N. compressa</i>	92	151.	<i>Neomammillaria mazatlanensis</i>	137
88.	<i>Neomammillaria compressa</i>	92	152.	<i>Neomammillaria albicans</i>	138
89.	<i>Neomammillaria mystax</i>	93	153.	<i>Neomammillaria slevinii</i>	139
90.	<i>Neomammillaria petterssonii</i>	93	154.	<i>Neomammillaria hamata</i>	141
91.	<i>Neomammillaria eichlamii</i>	95	155.	<i>Neomammillaria wildii</i>	141
92.	<i>Neomammillaria karwinskiiana</i>	95	155a.	<i>Neomammillaria rekoi</i>	142
93.	<i>Neomammillaria standleyi</i>	96	156.	<i>Neomammillaria solisii</i>	143
94.	<i>Neomammillaria parkinsonii</i>	96	157.	<i>Neomammillaria solisii</i>	143
95.	<i>Neomammillaria evermanniana</i>	97	157a.	<i>Neomammillaria seideliana</i>	144
96.	<i>Neomammillaria collinsii</i>	99	158.	<i>Neomammillaria seideliana</i>	144
97.	<i>Neomammillaria geminispina</i>	99	159.	<i>Neomammillaria barbata</i>	145
98.	<i>Neomammillaria woburnensis</i>	100	160.	<i>Neomammillaria mercadensis</i>	146
99.	<i>Neomammillaria chinocephala</i>	100	161.	<i>Neomammillaria multihamata</i>	146
100.	<i>Neomammillaria pyrrocephala</i>	103	162.	<i>Neomammillaria longicoma</i>	147
101.	<i>Neomammillaria polyedra</i>	103	163.	<i>Neomammillaria bocasana</i>	147
102.	<i>Neomammillaria tenampensis</i>	104	164.	<i>Neomammillaria multiformis</i>	148

TEXT-FIGURES—continued.

	PAGE.		PAGE.
FIG. 165. <i>Neomammillaria longicoma</i>	150	FIG. 214. <i>Pseudorhipsalis himantoclada</i>	213
166. <i>Neomammillaria glochidiata</i>	150	215. Top of flowering branch of <i>P. himantoclada</i>	214
167. <i>Neomammillaria trichacantha</i>	151	216. Longitudinal section of flower of <i>Pseudo-</i>	
168. <i>Neomammillaria saffordii</i>	151	<i>rhipsalis himantoclada</i>	214
169. <i>Neomammillaria painteri</i>	152	217. Section of flower of <i>Pseudorhipsalis alata</i>	214
170. <i>Neomammillaria microcarpa</i>	152	218. Flowering branch of <i>Pseudorhipsalis alata</i>	214
171. <i>Neomammillaria wrightii</i>	153	219. Unusual form of <i>Hatiora salicornioides</i>	217
172. <i>Neomammillaria mainae</i>	153	220. <i>Hatiora bambusoides</i>	218
172 <i>a</i> . <i>Neomammillaria boedekeriana</i>	154	221. <i>Rhipsalis cereuscula</i>	223
173. <i>Neomammillaria microcarpa</i>	155	222. <i>Rhipsalis cassutha</i>	226
174. <i>Neomammillaria microcarpa</i>	155	223. <i>Rhipsalis shaferi</i>	228
175. <i>Neomammillaria sheldonii</i>	157	224. <i>Rhipsalis lumbricoides</i>	230
176. <i>Neomammillaria carretii</i>	157	225. <i>Rhipsalis loefgrenii</i>	233
177. <i>Neomammillaria zephyranthoides</i>	159	226. <i>Rhipsalis sulcata</i>	235
178. <i>Neomammillaria bombycina</i>	160	227. <i>Rhipsalis gibberula</i>	235
179. <i>Neomammillaria occidentalis</i>	160	228. <i>Rhipsalis micrantha</i>	239
179 <i>a</i> . <i>Neomammillaria occidentalis</i>	161	229. Top of fruiting branch of <i>R. ramulosa</i>	241
180. <i>Neomammillaria fasciculata</i>	162	230. <i>Rhipsalis platycarpa</i>	242
181. <i>Neomammillaria longiflora</i>	162	231. <i>Rhipsalis crispimarginata</i>	244
182. Fruit, spine-cluster, and seed of <i>Neomam-</i>		232. <i>Rhipsalis crispata</i>	245
<i>millaria nelsonii</i>	163	233. <i>Rhipsalis cuneata</i>	246
183. Seed and spine-cluster of <i>N. longiflora</i>	163	233 <i>a</i> . <i>Neoabbottia paniculata</i>	248
184. <i>Neomammillaria xanthina</i>	164	234. Fruit, stem, and seeds of <i>Opuntia wet-</i>	
184 <i>a</i> . <i>Neomammillaria milleri</i>	176	<i>morei</i>	255
185. Flowering branch of <i>Zygocactus truncatus</i>	178	235. <i>Opuntia impedata</i>	257
186. Fruiting joint of <i>Zygocactus truncatus</i>	178	236. <i>Opuntia pisciformis</i>	258
187. <i>Zygocactus truncatus</i>	179	237. <i>Opuntia eburnispina</i>	260
188. <i>Epiphyllanthus obovatus</i>	181	238. <i>Opuntia macbridei</i>	261
189. <i>Epiphyllanthus obovatus</i>	181	239. <i>Cereus trigonodendron</i>	267
190. <i>Epiphyllanthus microsphaericus</i>	182	240. Flower of <i>Cephalocereus purpusii</i>	269
191. <i>Epiphyllanthus candidus</i>	182	241. Flower of <i>Cephalocereus purpusii</i>	269
192. <i>Schlumbergera gaertneri</i>	183	242. <i>Cephalocereus collinsii</i>	270
193. <i>Schlumbergera gaertneri</i>	184	243. <i>Pachycereus pringlei</i>	271
194. Top of fruiting branch of <i>Epiphyllum</i>		244. <i>Pachycereus lepidanthus</i>	272
<i>phyllanthus</i>	186	245. <i>Pachycereus lepidanthus</i>	272
195. Seedling of <i>Epiphyllum phyllanthus</i>	188	246. <i>Lemaireocereus eichlamii</i>	273
196. <i>Epiphyllum pumilum</i>	189	247. <i>Lemaireocereus beneckeii</i>	274
197. <i>Epiphyllum caudatum</i>	190	248. Branch, old flowers, and seeds of <i>Penio-</i>	
198. <i>Epiphyllum darrahii</i>	191	<i>cereus johnstonii</i>	275
199. <i>Epiphyllum crenatum</i>	193	249. <i>Dendrocereus nudiflorus</i>	276
200. <i>Epiphyllum macropterum</i>	194	250. <i>Borzicactus fieldianus</i>	277
201. <i>Epiphyllum guatemalense</i>	195	251. <i>Borzicactus fieldianus</i>	277
202. <i>Epiphyllum stenopetalum</i>	196	252. <i>Borzicactus fieldianus</i>	278
203. Tip of branch with flower of <i>Disocactus</i>		253. <i>Borzicactus fieldianus</i>	278
<i>biformis</i>	201	254. <i>Borzicactus fieldianus</i>	278
204. <i>Disocactus biformis</i>	201	255. <i>Binghamia multangularis</i>	279
205. <i>Disocactus eichlamii</i>	202	256. Flower of <i>Neoabbottia paniculata</i>	280
206. <i>Chiapasia nelsonii</i>	203	257. Fruit of <i>Neoabbottia paniculata</i>	280
207. <i>Nopalxochia phyllanthoides</i>	205	258. <i>Neoabbottia paniculata</i>	281
208. <i>Wittia amazonica</i>	206	259. <i>Neoabbottia paniculata</i>	282
209. <i>Wittia panamensis</i>	207	260. <i>Neoabbottia paniculata</i>	282
210. <i>Rhipsalidopsis rosea</i>	209	261. <i>Selenicereus nelsonii</i>	284
211. <i>Acanthorhipsalis micrantha</i>	210	262. <i>Cactus oaxacensis</i>	289
212. <i>Acanthorhipsalis crenata</i>	211	263. Hillside covered with giant cactus, <i>Car-</i>	
213. <i>Acanthorhipsalis monacantha</i>	212	<i>negia gigantea</i>	290

THE CACTACEAE

Descriptions and Illustrations of Plants of the Cactus
Family

DESCRIPTIONS AND ILLUSTRATIONS OF PLANTS OF THE CACTUS FAMILY.

Tribe 3. CEREEAE.

Subtribe 6. CORYPHANTHANA.

Terrestrial, spiny, low cacti, mostly globose, sometimes cylindric, rarely elongated, 1-jointed, solitary or cespitose, tuberculate, the tubercles numerous; tubercles usually arranged in spirals; juice watery or milky; flowers always solitary at areoles, either at top or side of plant, but never at spine-areoles, large or small, regular (except in the genus *Cochemia*): ovary naked or bearing a few scales; fruit a green or red indehiscent berry (except in the genus *Bartschella*); seeds small, brown or black.

We recognize 14 genera.

Key to Genera.

- A. Ovary more or less scaly (not known in *Mamillopsis*).
 - Flower campanulate with short tube.
 - Some of spines hooked 1. *Ancistrocactus* (p. 3)
 - None of spines hooked (see species No. 2 in *Neolloydia*).
 - Tubercles not deeply grooved; fruit scaly 2. *Thelocactus* (p. 6)
 - Tubercles deeply grooved; fruit nearly naked 3. *Neolloydia* (p. 14)
 - Flower-tube elongated, scaly 4. *Mamillopsis* (p. 19)
- AA. Ovary naked or nearly so.
 - B. Flowers irregular 5. *Cochemia* (p. 21)
 - BB. Flowers regular.
 - C. Flowers central, borne in axils of young, usually nascent, tubercles, large (except in genus No. 8); tubercles containing a watery Juice; fruit dull green or red; seeds brown or black.
 - D. Tubercles grooved on upper side; flowers borne at base of groove.
 - Seeds mostly light brown; Fruit greenish or yellowish even when mature, ripening slowly 6. *Coryphantha* (p. 23)
 - Seeds black to (dark brown; fruit red, maturing rapidly.
 - Tubercles long, not numerous, not persisting as woody knobs; aril of seed large 7. *Neobesseya* (p. 51)
 - Tubercles short, numerous, persisting after spines fall off as woody knobs; aril of seed small 8. *Escobaria* (p. 53)
 - DD. Tubercles not grooved above.
 - Fruit circumscissile; tubercles fleshy; spines acicular 9. *Bartschella* (p. 57)
 - Fruit not circumscissile; tubercles woody; spines peculate 10. *Pelecyphora* (p. 59)
 - CC. Flowers lateral, borne in axils of old and mature tubercle;; these never grooved above.
 - Seeds with a large corky aril 11. *Phellosperma* (p. 60)
 - Seeds without a corky aril.
 - Flowers large with an elongated tube; tubercles elongated, flabby 12. *Dolichothele* (p. 61)
 - Flowers small campanulate; tubercles not flabby.
 - Hilum of seed large; tubercles lactiferous; spines pectinate. 13. *Solisia* (p. 64)
 - Hilum of seed minute; tubercles sometimes lactiferous, but not in species with black seeds; spines not pectinate. 14. *Neomammillaria* (p. 65)

1. ANCISTROCACTUS gen. nov.

Small, globular or short-cylindric plants, indistinctly ribbed, strongly tubercled, very spiny, one of central spines always hooked; flowering tubercles more or less grooved on tipper side; flowers rather small, short, funnelform, borne at top of plant; ovary small, bearing a few thin scales, these always naked in their axils; fruit oblong, greenish, juicy, thin-walled, usually naked below but with a few broad cordate, thin-margined scales above; seeds globular, rather large, brownish to black, the papillae low, flattened; hilum large, depressed, sub-basal, surrounded by a thick rim.

Type species: *Echinocactus megarhizus* Rose.

Engelmann in describing *Echinocactus scheeri*, one of the species of this genus, refers to its anomalous characters when he says:

"Seeds are large, about 1 line long, 0.8 line in diameter, with very minute and flattened tubercles, brown (the only *Echinocactus* with seeds of that color known to me); hilum large and circular, surrounded by a thick rim; albumen very small; embryo curved but cotyledons small, connate, more like those of a *Mammillaria*, separating on the curvature and not at the end of the hook, as in all other hooked embryos of Cactaceae known to me." (Cact. Mex. Bound. 19. 1859.)

The generic name is from ἄγκιστρον fish-hook, and κάκτος cactus, referring to the long, hooked central spines.

Ancistrocactus was used by Schumann for a subgenus of *Echinocactus*. We recognize three species in the genus, occurring in southern Texas and northern Mexico.

Coulter (Contr. U. S. Nat. Herb. 3: 368, 369) calls attention to grooved areoles of *Echinocactus breviamatus* resembling those of *Coryphantha* and *Echinocactus scheeri*.

KEY TO SPECIES.

- Radial spines 20 or more, strongly appressed, pectinate; flowering areoles naked 1. *A. megarhizus*
 Radial spines 18 or fewer, more or less spreading, hardly pectinate; flowering areoles woolly.
 Groove half length of tubercle; flower greenish; radial spines 15 to 18 2. *A. scheeri*
 Groove extending full length of tubercle; flower rose-colored; radial spines usually 12 3. *A. breviamatus*

1. *Ancistrocactus megarhizus* (Rose).

Echinocactus megarhizus Rose, Contr. U. S. Nat. Herb. 12: 290. 1909.

Solitary or in clusters of 3 or 4; plant body nearly globular or a little elongated, 5 to 8 cm. high, usually solitary, from large and fleshy roots; ribs spiral, divided into dark-green tubercles, 4




FIG. 1.—*Ancistrocactus megarhizus*.

to 5 cm. high; radial spines 20 or more, pectinate, at first pale yellow, in age white; in seedlings the spines pubescent; central spines usually 4, the 3 upper similar to the radials, although a little stouter and in young areoles not easily distinguished from them, the lower central spines stout and strongly hooked, 15 mm. long; flowers not seen; fruit green, suggesting that of a *Coryphantha*, clavate, bearing a few naked scales near top; seed black, smooth, shining.

Type locality: Near Victoria, Mexico.

Distribution: Known only from the type locality.


Text-figure 1 is from a photograph of the type specimen collected by Dr. Edward Palmer.

2. *Ancistrocactus scheeri* (Salm-Dyck).

Echinocactus scheeri * Salm-Dyck, Cact. Hort. Dyck. 1849. 155. 1850.

Globular to clavate, 3.5 to 5 cm. long; ribs usually 13, indistinct, somewhat spiraled, strongly divided into stout, terete tubercles grooved only to middle; radial spines 15 to 18, spreading, 12

* This name was originally spelled *Echinocactus scheerii*.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Fruit of plant of *Coryphantha neo-mexicana*.
- 1a. Fruit of same.
2. Top of flowering plant of *Ancistrocactus scheeri*.
3. Flowering plant of *Cochemiea poselgeri*.

- 3a. Fruit of same.
- 3b. Seed of same.
4. Flowering plant of *Coryphantha cornifera*.

mm. long or less, white to straw-colored; central spines 3 or 4, the lowest one strongly hooked; flowers small, 2.5 cm. long, greenish yellow; ovary small, nearly naked; seeds large (about 2 mm, long), brown and minutely tuberculate (according to Coulter).

Type locality: Not cited.

Distribution: Southern Texas and northern Mexico.

It is probable that this species is based on Potts's specimen from Chihuahua and, if so, may be a different species from the one described by Engelmann, which he said was "a most elegant little species, one and a half to two inches high; larger spines black and white variegated." We have not seen Potts's plant, but it was referred here by Hemsley.

Illustrations: Cact. Mex. Bound. pl. 17; Rümpler, Sukkulente f. 105; Cact. Journ. 1: pl. for March; Schelle, Handb. Kakteenk. 156. f. 84, as *Echinocactus scheeri*.

Plate 11, figure 2, shows a plant collected by Dr. Rose at Laredo, Texas, in 1913, which flowered in the New York Botanical Garden in 1914. Text-figure 2 is from a photograph taken by Robert Runyon of a plant collected in 1921 near Brownsville, Texas.




FIG. 2.—*Ancistrocactus scheeri*.



FIG. 3.—*Ancistrocactus brevihamatus*.

3. *Ancistrocactus brevihamatus* (Engelmann).

Echinocactus brevihamatus Engelmann, Proc. Amer. Acad. 3: 271. 1856.

Echinocactus scheeri brevihamatus Weber in Schumann, Gesamtb. Kakteen 336. 1898.

Globose to obovoid, 5 to 10 cm. high, 5 to 7.5 cm. in diameter, dark green; ribs usually 13, compressed, strongly tubercled; tubercles grooved on upper side from spine-cluster to base, the groove woolly; radial spines 10 to 14, terete, white, 10 to 20 mm. long; central spines 4, the lower one porrect, hooked at apex; flowers rose-colored, 25 to 32 mm. long, not so broad as long; inner perianth-segments 15 mm. long, 4 mm. broad; mid-rib darker colored than margins; fruit about 1.5 cm. long, thin-walled, nearly naked; seeds brownish black, about 2 mm. long, smooth or with low flattened papillae, with a deep-set basal hilum.

Type locality: On the San Pedros, Texas.

Distribution: Southern Texas.

Illustrations: Haage and Schmidt, Haupt-Verz. Cact. 1908: 226; Schelle, Handb. Kakteenk. 157. f. 85, as *Echinocactus scheeri brevihamatus*; Cact. Mex. Bound. pl. 18, 19; Ann. Rep. Smiths. Inst. 1908: pl. 3, f. 3; Förster, Handb. Cact. ed. 2. 16. f. 64; Rümpler, Sukkulente 186. f. 104; Blanc, Cacti 41. No. 414; Engler and Prantl, Pflanzenfam. 3^{6a}: 162. f. 56, c, as *Echinocactus brevihamatus*.

Text-figure 3 is a reproduction of plate 18 of the Cactaceae of the Mexican Boundary Survey.

2. THELOCACTUS (Schumann) Britton and Rose, Bull. Torr. Club 49: 251. 1922.

Cacti of medium size, globular or somewhat depressed, spiny, often densely so; ribs few, low or even indefinite, divided into large, often spiraled, tubercles; flowering tubercles more or less grooved above; flowers from near center of plant, borne on very young tubercles, rather large for the subtribe, campanulate, diurnal; scales on ovary usually few, their axils naked; fruit so far as known dry, dehiscing by a basal pore; seeds black, finely tuberculate, with a large basal hilum.

Type species: *Echinocactus hexaedrophorus* Lemaire.

The generic name is from *οηλή* nipple, and cactus, referring to the tubercled ribs. *Thelocactus* was used for a subgenus of *Echinocactus* by Schumann; he described it with "ribs mostly divided into spirally disposed tubercles or mamillae, not protruding like a chin at base; spines straight or slightly curved." He referred to the group a number of diverse species representing several generic types, some of which we took up in Volume III.

We recognize 12 species, all native of Mexico.

To this genus we have referred the *Echinocacti* of previous authors which seem to intergrade with the *Coryphanthanae*. The group is perhaps complex and may contain two or more distinct genera, but most of the species are little known.

KEY TO SPECIES.

- Ribs indefinite, strongly tubercled.
 - Spines all straight.
 - Tubercles not flattened laterally; radial spines 6 to 9..... 1. *T. hexaedrophorus*
 - Tubercles flattened laterally; spines 1 to 5.
 - Flowers white..... 2. *T. rinconensis*
 - Flowers not white.
 - Flowers salmon to yellow..... 3. *T. lophothele*
 - Flowers rose-purple..... 4. *T. phymatothele*
 - Some of spines curved outward..... 5. *T. buekii*
 - Ribs definite, but more or less divided into tubercles.
 - Flowers yellowish.
 - Ribs 8 to 13..... 6. *T. leucacanthus*
 - Ribs 20 to 25..... 7. *T. nidulans*
 - Flowers red to purple.
 - Spines all straight.
 - Spines subulate..... 8. *T. fossulatus*
 - Spines acicular..... 9. *T. tulensis*
 - Spines more or less curved.
 - Spines 8 or fewer..... 10. *T. lloydii*
 - Spines numerous.
 - Central spines flexible, usually straight, porrect or ascending..... 11. *T. bicolor*
 - Central spines subulate, rigid, some of them curved and reflexed..... 12. *T. pottsii*

1. *Thelocactus hexaedrophorus* (Lemaire) Britton and Rose, Bull. Torr. Club 49: 251. 1922.

Echinocactus hexaedrophorus Lemaire, Cact. Gen. Nov. Sp. 27. 1839.

Echinocactus hexaedrophorus roseus Lemaire in Labouret, Monogr. Cact. 251. 1853.

Echinocactus hexaedrophorus labouretianus Schumann, Gesamtb. Kakteen 438. 1898.

Echinocactus hexaedrophorus major Quehl in Schumann, Gesamtb. Kakteen 438. 1898.

Globose or somewhat flattened above or umbilicate, glaucous, strongly tubercled, not ribbed, 13 to 14 cm. in diameter; tubercles prominent, somewhat 6-sided, 27 mm. broad at base, arranged in indefinite spirals; radial spines 6 to 9, spreading, unequal, 11 to 18 mm. long, rigid, straight, subulate, annulate; central spine much stouter than the radials, erect, 2.3 to 3 cm. long; flowers large, 5.5 cm. long and broader than long when expanded; perianth-segments oblong, purplish; stigma-lobes yellowish white.

Type locality: Tampico, Mexico.

Distribution: Central Mexico.

Schumann refers a plant from San Luis Potosí* to this species. The type, however, is said to have come from Tampico on the coast, while San Luis Potosí is on the table-land at an altitude of 7,000 feet or more, and such an altitudinal distribution is not to be expected. It is possible, but hardly probable, that the plant was actually collected at San Luis Potosí but shipped from Tampico, the port of San Luis Potosí, as such mistakes were common in the early shipments of cacti. Thus, species are attributed to Buenos Aires which came from

* This plant of the table-land is *Echinocactus fossulatus* Scheidweiler.

northwestern Argentina, and *Echinocactus insculptus*, referred to below, although reported from Buenos Aires, is really of Mexican origin.

Echinocactus insculptus Scheidweiler (Hort. Belge 4: 120. pl. 7. 1837) is referred here by Schumann, but the illustration indicates a very different plant.

Echinocactus labouretianus, referred by Schumann (Gesamtb. Kakteen 438. 1898) to Cels's Catalogue, probably never described, is to be referred here.

Illustrations: Cact. Journ. 1: 181; Lemaire, Icon. Cact. pl. 4; Dict. Gard. Nicholson 1: f. 690; Balt. Cact. Journ. 2: 196; Rümpler, Sukkulente 182. f. 101; Knippel, Kakteen pl. 12; Amer. Gard. 11: 461; Blanc, Cacti 45. No. 508; Schumann, Gesamtb. Kakteen 437. f. 76; Watson, Cact. Cult. 105. f. 36; ed. 3. f. 25, as *Echinocactus hexaedrophorus*.




FIG. 4.—*Thelocactus rinconensis*.




FIG. 5.—*Thelocactus phymatothele*.

2. *Thelocactus rinconensis* (Poselger).

Echinocactus rinconensis * Poselger, Allg. Gartenz. 23: 18. 1855.

Simple, globose or somewhat depressed, 6 to 8 cm. high, 12 cm. in diameter; ribs somewhat spiraled, strongly tubercled; tubercles more or less flattened laterally, somewhat angled; spines usually only 3, acicular, 1.5 cm. long; flowers white, 4 cm. long; inner perianth-segments lanceolate, acute.

Type locality: Near Rinconada, Mexico.

Distribution: Nuevo Leon, Mexico.

We do not know this species definitely, but we suspect that the plant collected and illustrated by Safford as *Echinocactus lophothele* belongs here.

Illustrations: Schumann, Gesamtb. Kakteen 433. f. 75; Schelle, Handb. Kakteenk. 197. f. 130, as *Echinocactus rinconadensis*; (?) Ann. Rep. Smiths. Inst. 1908: pl. 3, f. 1, as *Echinocactus lophothele*.

Text-figure 4 is reproduced from the first illustration cited above.

3. *Thelocactus lophothele* (Salm-Dyck) Britton and Rose, Bull. Torr. Club 49: 251. 1922.

Echinocactus lophothele Salm-Dyck, Allg. Gartenz. 18: 395. 1850.

Simple, or in its native state cespitose, globose, sometimes depressed or short-cylindric, up to 25 cm. high, glaucous; ribs indefinite, strongly tuberculate; tubercles flattened; areoles depressed, gray-

* Because this species came from Rinconada, Schumann (Engler and Prantl, Pflanzenfam. 3^{6a}: 189. 1894) has changed the name to *Echinocactus rinconadensis*.

ish lanate when young; radial spines 3 to 5, stout, purplish brown, 1 to 3 cm. long; central spines wanting or solitary; flowers salmon to yellow, about 5 cm. broad; perianth-segments nearly linear, acute; scales of ovary glabrous, 6 mm. long.

Type locality: Near Chihuahua.

Distribution: Chihuahua, Mexico.

Our description is drawn mostly from the figure in Blühende Kakteen, plate 126.

We have seen flowering specimens of what is called this species at La Mortola, Italy. Although the type came from Chihuahua, we have seen no plant from that region which answers it.

There is a plant in collections, passing as *Echinocactus lophothele longispinus* (Monatsschr. Kakteenk. 15: 138. 1905), which we do not know.

Illustrations: Schelle, Handb. Kakteenk. 196. f. 129; Blühende Kakteen 3: pl. 126; Weinberg, Cacti 12; Blanc, Cacti 48. No. 560, as *Echinocactus lophothele*.

4. *Thelocactus phymatothele* (Poselger).

Echinocactus phymatothelos * Poselger in Förster, Handb. Cact. ed. 2. 602. 1885.

Simple, depressed-globose, 5 cm. high, 9 to 10 cm. in diameter; ribs 13, glaucous-green, divided into low irregular tubercles, these somewhat flattened and pointed; spines usually 1 to 3, sometimes wanting, subulate, rigid, 2 cm. long, brown, spreading; flowers 6 cm. broad; inner perianth-segments rose-purple to pinkish, narrow, acute; scales on ovary and flower-tube acute.

Type locality: Not cited.

Distribution: Mexico.

This plant is evidently related to *Thelocactus lophothele*.

Illustrations: Möllers Deutsche Gärt. Zeit. 25: 474. f. 6, No. 24; Blühende Kakteen 3: pl. 130, as *Echinocactus phymatothelos*.

Text-figure 5 is reproduced from the second illustration above cited.

5. *Thelocactus buekii* (Klein).

Echinocactus buekii † Klein, Gartenflora 8: 257. 1859.

Stems simple, deep green; tubercles distinct, somewhat pointed, angled; spines about 7, reddish, unequal, some of them outwardly curved, the longer ones much elongated; flowers dark red; inner perianth-segments narrow.

Type locality: Mexico.

Distribution: Known only from the type locality.

Schumann refers this species to *Echinocactus tulensis*, but it is clearly different from his illustration of that species. Its relationship must be rather with *Thelocactus rinconensis* (see Schumann's figure, No. 75).

This plant is probably named for Dr. Johannes Nicolaus Buck, a botanist and physician of Frankfurt, Germany, and author of the Index to De Candolle's Prodrum.

Illustration: Gartenflora 8: pl. 266, as *Echinocactus buekii*.

Text-figure 6 is reproduced from the illustration cited above.

6. *Thelocactus leucacanthus* (Zuccarini).

Echinocactus leucacanthus Zuccarini in Pfeiffer, Enum. Cact. 66. 1837.

Cereus tuberosus Pfeiffer, Enum. Cact. 102. 1837.

Cereus maelenii Pfeiffer, Allg. Gartenz. 5: 378. 1837.

Echinocactus porrectus Lemaire, Cact. Aliq. Nov. 17. 1838.

Echinocactus subporrectus Lemaire, Cact. Aliq. Nov. 25. 1838.

Echinocactus maelenii Salm-Dyck, Cact. Hort. Dyck. 1842. 18. 1843.

Mammillaria maelenii Salm-Dyck, Cact. Hort. Dyck. 1844. 14. 1845.

Echinocactus leucacanthus Tuberosus Förster, Handb. Cact. 287. 1846.

Echinocactus leucacanthus crassior Salm-Dyck, Cact. fort. Dyck. 1849. 35. 1850.

Echinocactus theloideus Salm-Dyck, Allg. Gartenz. 18: 396. 1850.

* This is the original spelling of the name, but it is sometimes written *Echinocactus phymatothele*, the ending being the usual one for specific names of this kind.

† The original spelling of this name was *buckii*, but on the accompanying plate it was *buekii*.

‡ This name is spelled *macleanii* by Hemsley (Biol. Centr. Amer. Bot. 1: 534. 1880).

Densely cespitose, short-cylindric, 10 to 15 cm. long; ribs 8 to 13, sometimes spiraled, obtuse, tubercled; radial spines 7 to 20, at first light yellow, in age gray, spreading or recurved, unequal, the longer ones 4 cm. long, more or less annulate; central spine solitary, at first blackish, but in age gray, up to 5 cm. long; flowers yellow, 5 cm. long; inner perianth-segments numerous, lanceolate, acute; ovary and flower-tube bearing broad imbricated scales.

Type locality: Near Zimapán, Mexico.

Distribution: Zimapán and Ixmiquilpan, Mexico.

We are inclined to refer here *Echinocactus ehrenbergii* Pfeiffer (Allg. Gartenz. 6: 275. 1838), which, according to Schumann, also came from Ixmiquilpan, Mexico. In his monograph Schumann describes the flowers as yellow like those of *E. leucacanthus*, but in his English Keys he says that the flowers are rose-red. Dr. Rose, who collected in this region in 1905, found only one species of this relationship.




FIG. 6.—*Thelocactus buekii*.




FIG. 7.—*Thelocactus leucacanthus*.

Echinocactus tuberosus Salm-Dyck (Förster, Handb. Cact. 287. 1846) is known only as a synonym.

Echinocactus tuberosus subporrectus (Förster, Handb. Cact. 523. 1846) belongs here.

Illustrations: Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 14; Abh. Bayer. Akad. Wiss. München 2: pl. 2, f. 10; pl. 3, f. 4, as *Echinocactus leucacanthus*.

Figure 7 is from a photograph of the plant collected by Dr. Rose at Ixmiquilpan in 1905.

7. *Thelocactus nidulans* (Quehl).

Echinocactus nidulans Quehl, Monatsschr. Kakteenk. 22: 119. 1911.

Simple, depressed-globose, 10 cm. high, sometimes 20 cm. in diameter, gray, usually glaucous; ribs 20 to 25, rather indistinct, divided into tubercles; spines about 15, all similar, 2 to 6 cm. long; flowers 4 cm. long, yellowish white.

Type locality: Mexico.

Distribution: Mexico, but known only from cultivated plants.

Illustrations: Monatsschr. Kakteenk. 22: 51; Alianza Científica Universal 3: 114, as *Echinocactus nidulans*.

Figure 8 is from a photograph given to Dr. Rose by Frantz de Laet in 1912.

8. *Thelocactus fossulatus* (Scheidweiler).

Echinocactus fossulatus Scheidweiler, Allg. Gartenz. 9: 49. 1841.

Echinocactus hexaedrophorus subcostatus Salm-Dyck, Cact. Hort. Dyck. 1849. 34. 1850.

Echinocactus hexaedrophorus fossulatus Salm-Dyck in Labouret, Monogr. Cact. 251. 1853.

Globose to much depressed, 10 to 15 cm. in diameter; ribs usually 13, slightly glaucous, bronzed; tubercles large, somewhat flabby, more or less compressed, dorsally somewhat angled; flowering areoles narrow, sometimes extending forward to next tubercle; radial spines 4 or 5, unequal, 1 to 3.5 cm. long, brown; central spine solitary, 3 to 4.5 cm. long, subulate, annulate; flowers nearly white or slightly tinged with pink; scales on flower-tube ovate, their scarious margins slightly ciliate.

Type locality: Near San Luis Potosí, Mexico.

Distribution: San Luis Potosí, Mexico.




Fig. 8.—*Thelocactus nidulans*.




Fig. 9.—*Thelocactus fossulatus*.

Somewhat similar to the foregoing species is C. A. Purpus's No. 15 from Minas de San Rafael, Mexico. This plant has more rounded tubercles, only 4 spines, these all radial and 2 cm. long or less, somewhat flattened.

Thelocactus fossulatus is certainly distinct from *Thelocactus hexaedrophorus*, differing in the arrangement of the tubercles and in the color of the flowers. The former is from an altitude of 7,000 feet, while the other is from near sea-level.

Echinocactus drageanus (Moerder, Rev. Hort. 67: 186. 1895) and *E. droegeanus* Hildmann (Schumann, Gesamtb. Kakteen 438. 1898) probably belong here, although the latter is referred by Schumann to *Echinocactus hexaedrophorus*. This may be the plant, judging from the name and authorities mentioned, which Schelle (Handb. Kakteenk. 257. 1907) refers to as *Mammillaria rhodantha droegeana* Schumann (*M. droegeana* Hildmann). Schelle questions whether it may not be a distinct species, presumably a *Mammillaria*.

Illustrations: Scientific Amer. 124: 492, as *Echinocactus*; Curtis's Bot. Mag. 73: pl. 4311; Ann. Rep. Smiths. Inst. 1908: pl. 13, f. 3, as *Echinocactus hexaedrophorus*; Pfeiffer, Abbild. Besch. Cact. 2: pl. 13, as *Echinocactus fossulatus*; Monatsschr. Kakteenk. 27: 41, as *Echinocactus hexaedrophorus droegeanus*.

Figure 9 is from a photograph of a plant collected by Dr. Edward Palmer at San Luis Potosí, Mexico, in 1905.

9. *Thelocactus tulensis* (Poselger).*Echinocactus tulensis* Poselger, Allg. Gartenz. 21: 125. 1853.

Plant simple to abundantly cespitose, globular to short-cylindric, up to 25 cm. high; ribs 8 to 13, strongly tubercled; radial spines 6 to 8, more or less spreading, 10 to 15 mm. long, brownish; central spines solitary or sometimes 2, 3 cm. long; flowers 2.5 cm. long, rose-colored; inner perianth-segments linear-oblong, acute.

Type locality: Near Tula, Tamaulipas, Mexico.

Distribution: Tamaulipas, Mexico.

We have not seen this plant but we have seen two good illustrations. It is closely related to *Thelocactus hexaedrophorus*.

Illustrations: Blühende Kakteen 1: pl. 18; Schumann, Gesamtb. Kakteen 431. f. 74, as *Echinocactus tulensis*.

Figure 10 is reproduced from the first illustration cited above.




FIG. 10.—*Thelocactus tulensis*.




FIG. 11.—*Thelocactus bicolor*.

10. *Thelocactus lloydii* sp. nov.

Plants simple, depressed-globose, 8 to 12 cm. broad, pale bluish green, strongly tubercled and strongly armed; tubercles conspicuous but low, often wider than long, sometimes 4 cm. wide; flowering groove rather conspicuous but narrow, extending from spines to about half-way to axil of tubercle; spines usually 8, sometimes with a smaller accessory one, all ascending from base and curved outward from middle, terete or somewhat angled at base, often highly colored below with sharp yellowish-crimson tips, the longer ones 6 cm. long; outer perianth-segments very pale purple, never deep purplish pink; filaments white; anthers deep yellow; style yellowish, pinkish at top; stigma-lobes pinkish yellow.

Collected by F. E. Lloyd in northern Zacatecas, Mexico, May 25, 1908 (No. 33).

11. *Thelocactus bicolor* (Galeotti) Britton and Rose, Bull. Torr. Club 49: 251. 1922.

Echinocactus bicolor Galeotti in Pfeiffer, Abbild. Beschr. Cact. 2: pl. 25. 1848.

Echinocactus rhodophthalmus Hooker in Curtis's Bot. Mag. 76: pl. 4486. 1850.

Echinocactus rhodophthalmus ellipticus Hooker in Curtis's Bot. Mag. 78: pl. 4634. 1852.

Echinocactus ellipticus Lemaire, Jard. Fleur. 3: pl. 270. 1853.

Echinocactus bicolor schottii Engelm., Proc. Amer. Acad. 3: 277. 1856.

Echinocactus bolansis Runge, Gartenflora 38: 106. 1889.

Echinocactus bicolor bolansis Schumann, Gesamtb. Kakteen 303. 1898.

Echinocactus bicolor tricolor Schumann, Gesamtb. Kakteen 303. 1898.

Echinocactus schottii Small, Fl. Southeast. U. S. 814. 1903.

Plants simple, globose to conic, glaucous, small, up to 3 cm. high, very spiny; ribs usually 8, broad, somewhat tubercled; areoles approximate; spines highly colored, sometimes bright red or yellowish or red and yellow; radial spines 9 to 18, widely spreading or sometimes bent backward at tip, 3 cm. long or less; central spines usually 4, ascending or porrect, all straight, 3 to 5 cm. long, subulate; flowers large, 5 to 6 cm. long and fully as broad when expanded; outer perianth-segments pale purple; inner perianth-segments deep purplish pink, oblong, acute; scales on ovary and flower-tube imbricated, ovate, with scarious and ciliate margins; filaments white to purple; stigma-lobes pale to pinkish yellow; fruit small, about 1 cm. long, dehiscing by a large irregular basal opening; seeds 2 mm. long, black, broader at apex, tuberculate with a circular and depressed basal hilum.

Type locality: Mexico.

Distribution: Southern Texas to central Mexico.

Echinocactus tricolor, *E. castaniensis*, and *E. bicolor montemorelanus* Weber (all in Dict. Hort. Bois 465. 1896) are usually referred here but were never described.

Illustrations: Jard. Fleur. 3: pl. 270, as *Echinocactus ellipticus*; Gartenflora 38: 106. f. 21, as *Echinocactus bolansis*; Curtis's Bot. Mag. 76: pl. 4486; Jard. Fleur. 1: pl. 101; Loudon, Encycl. Pl. ed. 3. 1377. f. 19375; Gard. Mag. Bot. 1: 40, as *E. rhodophthalmus*; Curtis's Bot. Mag. 78: pl. 4634, as *E. rhodophthalmus ellipticus*; Karsten and Schenck, Vegetationsbilder 2: pl. 20, c; Pfeiffer, Abbild. Beschr. Cact. 2: pl. 25; Schumann, Gesamtb. Kakteen Nachtr. 87. f. 14; Ann. Rep. Smiths. Inst. 1908: pl. 13, f. 2; Blühende Kakteen 2: pl. 74; Monatsschr. Kakteenk. 12: 7; 29: 81; Schelle, Handb. Kakteenk. 157. f. 86; Blanc, Cacti 41. No. 412, as *E. bicolor*.

Figure 11 is from a photograph taken by Robert Runyon at Saltillo, Mexico, in 1921.

12. *Thelocactus pottsii* (Salm-Dyck).

Echinocactus pottsii * Salm-Dyck, Allg. Gartenz. 18: 35. 1850.

Echinocactus bicolor pottsii Salm-Dyck, Cact. Hort. Dyck. 1849. 173. 1850.

Echinocactus heterochromus Weber, Dict. Hort. Bois 466. 1896.

Globular or somewhat depressed, 10 to 15 cm. in diameter, somewhat glaucous, yellowish; ribs 8 or 9, broad and obtuse, more or less distinctly tubercled; areoles large, closely set on old plants, densely felted when young, naked in age; spines variable as to number, shape, size, and color; radial spines 7 to 10, acicular, usually terete, straight or incurved, more or less banded with red and white or pale yellow, 1 to 3 cm. long; central spines several, stout-subulate, more or less flattened, 3 or 4 cm. long, often white, but sometimes banded with red; flowers, 5 to 6 cm. long; scales on ovary and flower-tube ovate, greenish; margins thin and ciliate; inner perianth-segments light purple, darker at base, oblong; stigma-lobes yellow; fruit globose, small, 1.5 cm. in diameter; seed tuberculate, black, truncate at base, ridged on back; hilum basal, white, circular.

Type locality: Near Chihuahua City.

Distribution: Chihuahua to Coahuila, Mexico.

There are three illustrations passing as *Echinocactus pottsii*, none of which agrees with the original description of Salm-Dyck. Two of these are in Nicholson's Dictionary (Dict. Gard. 4: 540. f. 23 and Suppl. f. 359) where the species is described as follows: flowers yellow, about 2 inches across, short-tubed, several expanding together at the top of the stem; stem globular, 1½ feet in diameter: ridges about a dozen, rounded and even, with acute sinuses; spines 1 inch long, bristle-like, arranged in clusters of 7 or 9, with a cushion of white wool at the base.

Nicholson indicates that his plant of *E. pottsii* was from California and introduced into cultivation in 1840. There is no Californian species which answers this description or illustration.

The other illustration is Schumann's (Gesamtb. Kakteen 328. f. 7), which is somewhat similar to the above. Schumann states that the radial spines are commonly 6, spreading and yellow; central spines solitary. We are not able to identify this illustration; it suggests some *Echinocereus* as much as it does an *Echinocactus*.

* Salm-Dyck (Cact. Hort. Dyck. 1849. 35. 1850) credits this name to Scheer.

Illustrations: Knippel, Kakteen pl. 7, in part; Schelle, Handb. Kakteenk. 144. f. 70, as *Echinocactus heterochromus*; Dict. Gard. Nicholson 4: 540. f. 23; Suppl. 336. f. 359; Schumann, Gesamtb. Kakteen 328. f. 57; Garden 2: 521; Monatsschr. Kakteenk. 30: 53; Schelle, Handb. Kakteenk. 155. f. 82; Watson, Cact. Cult. 117. f. 43, as *Echinocactus pottsii*.

PUBLISHED SPECIES, POSSIBLY OF THIS RELATIONSHIP.

ECHINOCACTUS CONOTHELOS Regel and Klein, Ind. Sem. Hort. Petrop. 1860: 48. 1860.

Ovoid to subcylindric, 10 cm. high, 7.5 cm. in diameter, grayish green; ribs somewhat spiraled, somewhat tubercled at base, the lower tubercles 12 to 20 mm. long; upper areoles oblique, white-tomentose; radial spines 14 to 16, white, spreading to recurved, 8 to 10 mm. long; central spines 2 to 4, erect or a little spreading and recurved, stouter and longer than the radials, 13 to 34 mm. long; flowers and fruit unknown.

Type locality: Near Tanquicillos and Jaumave, Mexico.

This plant was collected by Karwinsky and is known only from his collection. The authors refer the species to Salm-Dyck's section of the *Theloidei*, which, however, is a very diverse group containing representatives of several genera. Schumann was unable to place the species; it may be related to some species of *Thelocactus*.

ECHINOCACTUS HEXAEDRUS Scheidweiler, Bull. Acad. Sci. Brux. 6: 89. 1839.

Globose to oblong-ovate, glaucous; ribs 18, tuberculate; tubercles 6-angled, gibbous below areoles; areoles oblong, lanate; spines 13, white with purplish bases; lowermost spine longest; central spines 2, either straight or recurved; flowers and fruit unknown.

Type locality: San Luis Potosí.

ECHINOCACTUS SAUSSIERI Weber, Dict. Hort. Bois 468. 1896.

Depressed-globose, 15 to 20 cm. in diameter; ribs spiraled, strongly tubercled; radial spines 9, grayish white, 15 mm. long; central spines 4, acicular, 3 to 4 cm. long; flowers purplish, 4 cm. in diameter; inner perianth-segments lanceolate; stamens and style yellow.

Type locality: Matehuala, state of San Luis Potosí, Mexico.

We know this species from the brief description only and are unable to determine its relationship.

ECHINOCACTUS SMITHII Mühlenpfordt in Otto and Dietrich, Allg. Gartenz. 14: 370. 1846.

Simple, globose to cylindric, 7 cm. in diameter; ribs 21, often spiraled, strongly tubercled, glaucous; radial spines 20 to 27, setaceous, white, 16 mm. long; central spines, the upper one flattened, white with brown or black tips; flowers reddish, 3.5 cm. long; fruit globular, 8 mm. in diameter; seed nearly globular, flattened at the hilum.

Type locality: San Luis Potosí.

We know this species from description only and are unable to assign it to any genus.

ECHINOCACTUS VARGASII Regel and Klein, Ind. Sem. Hort. Petrop. 1860: 48. 1860.

Globose, 5 cm. high, 6 cm. in diameter; tubercles rather large, somewhat angled, arranged in spirals; radial spines 5 or 6, terete, subulate, brownish, 2 to 6 mm. long; central spine 1, erect, 12 mm. long; flowers and fruit unknown.

Type locality: Mexico, near Rio Blanco.

Schumann did not know this plant nor do we, but to us it suggests a *Thelocactus*. The authors of the species compared it with *Echinocactus poselgerianus*, now referred to *Coryphantha*, and with *E. phymatothelos*.

3. NEOLLOYDIA Britton and Rose, Bull. Torr. Club 49: 251. 1922.

Small, more or less cespitose cacti, fibrous-rooted, cylindric, densely spiny, tubercled; tubercles more or less arranged on spiraled ribs, grooved above; radial spines numerous, widely spreading; central spines one to several, much stouter and longer than radials; flowers large, pink or purple, subcentral from axils of nascent tubercles, their segments widely spreading; fruit compressed-globose, dull-colored, thin-walled, becoming papery, dry, with few scales or none; seeds globose, black, dull, tuberculate-roughened, with a large white basal scar; embryo straight in typical species.

Type species: *Mammillaria conoidea* De Candolle.

We recognize 7 species from central and northern Mexico and Texas, which have been transferred from *Echinocactus* and *Mammillaria*. The genus is dedicated to Professor Francis E. Lloyd, whose collections and observations have contributed highly important information to our investigations.

KEY TO SPECIES.

- Plants 3 cm. in diameter or less; central spines sometimes wanting 1. *N. pilispina*
- Plants larger; central spines always present.
 - Central spines curved or hooked 2. *N. clavata*
 - Central spines all straight.
 - Central spine solitary.
 - Central spine stiff, porrect 3. *N. horripila*
 - Central spine weak, ascending or connivent 4. *N. beguinii*
 - Central spines several.
 - Spines white or sometimes dark above 5. *N. ceratites*
 - Central spines or some of them black.
 - Radial spines 25 or more; Mexican species 6. *N. conoidea*
 - Radial spines 15 or less 7. *N. texensis*

1. *Neolloydia pilispina* (J. A. Purpus).

Mammillaria pilispina J. A. Purpus, Monatsschr. Kakteenk. 22: 150. 1912.

Plants cespitose, about 3 cm. in diameter; ribs indistinct, made up of very definite, somewhat angled tubercles young spine-areoles clothed with abundant, long, white wool covering top of plant;




FIG. 12.—*Neolloydia pilispina*.




FIG. 13.—*Neolloydia horripila*.

radial spines 6 or 7, 5 to 6 mm. long, weak and spreading, the upper ones longer and connivent over top of plant, 2 cm. long or more, white with blackish tips; central spines often wanting, sometimes one; flowers small, 1.5 to 2 cm. long, purplish; outer perianth-segments brownish.

Type locality: Minas de San Rafael, San Luis Potosí, Mexico.



Distribution. Known only from the type locality.

Figure 12 is from a photograph of a plant collected by C. A. Purpus at the type locality.

2. *Neolloydia clavata* (Scheidweiler).

- Mammillaria clavata* Scheidweiler, Bull. Acad. Sci. Brux. 5: 494. 1838.
Mammillaria stipitata Scheidweiler, Bull. Acad. Sci. Brux. 5: 495. 1838.
Mammillaria rhabdicantha Lemaire, Cact. Gen. Nov. Sp. 34. 1839.
Mammillaria ancistracantha Lemaire, Cact. Gen. Nov. Sp. 36. 1839.
*Mammillaria rhabdicantha humilior** Salm-Dyck in Förster, Handb. Cact. 244. 1846.
Mammillaria scolymoides rhabdicantha Salm-Dyck, Cact. Hort. Dyck. 1849. 132. 1850.
Echinocactus corniferus rhabdicanthus Poselger, Allg. Gartenz. 21: 102. 1853.
 ? *Mammillaria potosiana* Jacobi, Allg. Gartenz. 24: 92. 1856.
Mammillaria sulcoglandulifera Jacobi, Allg. Gartenz. 24: 92. 1856.
Coryphantha rhabdicantha Lemaire, Cactées 34. 1868.
Coryphantha ancistracantha Lemaire, Cactées 34. 1868.
Cactus ancistracanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus rhabdicanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus brunneus Coulter, Contr. U. S. Nat. Herb. 3: 117. 1894.
Cactus maculatus Coulter, Contr. U. S. Nat. Herb. 3: 117. 1894.
Mammillaria rhabdicantha† *ancistracantha* Schumann, Gesamtb. Kakteen 506. 1898.
Mammillaria radicansissima Quehl, Monatsschr. Kakteenk. 22: 164. 1912.

Plants simple, elongated, cylindric, 10 to 15 cm. high, dark bluish green; tubercles in rows of 5, 8, and 13, conic, grooved above, the axils when young bearing short white wool; glands in the groove 1 to several, large, red; radial spines 6 to 12, with reddish or black tips; central spine 1, somewhat longer than radials, curved or even hooked; flowers small for the genus, about 2 cm. long; outer perianth-segments linear, acute, entire, with broad brownish midrib; inner perianth-segments linear, entire, narrow, creamy white; stamens pinkish, much shorter than the perianth-segments; style pinkish; stigma-lobes 5 or 6, short, greenish.

FIG. 14.—*Neolloydia clavata*.FIG. 15.—*Neolloydia conoidea*.

Type locality: Not cited.

Distribution: San Luis Potosí, Mexico.

The two species of Coulter, *Cactus brunneus* and *Cactus maculatus*, as well as *Mammillaria radicansissima*, came from San Luis Potosí, and all seem to be so much alike that we do not hesitate to reduce them as above.

Echinocactus rhabdicanthus is credited by Schumann to Poselger, but he used the name *rhabdicanthus* only as a variety of *E. corniferus*. This binomial was used in 1850 by Salm-Dyck for a very different plant.

Mammillaria humilior Förster we have seen only in Schumann's Index (Gesamtb. Kakteen 824. 1898). He refers it to *M. rhabdicantha ancistracantha*.

* Schumann (Gesamtb. Kakteen 506, 824, Index, 1898), perhaps not intentionally, gives this name specific rank.

† Schumann has dropped the first "h" in *Mammillaria rhabdicantha* and he is followed by the Monatsschrift für Kakteenkunde.

Illustrations: Blühende Kakteen 1: pl. 7; Schumann, Gesamtb. Kakteen 505. f. 83, as *Mammillaria raphidacantha*; Blühende Kakteen 3: pl. 163; Monatsschr. Kakteenk. 22: 165, as *M. radicanissima*.

Figure 14 is from a photograph of a plant collected by Dr. Edward Palmer at San Luis Potosí, Mexico, in 1908 (No. 814).

3. *Neolloydia horripila* (Lemaire).

Mammillaria horripila Lemaire, Cact. Aliq. Nov. 7. 1838.

Echinocactus horripilus Lemaire, Cact. Gen. Nov. Sp. 91. 1839.

Echinocactus horripilus longispinus Monville in Labouret, Monogr. Cact. 265. 1853.

Simple or somewhat cespitose, globular to short-cylindric, 10 to 12 cm. high; tubercles glaucous, prominent, rounded at apex; radial spines 8 to 10, acicular, spreading, 15 mm. long, grayish; central spine solitary, straight, a little longer than the radials; flowers deep purple, 3 cm. long; inner perianth-segments narrowly oblong, acute; stigma-lobes, white.

Type locality: Not cited.

Distribution: Hidalgo, Mexico.

Lemaire first referred this plant to *Mammillaria*, but finally described it as an *Echinocactus* on account of its grooved tubercles; he believed that it was an intergrade between these two genera. As he states, its general appearance is that of a species of the so-called *Mammillaria*.

Echinocactus caespititius Pfeiffer is usually given as a synonym of this species, but it seems never to have been described. Schumann cites the place of publication as Salm-Dyck's Cactaceae of 1850 (p. 35), but it is given only as a synonym. It appeared also in Salm-Dyck's Cactaceae of 1845 (p. 17) and in Förster's Handbuch (p. 283), but also as a synonym.

Illustration: Blühende Kakteen 1: pl. 6, as *Echinocactus horripilus*.

Figure 13 is reproduced from the illustration above cited.

4. *Neolloydia beguinii* (Weber) Britton and Rose, Bull. Torr. Club 49: 252. 1922.

Echinocactus beguinii Weber in Schumann, Gesamtb. Kakteen 442. 1898.

Plant-body cylindric, 10 to 15 cm. high; ribs spiraled and divided at regular intervals into low tubercles resembling geometric figures, pale bluish green in color but nearly hidden by the dense covering of spines; radial spines 20 or more, white, but with dark tips; centrals usually single, longer and ascending; flowers appearing from top of plant, large, 3 to 4 cm. long, bright pink; stigma-lobes 7, long, white; ovary without scales; seeds black, tubercled, with a broad triangular hilum.

Type locality: Probably at Saltillo, in Coahuila, Mexico.

Distribution: Zacatecas and Coahuila, Mexico.

This plant is very distinct from *Echinomastus erectocentrus*, with which it was confused both by Coulter and by Schumann.

Mammillaria beguinii and *Echinocactus beguinii* Weber are referred by Weber (Dict. Hort. Bois 466. 1896) as synonyms of *Echinocactus erectocentrus*. The Index Kewensis (Suppl. 5) refers the former name to Schelle (Handb. Kakteenk. 200. 1907). The name *E. beguinii* has been previously used in Rebut's Catalogue and by Schumann (Monatsschr. Kakteenk. 5: 44. 1905), but not described.

5. *Neolloydia ceratites* (Quehl).

Mammillaria ceratites Quehl, Monatsschr. Kakteenk. 19: 155. 1909.

Simple or in small clusters, short-cylindric, 6 to 10 cm. high; tubercles somewhat 4-angled, more or less arranged in ribs; young areoles very woolly but becoming naked; radial spines 15 to 20, more or less spreading, white, 1.5 cm. long; central spines 5 or 6, longer and stouter than the radials, blackish above; flowers purple, 3 to 3.5 cm. long; perianth-segments oblong, acute.

Type locality: Mexico.

Distribution: Mexico.

Illustration: Monatsschr. Kakteenk. 19: 155, as *Mammillaria ceratites*.

Figure 16 is from a photograph of the type plant sent us by Mr. Quehl.

6. *Neolloydia conoidea* (De Candolle) Britton and Rose, Bull. Torr. Club 49 252. 1922.

Mammillaria conoidea De Candolle, Mém. Mus. Hist. Nat. Paris 17: 112. 1828.

Mammillaria grandiflora Otto in Pfeiffer, Enum. Cact. 33. 1837.

Mammillaria diaphanacantha Lemaire, Cact. Aliq. Nov. 93. 1838.

Mammillaria inconspicua Scheidweiler, Bull. Acad. Sci. Brux. 5: 49. 1838.

Mammillaria echinocactoides Pfeiffer, Allg. Gartenz. 8: 281. 1840.

Mammillaria scheeri Mühlenpfordt, Allg. Gartenz. 13: 346. 1845.

Mammillaria strobiliformis Engelman in Wislizenus, Mem. Tour North. Mex. 113. 1848.

Echinocactus conoideus Poselger, Allg. Gartenz. 21: 107. 1853.

Cactus conoideus * Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus echinocactoides * Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus grandiflorus Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Linnaeus, 1753.

Sometimes simple, hut usually cespitose, sometimes forming large clusters, often branching or budding above, short-cylindric; tubercles in 5 or 8 spiral rows, obtuse, their axils very woolly; spines very numerous, often completely covering the plant; radial spines white, 25 or more, widely




Fig. 16.—*Neolloydia ceratites*.



Fig. 17.—*Neolloydia conoidea*.

spreading, 8 to 10 mm. long; central spines several, stouter and longer than the radials, 1 to 3 cm. long, blackish; flowers large; outer perianth-segments dull purple without, lighter toward the margins inner perianth-segments rich purple; anthers orange; filaments pale yellow, purplish at base; style and stigma-lobes pale yellow, the latter 5 or 6; fruit compressed-globose, dull yellow, mottled with red, becoming dry and papery, then brown; seeds 1 mm. in diameter.

Type locality: Mexico.

Distribution: Northern Mexico.

Mammillaria canescens, listed by De Candolle (Prodr. 3: 460. 1828) as hardly known and given by Pfeiffer (Enum. Cact. 33. 1837) as a synonym of *M. grandiflora*, doubtless belongs here. A plant of this name was in the Berlin Botanical Garden in 1829 (Verh. Ver. Beförd. Gartenb. 6: 430. 1830).

The name *Coryphantha conoidea* occurs in C. R. Orcutt's Circular to Cactus Fanciers 1922.

Illustrations: De Candolle, Mém. Cact. pl. 2; Pfeiffer, Abbild. Beschr. Cact. 2: pl. 26; Blühende Kakteen 2: pl. 96; Schelle, Handl. Kakteenk. 238. f. 155; Ann. Rep. Smiths.

* Kuntze's spelling of these two names is as follows: *C. conodeus* and *C. echinocactodes*.

Inst. 1908: pl. 14, f. 1; Thomas, Zimmerkultur Kakteen 46, as *Mammillaria conoidea*; Monatsschr. Kakteenk. 6: 119, as *Mammillaria grandiflora*.

Figure 15 is from a photograph of a barren plant collected by Dr. Safford in Mexico in 1907 (No. 1334); figure 17 is from a photograph of a flowering plant collected by Dr. Chaffey in the state of Zacatecas, Mexico, July 4, 1910.

Related to the preceding is:

MAMMILLARIA CREBRISPINA De Candolle, Mém. Mus. Hist. Nat. Paris 17: 111. 1828.

Cactus crebrispinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

This plant was collected by Thomas Coulter but its identification is very uncertain. Pfeiffer thought that it was related to *Mammillaria conoidea* and perhaps it should be referred there.

Mammillaria polychlora Scheidweiler (Förster, Handb. Cact. 205. 1846) was given as a synonym of *M. crebrispina*.




FIG. 18.—*Neolloydia texensis*.

7. *Neolloydia texensis* sp. nov.

Globular to short-oblong, 4 to 6 cm. long; tubercles arranged in long spirals, somewhat imbricated, a little flattened dorsally; radial spines 10 to 15, white, widely spreading, about 1 cm. long; central spines 1 to 3, much stouter than the radials, elongated, 2 to 3 cm. long, black; flowers not seen; fruit small, globular, almost hidden by the spines, greenish, thin-walled, dry; seeds black, tuberculate, 1.5 mm. in diameter; hilum large, basal, white lunate.

Collected by MacDougal and Shreve at Sanderson, Texas, December 1920.

This seems to be the plant from Texas referred by Engelmann to *Mammillaria scolymoides* but it probably is not that species which came from central Mexico. *M. scolymoides* probably should be considered a synonym of *Coryphantha cornifera*, the species of which Engelmann once thought that it might be only a form. Coulter (Contr. U. S. Nat. Herb. 3: 115. 1894) treats the Texan plant under the name of *Cactus scolymoides* but the range which he gives is too wide, and doubtless more than one species is involved, both in his description and range. The only specimen which we have seen of this species, except MacDougal and Shreve's plant, is one collected by Walter M. Evans in 1891, which is mixed with *Cactus echinus* and labeled as from near El Paso, Texas.

Figure 18 is from a photograph of plants collected by Dr. MacDougal and Dr. Shreve.

4. MAMILLOPSIS * (Morren) Weber.

Cespitose cacti, often forming large clusters, globular or short-cylindric, completely hidden under a mass of long, soft, white, hair-like spines; tubercles not arranged in ribs, more or less conic, not grooved above, spine-bearing at apex, their axils pubescent and bristly; radial spines numerous, weak, straight; central spines 4 to 6, with yellow, hooked tips; flowers from near top of plant but apparently from axils of old areoles, with a regular, straight, slender, scaly tube and a broad, spreading limb; perianth-segments oblong, obtuse; stamens and style erect, long-exserted beyond tube; scales on flower-tube orbicular, obtuse.


Schumann associated *Mammillaria senilis* Loddiges, the type of the genus, with species now referred to *Cochemiea*, treating them all as a subgenus of *Mammillaria*, but *Cochemiea* has an irregular flower and otherwise is different from this genus.

Morren first proposed the subgeneric name *Mamillopsis*, but Weber, we believe, was justified in recognizing the genus. He states, very properly, that the flowers are very unlike those of any of the species of *Mammillaria*. He also calls attention to the long-exserted stamens, and long and scaly flower-tube, and also to the fact that the filaments are borne in two series, one series being on the flower-tube. The ovary, too, seems to be scaly, and doubtless other differences will be recorded when the species are better known. Two species are here recognized, both from the high mountains of Mexico.

The generic name, *Mamillopsis*, means *Mammillaria*-like.

KEY TO SPECIES.

- Flowers 6 to 7 cm. long, orange-yellow 1. *M. senilis*
 Flowers 3 cm. long, deep red 2. *M. diguetii*



FIGS. 19 and 20.—*Mamillopsis senilis*.

1. *Mamillopsis senilis* (Loddiges) Weber.

Mammillaria senilis Loddiges in Salm-Dyck, Cact. Hort. Dyck. 1849. 82. 1850.
Cactus senilis Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Haworth, 1824.

Stems 6 to 15. cm. high, 3 to 6 cm. in diameter, the flesh juicy and drying red; tubercles 3 to 4 mm. long; spines 30 to 40, 2 cm. long; flowers 6 to 7 cm. long, 6 cm. broad, orange-yellow; perianth-segments oblong, acute, with serrated margin; stigma-lobes 6, spreading; fruit not known.

Type locality: Not cited.

Distribution: High mountains of Chihuahua and Durango.

* *Mamillopsis* has never been formally published as a genus, but it is mentioned by Weber as a synonym of *Mammillaria senilis* (Dict. Hort. Bois 805. 1898). It was proposed as a subgenus by Morren in 1874 (Belg. Hort. 24: 33).

This species was probably first collected by Seemann in the Sierra Madre of Mexico, where it was collected by Dr. Rose in 1897. It has frequently been introduced into cultivation but does not do well, soon dying out. It is able to stand considerable cold and in its home is usually covered with snow during the winter.

Salm-Dyck gave two varieties without descriptions, based on two unpublished names, when he first listed *Mammillaria senilis*, as follows: *M. senilis haseloffii* (Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850; *M. haseloffii* Ehrenberg, Allg. Gartenz. 17: 303. 1849) and *M. senilis linkei* (Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850; *M. linkei* Ehrenberg). The former, however, was published the previous year as *M. haseloffii* and has priority.

Illustrations: Fl. Serr. 21: pl. 2159; Rev. Hort. iv. 2: pl. 334; Belg. Hort. 24: pl. 3; Cact. Journ. 1: pl. for March; Contr. U. S. Nat. Herb. 5: pl. 62; Schelle, Handb. Kakteenk. 245. f. 163; Tribune Hort. 4: pl. 140; De Laet, Cat. Gén. 28. f. 41; Gartenwelt 14: 331; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 31; Succulenta 4: 80, as *Mammillaria senilis*.

Figure 19 is from a photograph of a flowering plant; figure 20 is from a photograph of two flowers of a plant obtained in the Sierra Madre, Mexico, by I. Ochoterena in 1911; figure 21 is reproduced from the third illustration cited above.




FIG. 21.—*Mamilloopsis senilis*.

2. *Mamilloopsis diguetii* (Weber).

Mammillaria senilis diguetii Weber, Bull. Mus. Hist. Nat. Paris 10: 383. 1904.

Plants densely cespitose, forming a hemispheric clump of about 35 globular heads, each 25 cm. in diameter; radial spines numerous, dark straw-colored; flowers 3 cm. long, about 2 cm. broad, deep red; ovary bearing small scales.

Type locality: Sierra de Nayarit, Jalisco.

Distribution: Jalisco to Sinaloa, Mexico.

This species, until recently, was known only from the single collection of L. Diguët made in March 1900; he found it in the mountains of Jalisco at an altitude of 2,500 meters. It has again been collected by J. G. Ortega in the Sierra de Chabarra, Concordia, Sinaloa, in 1921.

The type is in the Museum of Natural History of Paris and was studied there by Dr. Rose in May 1912; he believes that it is distinct from *M. senilis*, the spines being of a different color and much more rigid than in that species.

5. **COCHEMIEA** (K. Brandege) Walton, Cact. Journ. 2: 50. 1899.

Plant-body cylindric, often much elongated, the surface covered with spirally arranged tubercles, these not milky; tubercles not grooved above; spines both central and radial; flowers borne from axils of upper old tubercles, narrowly tubular, curved and bilabiate; perianth-segments in 2 series; stamens and style red, exserted; ovary naked; fruit indehiscent, globular, red, naked, bearing a large scar at top; seeds black, reticulated.

Type species: *Mammillaria halei* Brandege.

The genus was named for an Indian tribe which once inhabited Lower California. Mrs. Brandege, who first separated these species as a subgenus, describes the flowers as "scarlet, tubular, slender, somewhat curved, and oblique, with spreading unequal petaloid sepals, so making the flower apparently double as in *Cereus flagelliformis*."

Four species are known, all inhabiting Lower California.




FIG. 22.—*Cochemiea halei*.




FIG. 23.—*Cochemiea poselgeri*.

The fact that *Cochemiea* had been raised to generic rank, to which four species had been transferred, has been overlooked by all our botanical indexes. Walton's remarks in this connection are interesting:

"The plants so classed have flowers very elongated, tubular, with sepals placed as a second ring, removed some distance below the petals; they are oblique like *Epiphyllum truncatum* and *Cereus flagelliformis* and in fact more resemble those flowers than they do those of any *Mammillaria*, so much so that I think it would be best to drop the generic name of *Mammillaria* and simply adopt Mrs. Brandege's name of *Cochemiea* as a generic name."

Mrs. Brandege suggested (*Erythea* 5: 117), "It is possible that some of the elongated species of Mexico proper will be found to belong to this section when the flowers are better known." But we have seen no plants from the mainland of Mexico which suggest this relationship.

KEY TO SPECIES.

- Spines all straight. 1. *C. halei*
- Some or all of central spines hooked. 2. *C. poselgeri*
- Central spine normally solitary. 3. *C. setispina*
- Central spines normally 2 to 11 (sometimes only 1 in *C. setispina*). 4. *C. pondii*
- Central spines 1 to 4. 3. *C. setispina*
- Central spines 8 to 11. 4. *C. pondii*

1. *Cochemiea halei* * (Brandege) Walton, Cact. Journ. 2: 50. 1899.

Mammillaria halei Brandege, Proc. Calif. Acad. II. 2: 161. 1889.

Cactus halei Coulter, Contr. U. S. Nat. Herb. 3: 106. 1894.

Cespitose; stems nearly upright, often 30 to 50 cm. high, 5 to 7.5 cm. in diameter, almost entirely covered by the spines; tubercles short; axils of tubercles woolly but not setose; radial spines 10 to 20, 10 to 12 mm. long; central spines 3 or 4, 25 mm. long, all straight; flowers central or nearly so, 4 to 5 cm. long; filaments yellow; stigma-lobes scarlet; fruit scarlet, 12 mm. long; seeds reticulated.

Type locality: Magdalena Island, Lower California.

Distribution: Islands of southern Lower California.

This species was observed first by Mr. T. S. Brandege in 1889, while making a botanical excursion through Lower California, and described by him the same year. It has been reported from only two islands off the coast of Lower California but it is there very abundant. It has been introduced into Europe and is sometimes offered in the trade. It is remarkable for its very large slender flowers. An abundance of material was collected by Dr. Rose in 1911. The plant does not do well in cultivation.

The species was named for Mr. J. P. Hale, who had extensive domains in Lower California and who assisted Mr. Brandege while making explorations in 1889.

Illustrations: Proc. Calif. Acad. II. 2: pl. 6; Monatsschr. Kakteenk. 5: 89; Schumann, Gesamtb. Kakteen 510. f. 84; Thomas, Zimmerkultur Kakteen 47, as *Mammillaria halei*.

Figure 22 is from a photograph of a barren shoot of a specimen collected by C. R. Orcutt at Magdalena Bay, Lower California, 1917.

2. *Cochemiea poselgeri* (Hildmann).

Mammillaria poselgeri Hildmann, Garten-Zeitung 4: 559. 1885.

Mammillaria roseana Brandege, Zoe 2: 19. 1891.

Mammillaria radliana Quehl, Monatsschr. Kakteenk. 2: 104. 1892.

Cactus roseanus Coulter, Contr. U. S. Nat. Herb. 3: 105. 1894.

Cochemiea rosiana Walton, Cact. Journ. 2: 50. 1899.

Stems numerous from a central root, spreading or sometimes pendent from rocks or creeping over the ground, often 2 meters long, 4 cm. thick; areoles and upper axils white-woolly, the latter rarely setose; tubercles remote, somewhat flattened; radial spines 7 to 9, 9 to 12 mm. long, straw-colored; central spine 1, hooked, 25 mm. long; flowers appearing in the upper axils, 3 cm. long, scarlet; stamens and style exserted; fruit globular, 6 to 8 mm. in diameter.

Type locality: Cape Region, Lower California.

Distribution: At lower elevations in southern Lower California.

This cactus, according to Mr. Brandege, is one of the most showy of this region.

Mammillaria longihamata Engelm. was a manuscript name taken up by Coulter (Contr. U. S. Nat. Herb. 3: 105. 1894) as a synonym of *Cactus roseanus*.

Illustrations: Thomas, Zimmerkultur Kakteen 49; Monatsschr. Kakteenk. 2: 105, as *Mammillaria radliana*; Garten-Zeitung 4: 559. f. 131; Schelle, Handb. Kakteenk. 246. f. 164, as *M. poselgeri*.

Plate 11, figure 3, shows a plant collected by Dr. Rose at Cape San Lucas, Lower California, which flowered in the New York Botanical Garden in 1915; figure a shows the fruit and figure 3b the seed from a plant collected by Dr. Wm. S. W. Kew near La Junta, Lower California, November 10, 1920. Figure 23 is from a photograph of a plant collected by C. R. Orcutt near Magdalena, Lower California, and sent to the Bureau of Chemistry, U. S. Department of Agriculture, in 1917.

2. *Cochemiea setispina* (Coulter) Walton, Cact. Journ. 2: 51. 1899.

Cactus setispinus Coulter, Contr. U. S. Nat. Herb. 3: 106. 1894.

Mammillaria setispina Engelm. in K. Brandege, Erythea 5: 117. 1897.

Stems ascending, 30 cm. high; tubercles short; axils of tubercles woolly but not setose; radial spines 10 to 12, white with black tips, widely spreading, unequal, 10 to 34 cm. long, slender; central spines 1 to 4, stouter than the radials, one of them strongly hooked; flowers not definitely known but probably large; fruit obovoid, 3 cm. long, scarlet; seeds black and pitted.

* Walton published this name as *Cochemiea hallei*.

Type locality: San Borgia, Lower California.

Distribution: Interior of southern Lower California.

We have not seen living specimens of the species. Dr. Rose obtained a small specimen from L. Quehl at Halle in 1912.

The type of this species, now in the herbarium of the Missouri Botanical Garden, was collected by William Gabb in 1867, while Brandegee obtained specimens in 1889. Dr. C. A. Purpus found it near Calmalli and wrote of it as follows (Cact. Journ. 2: 54. 1899):

"My next trip was to a chain of granite mountains about 20 miles from Calmalli.

"I was very much surprised to find on the slope of the mountains *Mamillaria setispina* Engelm., which until now I had not been able to collect as a living specimen. I came upon it afterwards also in gneiss, trachyt, porphur, and in a sandstone conglomerate. Ground composed of granite gravel appears to suit it best."

4. *Cochemiea pondii* (Greene) Walton, Cact. Journ. 2: 51. 1899.

Mamillaria pondii Greene, Pittonia 1: 268. 1889.

Cactus pondii Coulter, Contr. U. S. Nat. Herb. 3: 102. 1894.

Stems at first upright, cylindric, simple or few-branched, 7 cm. to 3 dm. high, hidden under a dense covering of spines; axils of tubercles setose; young areoles white-tomentose; radial spines white, whitish or sometimes brownish, 15 to 25, spreading; central spines 8 to 11, much longer and stouter than the radials, the longest 3 cm. long, 1 or 2 hooked; flowers slender, 5 cm. long, bright scarlet; stamens exserted; fruit purplish red, 18 mm. long, ovoid to obovoid.


Type locality: Cedros Island.

Distribution: Islands off the western coast of northern Lower California.

This plant was found in great abundance on Cedros Island by Dr. Rose in 1911 (No. 16090) and a number of living specimens was brought to Washington and New York. These have been in cultivation for more than ten years but have never flowered. It is not often met with in cultivation.

The species was named for Charles Fremont Pond, U. S. N., who collected plants on Cedros and other islands off the coast of Lower California in 1889.

Figure 24 shows the fruit and figure 24a the seed from specimens obtained at the type locality by Dr. Rose in 1911.



FIGS. 24 and 24a.—Fruit and seed of *Cochemiea pondii*.

6. *CORYPHANTHA* (Engelmann) Lemaire, Cactées 32. 1868.

Plant body globular to cylindric, either solitary or cespitose; tubercles, except the very earliest ones, grooved on upper surface* from apex to base; flowers from near top of plant and from base of young and growing tubercles, large and showy, generally yellow, sometimes purple or red; ovary naked or, occasionally, bearing a few scales in some species; perianth long-persistent †; fruit large, ‡ ripening slowly, ovoid to oblong, greenish or yellowish; seeds brown (black and angled in *Coryphantha cubensis*), lightly reticulated or nearly smooth, thin-shelled, with a central or subventral hilum; embryo curved, at least in some species.

Type species:§ *Mamillaria sulcolanata* Lemaire.

The generic name is from *κορυφή* top, and *ἄνθος* flower, referring to the insertion of the flowers at the top of the plant. We recognize 37 species in the genus. The genus *Coryphantha* was proposed by Lemaire in 1868, but he did not designate a type. The

* In *C. macromeris* the tubercle is grooved only for about half its length.

† We quote the following observation of Engelmann in this connection: "I have repeatedly observed, and in a considerable number of species, that the red berries of the *Mamillariae* are always destitute of the remnants of the perigone, but the green fruits always are topped with it (Mem. Tour North. Mex. 21).

The only fruit which we have seen of *C. nickel* was globose and small, 5 to 7 mm. in diameter, but the species otherwise of this alliance.

§ See Britton and Milispough, Bahama Flora 295. 1920.

name, however, comes from Engelmann, who first used it as a subgenus of *Mammillaria* (Proc. Amer. Acad. 3: 264. 1856).

The position of this group has always been puzzling to cactus students. Dr. Poselger believed that it was a section of *Echinocactus* and transferred certain of these species which had been described under *Mammillaria* to *Echinocactus*. In its vertical, nearly central flowers it does approach the *Echinocactanae*, but otherwise it is quite distinct.

In the origin of their large flowers, in the shape and structure of their fruit, and in the color and form of their seeds the species compose a rather natural group, but they are diverse in form and armament. The species are most common in central Mexico, a few extending into the southern United States, and one extending into southern Canada.

The groove on the upper side of the tubercle which is so characteristic of the genus does not occur on seedlings or on very young plants, but it is always found on old flowering plants and seems to be associated with the inflorescence, for the flowers appear only in the axils of grooved tubercles and originate at the bottom of this groove. Plants which grow in conservatories for a long time without flowering lose this groove; * we have had one plant of this kind under observation for fifteen years.

KEY TO SPECIES.

- A. Seeds brown, not angled; flowers usually large.
 - B. Tubercles grooved to middle or a little below; ovary bearing scales with woolly axils. Series *Macromeres*.
 - Tubercles elongated, bright green 1. *C. macromeris*
 - Tubercles short, grayish green 2. *C. runyonii*
 - BB. Tubercles grooved from tip to base except in young plants; ovary naked.
 - C. Grooves of tubercles bearing large yellow or red glands. Series *Recurvatae*.
 - Flowers white 3. *C. ottonis*
 - Flowers not white.
 - Stems globular.
 - Radial spines more or less recurved 4. *C. recurvata*
 - Radial spines spreading or ascending.
 - Spines dark, sometimes black 5. *C. poselgeriana*
 - Spines yellow or sometimes tinged with red.
 - Central spines slender and flexible 6. *C. muehlenpfordtii*
 - Central spines stout and rigid.
 - Radial spines subulate 7. *C. guerkeana*
 - Radial spines acicular 8. *C. echinoidea*
 - Stems cylindric.
 - Stems bluish green 9. *C. clava*
 - Stems yellowish green.
 - Central spine usually one.
 - Glands in groove red 10. *C. octacantha*
 - Glands in groove yellow 11. *C. exsudans*
 - Central spines 2 12. *C. erecta*
 - CC. Grooves of tubercles without large glands. Series *Sulcolanatae*.
 - D. Outer perianth-segments not ciliate.
 - E. Flowers purplish or rose 13. *C. elephantidens*
 - EE. Flowers yellow or white.
 - F. Tubercles very large, broader than high 14. *C. bumamma*
 - FF. Tubercles of medium size, if large, longer than broad.
 - G. Plants large for this genus (often 8 cm. in diameter); seeds 3 mm. in diameter 15. *C. robustispina*
 - GG. Plants much smaller than in *C. robustispina*; seeds 2 mm. in diameter or less.
 - H. Central spines usually wanting.
 - Secondary cluster of spines developed in upper part of areoles and connivent at top! 16. *C. connivens*
 - Secondary cluster of spines not developed.
 - Spines pectinate 17. *C. pectinata*
 - Spines not pectinate.
 - Spines 14 or more.
 - Spines slender with long black tips 18. *C. nickelsae*
 - Spines rather short with light tips.
 - Spines subulate 19. *C. compacta*
 - Spines acicular 20. *C. radians*
 - Spines fewer than 15.
 - Spines slender and weak 21. *C. sulcolanata*
 - Spines not slender 22. *C. retusa*

* *Mammillaria potosiana* and *M. polymorpha* seem to have been based on such plants.

- HH. Central spines one to several.
 I. Central spines strongly hooked23. *C. palmeri*
 II. Central spines straight or at most curved.
 J. Central spines more or less curved.
 Central spine one, sometimes more in No. 25.
 Radial spines nearly as long as central24. *C. cornifera*
 Radial spines about half as long as central25. *C. salm-dyckiana*
 Central spines several.
 Radial spines 20 or more26. *C. pallida*
 Radial spines 12 or fewer27. *C. pycnanantha*
 JJ. Central spines straight.
 Radial spines, two kinds (to be looked for here). ... 5. *C. poselgeriana*
 Radial spines of one kind.
 Plant almost hidden under mass of spines;
 fruit oblong28. *C. echinus*
 Plant not hidden under mass of spines; fruit
 globular29. *C. durangensis*
 DD. Outer perianth-segments ciliate.
 Flowers yellow30. *C. chlorantha*
 Flowers purplish to pink.
 Inner perianth-segments linear or lanceolate.
 Stigma-lobes purple, apiculate31. *C. vivipara*
 Stigma-lobes white, obtuse or notched.
 Flowers 4 to 7 cm. broad, rose to purple.
 Plants mostly solitary; inner perianth-segments broadly linear ...32. *C. neo-mexicana*
 Plants mostly cespitose; inner segments linear-lanceolate33. *C. arizonica*
 Flowers very short, 3 cm. broad, light pink34. *C. deserti*
 Inner perianth-segments oblanceolate35. *C. aggregata*
 AAA. Seeds black, angled; flowers minute. Series *Cubenses*36. *C. cubensis*
 AA. Ungrouped species37. *C. sulcata*

1. *Coryphantha macromeris* (Engelmann) Lemaire, Cactées 35. 1868.

Mammillaria macromeris Engelmann in Wislizenus, Mem. Tour North. Mex. 97. 1848.

Mammillaria heteromorpha Scheer in Salm-Dyck. Cact. Hort. Dyck. 1849. 128. 1850.

Echinocactus macromeris Poselger, Allg. Gartenz. 21: 102. 1853.

Echinocactus heteromorphus Poselger, Allg. Gartenz. 21: 126. 1853.

Mammillaria dactylithele Labouret, Monogr. Cact. 146. 1853.

Cactus macromeris Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus heteromorphus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Plant branching at base, often many-headed, up to 2 dm. long; tubercles large, soft, loosely arranged, elongated, 12 to 30 cm. long, grooved on upper side about two-thirds of their length; spines 10 to 17, slender, the radials white; central spines several, black, the longer ones 5 cm. long; flowers large, purple, 6 to 8 cm. broad; scales on flower-tube ciliate; ovary bearing a few scales with hairy axils; fruit 15 to 25 mm. long; seeds globose, brown but sometimes described as yellow, smooth.

Type locality: Near Doñana, New Mexico.

Distribution: Southern New Mexico, western Texas, and Chihuahua, south to Zacatecas, Mexico.

This species and the following one are not closely related to the others of this genus. The tubercles are much more elongated and flattened, and the groove on the upper surface never extends to the base. Sometimes a branch or bulblet is produced instead of a flower.

Here may belong *Coryphantha heteromorpha* Lemaire (Cactées 34. 1868); this name is apparently erroneously referred to in the Index Kewensis (1: 624) as *Coryphantha heterophylla* (see *Ariocarpus fissuratus*, Cactaceae 3 83).

Mammillaria brownii Toumey was erroneously referred here by Schumann.

Mammillaria macromeris var. *longispina* and var. *nigrispina* are mentioned by Schelle (Handb. Kakteenk. 237. 1907).




FIG. 25.—Tubercles of *Coryphantha macromeris*.

Illustrations: Cact. Journ. 1: 43; Förster, Handb. Cact. ed. 2. 399. f. 41; Rümpler, Sukkulenten 205. f. 116; Dict. Gard. Nicholson 4: 564. f. 36; Suppl. 517. f. 552; Goebel, Pflanz. Schild. 1: pl. 1, f. 6; Amer. Gard. 11: 460; West Amer. Sci. 13: 39; Cact. Mex. Bound. pl. 14, 15; Cycl. Amer. Hort. Bailey 2: f. 746a, 1355; Stand. Cycl. Hort. Bailey 4: f. 2314; Gartenflora 42: 543. f. 111; Schelle, Handb. Kakteenk. 237. f. 152; Balt. Cact. Journ. 1: 21; Watson, Cact. Cult. 165. f. 64; ed. 3. f. 41, as *Mammillaria macromeris*.

Figure 25 is from a drawing of two tubercles, showing the grooves on the upper side, of a plant sent by Mrs. S. L. Pattison from western Texas. At the base of one is shown the flower-scar; in the other is a small bud.

2. *Coryphantha runyonii* sp. nov.

Forming low clumps, sometimes 5 dm. in diameter, grayish green, with a thick, elongated tap-root; tubercles rather short, 1 to 2 cm. long, terete or somewhat flattened, grooved on the upper half, rarely more, but never to the base; radial spines 6 or more, spreading, acicular, very variable in length, 3 cm. long or less, sometimes all yellow or sometimes one or more in a cluster brown, otherwise yellow; central spines on young plant solitary, dark brown to black but in old plants sometimes 2 or 3, somewhat angled, up to 6 cm. long; flowers large, purple, 5 cm. broad; outer perianth-segments ciliate; inner perianth-segments spatulate, oblong, acute; fruit green; seeds brown.

Found along the Rio Grande from Brownsville to Rio Grande City. This species has been repeatedly observed by Robert Runyon, from whom we received living plants in 1921 (No. 15, type) and 1922.

Mr. Runyon wrote us about the plant as follows:

"I also inclose you herewith two photographs of the plant you have called *Coryphantha runyonii*. I first became interested in this plant about two years ago when I saw it growing near Rio Grande, Texas. It was found at one place only, but in abundance. It grows on the gravel hillside and down in the lower land in a kind of white silt soil.

"The fruit is green and the flowers are a very pretty pink to a purple with a delicate fringed petal. The tubercles are very irregular. The largest plants are about 18 inches in diameter and would weigh not less than fifty pounds."

Plate 1, figure 1, is from a photograph sent us by Robert Runyon.

3. *Coryphantha ottonis* (Pfeiffer) Lemaire, Cactées 34. 1868.

Mammillaria ottonis Pfeiffer, Allg. Gartenz. 6: 274. 1838.

Echinocactus ottonianus Poselger, Allg. Gartenz. 21: 102. 1853.

Cactus ottonis Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Lehmann, 1827.

Mammillaria bussleri Mundt in Schumann, Monatsschr. Kakteenk. 12: 47. 1902.

Mammillaria golziana Haage Jr., Monatsschr. Kakteenk. 19: 100. 1909.

Simple, globular to short-cylindric, 12 cm. high or less, 8 cm. in diameter, glaucous to grayish green; radial spines 8 to 12, nearly equal, 8 to 10 mm. long; central spines 3 or 4, longer and a little stouter than the radials; axils of flowering tubercles woolly; flowers white, 4 cm. long; outer perianth-segments oblong, obtuse; inner perianth-segments apiculate; stigma-lobes 10, green.

Type locality: Mineral del Monte, Mexico.

Distribution: Central Mexico.

The name here used was proposed by Lemaire (Cactées 34) in 1868 but not formally published. *Mammillaria ottonis tenuispina* Pfeiffer is sometimes used but we have seen no formal description.

Nicholson (see also Watson, Cact. Cult. 168. f. 66; and ed. 3. f. 40) describes and illustrates (Dict. Gard. Nicholson Suppl. 517. f. 553) under this name a very peculiar specimen in which the flowers are borne away from the top of the plant; it is doubtless not congeneric with this species. Nicholson's description is here quoted:

"Flowers white, large for the size of the plant. May and June. Stem small, compressed, 3 in. across, with numerous compressed tubercles, and short hair-like spines (Mexico. 1834. See fig. 553). There is another species called *M. ottonis*, having a large spiny stem."

Here we believe belong some of the plants which are passing as *Mammillaria golziana*. Very different, however, are the two published illustrations of Kunze (Cact. 1910 and Monatsschr. Kakteenk. 19: 101. 1909), which also seem to differ from each other.

Illustrations: Monatsschr. Kakteenk. 12: 47, as *Mammillaria bussleri*; Monatsschr. Kakteenk. 27: 3. f. a, as *Mammillaria golziana*; Monatsschr. Kakteenk. 27: 3. f. b, as *Mammillaria ottonis*.


4. *Coryphantha recurvata* (Engelmann).

Mammillaria recurvispina Engelmann, Proc. Amer. Acad. 3: 266. 1856. Not De Vriese, 1839.

Mammillaria recurvata Engelmann, Trans. St. Louis Acad. 2: 202. 1863.

Cactus recurvatus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

*Cactus engelmannii** Kuntze, Rev. Gen. Pl. 1: 260. 1891.



FIGS. 26 and 27.—*Coryphantha recurvata*.

Plant-body depressed-globose, 10 to 20 cm. in diameter, often forming large masses 30 to 90 cm. in diameter and sometimes with over 50 heads; tubercles low; radial spines about 20, yellow to gray, with dark tips, pectinate, recurved; central spines 1, rarely 2, longer and darker than the radials, 12 to 20 mm. long, more or less reflexed, often appressed; flowers 25 to 35 mm. long, said to be brownish outside; inner perianth-segments lemon-yellow; fruit not known.

Type locality: Sonora. Explained in the Cactaceae of the Mexican Boundary to be eastern parts of Pimeria Alta in Sonora, especially in the Sierra del Pajarito.

Distribution: Arizona and Mexico, especially along the United States-Mexican Boundary near Nogales.

Engelmann describes a peculiar flowering habit for *Coryphantha* when he says that the flowers originate in the base of the grooves of full-grown tubercles, being scattered over the top of the plant. We have also noticed this character; not only are the flowers borne in the axils of mature tubercles, but they are produced in great abundance in a circle 5 to 6 cm. in diameter.

* It is possible that Lemaire also gave the name *Coryphantha engelmannii* for *Mammillaria recurvispina*, though this is not shown by the text.

† See Cact. Mex. Bound. 12. 1859.

Otto Kuntze made the binomial *Cactus engelmannii* because, as he states, the name *Mammillaria recurvispina* De Vriese had priority over Engelmann's name. Engelmann, however, had long before renamed his plant.

Mammillaria nogalensis Runge (Schumann, Gesamtb. Kakteen 494. 1898) has been referred here as a synonym, but this name had already been used by Walton.

Illustrations: Schelle, Handb. Kakteenk. 239. f. 156, as *Mammillaria recurvata*; Cact. Journ. 1: pl. for March; 2: 148; pl. for September, as *M. nogalensis*.

Figure 26 is from a photograph by Dr. MacDougal at Calabasas, showing a clump; figure 27 is from a photograph of a plant sent by F. J. Dyer from Nogales.

5. *Coryphantha poselgeriana* (Dietrich).

Echinocactus poselgerianus Dietrich, Allg. Gartenz. 19: 346. 1851.*

Echinocactus saltillensis Poselger, Allg. Gartenz. 21: 101. 1853.

Echinocactus salinensis Poselger, Allg. Gartenz. 21: 106. 1853.

Mammillaria difficilis Quehl, Monatsschr. Kakteenk. 18: 107. 1908.

Mammillaria valida J. A. Purpus, Monatsschr. Kakteenk. 21: 97. 1911. Not Weber, 1898.

Plant-body large for the genus, globular, bluish green; tubercles large, closely packed together and at base strongly angled; radial spines of two kinds, the 4 or 5 lower ones spreading, subulate, reddish to black, about as long as the single central one (2 to 4 cm. long); the upper radials, 5 to 8, ascending together, yellowish with black tips, weak, acicular; flower large, 4 to 5 cm. long and nearly as broad when expanded, flesh-colored, the segments spatulate, usually rounded at apex; fruit oblong, 15 mm. long; seeds brownish.

Type species: Near Saltillo, Mexico.

Distribution: States of Nuevo Leon, Coahuila, and Zacatecas, Mexico.

Two different plants have been passing under the name *Echinocactus saltillensis*. The one now in the trade, called *E. ingens* var. *saltillensis* by Schumann, is a very large plant and is a true *Echinocactus* which we have already elsewhere described as *E. palmeri*;† the other, which is the one originally described by Poselger, is a small globular *Coryphantha* and has usually been taken for *Mammillaria scheeri*, more recently described as *M. valida*.

The clusters of connivent weak spines, so characteristic of this species, are not always shown in young plants and this may account for certain seeming discrepancies in the original descriptions. The nascent spines are sometimes red, bleaching white; the gland in the groove of the tubercle is bright red.

Illustrations: De Laet, Cat. Gén. f. 44; Schelle, Handb. Kakteenk. 239. f. 157; Tribune Hort. 4: pl. 139; Rev. Hort. Belg. 40: after 196, as *Mammillaria radians*; Monatsschr. Kakteenk. 21: 99, as *Mammillaria valida*; (?) Blanc, Cacti 50. No. 599; (?) Cact. Journ. 2: 55, as *Echinocactus poselgerianus*; Monatsschr. Kakteenk. 18: 107, as *Mammillaria difficilis*; Rother, Praktischer Leitfaden Kakteen 31, as *Echinocactus scheeri*.

6. *Coryphantha muehlenpfordtii* (Poselger).

Mammillaria scheeri Muehlenpfordt, Allg. Gartenz. 15: 97. 1847. Not Muehlenpfordt, 1845.

Echinocactus muehlenpfordtii Poselger, Allg. Gartenz. 21: 102. 1853.

Mammillaria scheeri valida Engelmann, Proc. Amer. Acad. 3: 265. 1856.

Coryphantha scheeri Lemaire, Cactées 35. 1868.

Cactus scheeri Kuntze, Rev. Gen. Pl. 1: 261. 1891*

Plants nearly globular, usually simple, short-oblong, 20 cm. long, 7.5 to 15 cm. in diameter; tubercles large, 1 to 2.5 cm. long; axils of young tubercles grooved and young spine-areoles very woolly; grooves bearing large dark-colored glands; spines variable, reddish to yellow with brown to black tips; radials 6 to 16, usually about 2 cm. long, straight; central spines 1 to 4, subulate, stouter than the radials, 3 to 3.5 cm. long, from nearly straight to curved at tip or even strongly hooked; flowers yellow, 6 cm. long; scales on flower-tube and outer perianth-segments more or less lacerated; inner perianth-segments oblong, entire, acute; fruit greenish, oblong, 3 to 3.5 cm. long, naked; seeds large, 3 mm. long, brown, shining, smooth.

Type locality: Mexico.

Distribution: Northern Chihuahua, western Texas, and southern New Mexico.

* We have not seen the type of this species but Bödeker has sent us a copy of the photograph of it left by Poselger.

† See Contr. U. S. Nat. Herb. 12: 290. 1909; Britton and Rose, Cactaceae 3: 172. 1922.

There has been considerable confusion regarding this species, which was first described as *Mammillaria scheeri* by Mühlenpfordt in 1847, but this proved to be a homonym. This led Poselger in 1853, when he transferred the species to *Echinocactus*, to publish it as *E. muehlenpfordtii*.

Dr. Engelmann in 1856 described a variety of *Mammillaria scheeri*, calling it *valida*. Some time afterwards he compared this variety with the type of the species and decided that they were the same. We have examined several specimens from near the type locality of the variety *valida*, which is near El Paso, Texas.

It is possible that Scheer's plant was a very young one, which might account for the differences in form and spines. The *Mammillaria scheeri* of Schumann's Monograph is a complex of 4 or 5 distinct species.

Illustrations: Allg. Gartenz. 15: 97. pl. 2; Förster, Handb. Cact. ed. 2. 406. f. 44; Schumann, Gesamtb. Kakteen 485. f. 80; Monatsschr. Kakteenk. 8: 23; 10: 127; Schelle, Handb. Kakteenk. 237. f. 153, as *Mammillaria scheeri*.




FIG. 28.—*Coryphantha muehlenpfordtii*.




FIG. 29.—*Coryphantha bumamma*.

Figure 28 is from a photograph of a plant collected in western Texas by Mrs. S. L. Pattison in 1920.

7. *Coryphantha guerkeana* (Bödeker).

Mammillaria guerkeana Bödeker, Monatsschr. Kakteenk. 24: 52. 1914.

Plant-body globular, 6 to 7 cm. in diameter; tubercles bluish green, somewhat broader than thick, bearing a large red gland at base of groove and sometimes at top; radial spines 9 to 12, yellow when young, spreading, bulbous at base, rather stout; central spines 3 or 4, rarely one of them stouter, often bent slightly at tip; flowering areoles very woolly; ovary oblong, naked; flower and fruit not seen.

Type locality: Mexico.

Distribution: Durango, Mexico.

This species is near *Coryphantha poselgeriana*, but is smaller and has different spines.

We have seen photographs of the type and have spine-clusters, all obtained from L. Quehl of Halle. We would also refer here specimens obtained by Dr. E. Palmer near Durango City in 1906 (No. 456).

Illustrations: Monatsschr. Kakteenk. **24**: 53, as *Mammillaria guerkeana*; Alianza Científica Universal **3**: pl. opp. 119, as *Mammillaria valida*.

8. *Coryphantha echinoidea* (Quehl).

Mammillaria echinoidea Quehl, Monatsschr. Kakteenk. **23**: 42. 1913.

Plant solitary, globular or a little broader than high, 5 to 6 cm. in diameter, very woolly at apex; tubercles conic, 1.5 cm. high, 1.2 cm. broad at base; groove with 1 to 3 small, grayish glands; areoles elliptic, woolly when young, glabrate in age; radial spines 20 to 25, 1.5 cm. long, white with darker tips; central spines 1 to 3, a little stouter than the radials, one of them porrect, horn-colored; flowers rose-colored, 6 to 8 cm. broad; perianth-segments oblong, broad at apex, denticulate, sometimes mucronate; filaments numerous, red; fruit and seed unknown.

Type locality: Durango.

Distribution: Durango, Mexico.

Illustration: Monatsschr. Kakteenk. **23**: 42, as *Mammillaria echinoidea*.

9. *Coryphantha clava* (Pfeiffer) Lemaire, Cactées 34. 1868

Mammillaria clava Pfeiffer, Allg. Gartenz. **8**: 282. 1840.

Mammillaria schlehtendalii Ehrenberg, Linnaea **14**: 377. 1840.

Mammillaria schlehtendalii levior Salm-Dyck, Cact. Hort. Dyck. 1849. 127. 1850.

Echinocactus clavus Poselger, Allg. Gartenz. **21**: 125. 1853.

Echinocactus schlehtendalii Poselger, Allg. Gartenz. **21**: 125. 1853.

Cactus clavus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.

Cactus schlehtendalii Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

Plant-body club-shaped, deep green; axils of tubercles with white wool and with a red gland at base of groove; tubercles erect, elongated, somewhat 4-sided; spine-areoles white-villous; radial spines usually 7, straight, horn-colored, about equal; central spine 1, a little longer and stouter than the others; flowers very large, sometimes 9 cm. broad, pale yellow, with the outer segments tinged with red; perianth-segments glossy, linear-oblong to spatulate, outer ones entire, inner ones serrate and mucronate at apex; filaments orange; stigma-lobes 6, linear, yellow.

Type locality: Mexico.

Distribution: Mexico.

Coryphantha schlehtendalii Lemaire (Cactées 34. 1868) is usually given as a synonym of this species.

Illustrations: Curtis's Bot. Mag. **74**: pl. 4358; Loudon, Encycl. Pl. ed. 3. 1379. f. 19390, as *Mammillaria clava*.

10. *Coryphantha octacantha* (De Candolle).

Mammillaria octacantha De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 113. 1828.

Mammillaria leucacantha De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 113. 1828.

Mammillaria lehmanni Otto in Pfeiffer, Enum. Cact. **23**. 1837.

Mammillaria macrothele Martius in Pfeiffer, Enum. Cact. **24**. 1837.

Mammillaria plaschnickii Otto in Pfeiffer, Enum. Cact. **24**. 1837.

Mammillaria aulacothele Lemaire, Cact. Aliq. Nov. **8**. 1838.

Mammillaria biglandulosa Pfeiffer, Allg. Gartenz. **6**: 274. 1838.

Mammillaria sulcimamma Pfeiffer, Allg. Gartenz. **6**: 274. 1838.

Mammillaria lehmannii sulcimamma Miquel, Linnaea **12**: 9. 1838.

Mammillaria martiana Pfeiffer, Linnaea **12**: 140. 1838.

? *Mammillaria thelocamptos* Lehmann, Linnaea **13**: Litt. 101. 1839.

Mammillaria aulacothele multispina Scheidweiler, Bull. Acad. Ci. Brux. **6**: 92. 1839.

Mammillaria aulacothele spinosior Monville in Lemaire, Cact. Gen. Nov. Sp. **93**. 1839.

Mammillaria aulacothele sulcimamma Pfeiffer in Walpers, Bot. Repert. **2**: 302. 1843.

Mammillaria aulacothele flavispina Salm-Dyck, Cact. Hort. Dyck. 1844. 13. 1845.

Mammillaria polymorpha Scheer in Mühlenpfordt, Allg. Gartenz. **14**: 373. 1846.

Mammillaria macrothele lehmanni Salm-Dyck, Cact. Hort. Dyck. 1849. 19. 1850.

Mammillaria macrothele biglandulosa Salm-Dyck, Cact. Hort. Dyck. 1849. 19. 1850.

Mammillaria plaschnickii straminea Salm-Dyck, Cact. Hort. Dyck. 1849. 19. 1850.

Echinocactus macrothele Poselger, Allg. Gartenz. **21**: 125. 1853.

Echinocactus plaschnickii Poselger, Allg. Gartenz. **21**: 125. 1853.

Echinocactus macrothele lehmanni Poselger, Allg. Gartenz. **21**: 125. 1853.

Echinocactus macrothele biglandulosus Poselger, Allg. Gartenz. **21**: 125. 1853.

Coryphantha lehmanni Lemaire, Cactées. **34**. 1868.

Coryphantha aulacothele Lemaire, Cactées. **34**. 1868.

Cactus macrothele Kuntze, Rev. Gen. Pl. **1**: 260. 1891.

Cactus aulacothele Kuntze, Rev. Gen. Pl. **1**: 260. 1891.

Cactus biglandulosus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus lehmannii Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus plachnickii Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus octacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus martianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plant-body simple, cylindric, 3 dm. high, 12 to 15 cm. in diameter; axils of tubercles bearing white wool, the groove with 1 or 2 red glands; tubercles elongated, up to 25 mm. long, spreading, somewhat 4-angled but with broad bases; radial spines 8, spreading, rigid, horn-colored with black tips, 10 to 12 mm. long; central spines 1 or 2, stouter than the radials, brownish, 25 mm. long; flowers about 6 cm. broad, straw-colored; perianth-segments linear-oblong, obtuse; filaments reddish; style red; stigma-lobe yellow.

Type locality: Mexico.

Distribution: Central Mexico.

Mammillaria polymorpha Scheer (Mühlenpfordt, Allg. Gartenz. 14: 373. 1846) is probably only an abnormal greenhouse form of this species.

Coryphantha aulacothele and *C. lehmanni* (Lemaire, Cactées 34. 1868) and *M. macrothele nigrispina* (Schelle, Handb. Kakteenk. 243. 1907) are only names but are usually referred here.

Mammillaria leucantha is credited to De Candolle by Steudel (Nom. ed. 2. 2: 97. 1841), but we have not seen such a name used by De Candolle. It may be a misspelling for *M. leucacantha*. Steudel refers the name to *M. lehmannii*, while the Index Kewensis states that it equals *M. recurva*.

Cereus lehmannii Hortus is cited by Förster (Handb. Cact. 245. 1846) as a synonym of *M. lehmanni*.

Illustrations: Loudon, Encycl. Pl. ed. 2 and 3. 1201. f. 17362; Curtis's Bot. Mag. 65: pl. 3634, as *Mammillaria lehmannii*; Monatsschr. Kakteenk. 20: 85; Krook, Handb. Cact. 38, as *Mammillaria aulacothele*; Schelle, Handb. Kakteenk. 242. f. 161; Förster, Handb. Cact. ed. 2. 391. f. 39, as *Mammillaria macrothele*.

11. *Coryphantha exsudans* (Zuccarini) Lemaire.*

Mammillaria exsudans Zuccarini in Pfeiffer, Enum. Cact. 15. 1837.
Mammillaria brevimamma Zuccarini in Pfeiffer, Enum. Cact. 34. 1837.
Mammillaria glanduligera Otto and Dietrich, Allg. Gartenz. 16: 298. 1848.
Mammillaria brevimamma exsudans Salm-Dyck, Cact. Hort. Dyck. 1849. 19. 1850.
Mammillaria asterias Cels in Salm-Dyck, Cact. Hort. Dyck. 1849. 129. 1850.
Echinocactus glanduligerus Poselger, Allg. Gartenz. 21: 102. 1853.
Echinocactus brevimammus Poselger, Allg. Gartenz. 21: 102. 1853.
Coryphantha glanduligera Lemaire, Cactées 34. 1868.
Coryphantha brevimamma Lemaire in Förster, Handb. Cact. ed. 2. 394. 1885.
Cactus brevimamma Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus exsudans Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus glanduliger Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Subcylindric, 4 cm. in diameter; tubercles dull green, thick, ovate; glands in the axils of the tubercles pale yellow; spine-areoles somewhat tomentose, becoming naked; radial spines 6 or 7, 6 to 10 mm. long, slender, straight, spreading, yellow; central spine 1, erect, yellow but brown at tip, perhaps hooked; flowers yellow.

Type locality: Between Ixmiquilpan and Zimapan.

Distribution: Central Mexico.

All the synonyms cited above may or may not belong here. Our description is compiled mostly from Pfeiffer's.

Mammillaria curvata (Pfeiffer, Enum. Cact. 15. 1837) was given as a synonym of *Mammillaria exsudans*.

Illustrations: Monatsschr. Kakteenk. 23: 147; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 30, as *Mammillaria glanduligera*.

* This binomial is credited to Lemaire by Rümpler (Förster, Handb. Cact. ed. 2. 395. 1885), but as a synonym of *Mammillaria brevimamma exsudans*.


12. *Coryphantha erecta* Lemaire, Cactées 34. 1868.*Mammillaria erecta* Lemaire in Pfeiffer, Allg. Gartenz. 5: 370. 1837.*Mammillaria ceratocentra* Berg, Allg. Gartenz. 8: 130. 1840.*Echinocactus erectus* Poselger, Allg. Gartenz. 21: 126. 1853.*Cactus erectus* Kuntze, Rev. Gen. Pl. 1: 260. 1891.*Cactus ceratocentrus* Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Plant-body cylindric, yellowish green; axils of young tubercles white-woolly; tubercles obliquely conic, somewhat rhombiform at base; radial spines 8 to 14, subulate, ascending, yellowish; central spines 2, upper one short, lower one curved; flowers large, yellow; perianth-segments very narrow.

Type locality: Mexico.

Distribution: State of Hidalgo.

The plant described by Schumann has four central spines and may not belong to this species; his illustration answers it fairly well but does not show 4 centrals. We have recently examined specimens labeled *Mammillaria erecta* which were sent by Carl Ackerman, employed at the Huntington estate near Los Angeles, California; his plants grow in clumps



FIGS. 30 and 31.—*Coryphantha erecta*.

one meter in diameter; the larger branches are prostrate below, ascending or erect above, 3 dm. long; the spine-areoles are circular, white-felted when young; the spines are glossy yellow, the radials widely spreading; central spines often wanting or sometimes solitary, porrect, and shorter than the radials.

Mammillaria evarescentis, according to Lemaire (Cact. Aliq. Nov. 4. 1838), was a garden name improperly applied to this species.

The three names *Mammillaria evanescens*, *M. evarescens*, and *M. evarascens* were listed as synonyms of *M. erecta* by Förster (Handb. Cact. 243. 1846).

Illustrations: Schumann, Gesamtb. Kakteen 504. f. 82; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 7; Lemaire, Icon. Cact. pl. 10, as *Mammillaria erecta*.

Figure 30 is from a photograph of a plant collected by Dr. Rose in Mexico in 1906 (No. 1072a) and figure 31 is from a photograph of a plant growing in the Huntington collection in southern California which was made by Ernest Branton.

13. *Coryphantha elephantidens* Lemaire, Cactées 35. 1868.*Mammillaria elephantidens* Lemaire, Cact. Aliq. Nov. 1. 1838.*Echinocactus elephantidens* Poselger, Allg. Gartenz. 21: 102. 1853.*Cactus elephantidens* Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Simple, subglobose, up to 14 cm. high and 19 cm. broad; tubercles very large, somewhat flattened, obtuse, 4 to 5 cm. long, densely woolly in the axils; areoles elliptic, when young woolly, in age naked; spines 8, all radial, somewhat unequal, subulate, the longest about 2 cm. long, spreading, when young brownish with yellowish bases, black at apex; flowers large, rose-colored, 11 cm. broad; perianth-segments numerous, narrowly oblong, apiculate.

Type locality: Not cited.

Distribution: Central Mexico, but Nicholson's Dictionary of Gardening says Paraguay in error.

This is a very characteristic plant but we know it only from illustrations. Walter Mundt once offered it for sale but his supply has been exhausted; he gives a good illustration of it in a group of cacti printed on his letter heads and he writes us that this plant has a large carmine flower.

Schelle (Handb. Kakteenk. 238. 1907) gives *M. elephantidens spinosissima* Rebut, without synonymy or description.

Illustrations: Dict. Gard. Nicholson 4: 563. f. 33; Suppl. 516. f. 550; Förster, Handb. Cact. ed. 2. 397. f. 40; Hort. Univ. 1: pl. 33; Pfeiffer, Abbild. Beschr. Cact. 2 pl. 20; Rümpler, Sukkulanten 206. f. 117; Garden 1: 396; Lemaire, Icon. Cact. pl. 2 [not pl. 3]; Herb. Génér. Amat. II. 2: pl. 17; Palmer. Cult. Cact. 111; Ann. Rep. Smiths. Inst. 1908: pl. 14, f. 3; Goebel, Pflanz. Schild. 1: f. 34; Blanc, Cacti 68. No. 1224; Watson, Cact. Cult. 159. f. 60; Bergen in Rother, Praktischer Leitfaden Kakteen 5 ed. 1. 65; ed. 3. f. 38, as *Mammillaria elephantidens*.

14. *Coryphantha bumamma* (Ehrenberg).

Mammillaria bumamma Ehrenberg, Allg. Gartenz. 17: 243. 1849.

Mammillaria elephantidens bumamma Schumann, Keys Monogr. Cact. 43. 1903.

Globular or somewhat depressed; tubercles few, very large, rounded at apex, bluish green, very woolly in their axils when young but glabrate in age; spines 5 to 8, subulate, grayish brown, more or less recurved, 2 cm. long or more, all radial; flower large, yellow, 5 to 6 cm. broad; inner perianth-segments narrowly oblong, obtuse or retuse.

Type locality: Mexico.

Distribution: Mexico.

This plant is perhaps nearest *Coryphantha elephantidens*, to which it was referred as a variety, but the flowers are much smaller and nearly yellow. Mundt states that the flowers are smaller but bright rose with a dark stripe. His plant, however, is not now in his possession.

The plants are often much depressed, arising only a little above the surface of the ground, and are firmly anchored in the soil by a thick root, almost equal in diameter to that of the stem itself.

Dr. Rose made two collections in Mexico which we would refer here, one on the pedregal near Yautepec, Morelos (No. 8530), and the other at Iguala, Guerrero (No. 9320).

Illustration: Engler and Prantl, Pflanzenfam. 3^{6a}: 194. f. 67, as *Mammillaria bumamma*.

Plate v, figure 6, shows a plant collected by H. H. Rusby at Lemon Mountain, Guerrero, altitude 800 meters, July 28, 1910 (No. 4), which flowered in the New York Botanical Garden, September 11, 1911. Figure 29 is from a photograph showing a top view of a plant collected by Dr. C. Reiche at Iguala, Mexico, in 1921.

15. *Coryphantha robustispina* (Schott).

Mammillaria robustispina Schott in Engelmann, Proc. Amer. Acad. 3: 265. 1856.

Cactus robustispinus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria brownii Toumey, Bot. Gaz. 22: 253. 1896.

Cactus brownii Toumey, Bot. Gaz. 22: 253. 1896.

Stems solitary or clustered, globular or a little longer than thick, 5 to 15 cm. high, densely armed and almost hidden by the spines; tubercles large, 2.5 to 2.8 cm. long, arranged in 13 somewhat spiraled rows, fleshy, in age thickly set one against the other, becoming more or less dorsally flattened, pale, grayish green, narrowly grooved; radial spines 12 to 15, the 3 lower very stout, brownish,

the upper generally weaker, the 2 or 3 uppermost ones much weaker, clustered closely together and very pale, some of them sometimes crowded towards the center, the central spine solitary, very stout and erect or sometimes curved or even hooked, yellow, 3.5 cm. long; all the larger spines somewhat bulbous at base; flowers .5 to 6 cm. long, salmon-colored; ovary 20 to 25 mm. long, bearing 4 to 7 minute caducous scales; fruit narrowly oblong, 6 cm. long; seeds large, 3 mm. long, shining.

Type locality: Cited as Sonora in first publication of species; afterwards as south side of the Baboquivari Mountains in northern Sonora.

Distribution: Mountains of southern Arizona, southwestern New Mexico, and northern Sonora.

We have followed Mrs. K. Brandegee in referring *Mammillaria brownii* here, for not only do the original descriptions read much alike but the type localities for the two are in the same mountain range. *M. brownii* was described from a very small plant and differs considerably from mature individuals. Engelmann calls attention to the very large seeds, which he says are "larger than those of any other *Mammillaria* examined." He also states, "embryo with some albumen, curved; cotyledon foliaceous, approaching the structure of the seed of most *Echinocacti*."

Dr. Shreve reports that the flowers appear in the summer and the fruits, which follow, hold over the following winter, gradually drying up. The fruits do not open by a basal pore as in other related species.

We would refer here specimens from Lordsburg, New Mexico, and Bowie, Arizona, which, have heretofore been referred to *Mammillaria valida*, now *Coryphantha muehlenpfordtii*.

Illustrations: Bot. Gaz. 22: 254, as *Mammillaria brownii*; Cact. Journ. 1: 85; Cact. Mex. Bound. pl. 74, f. 8, as *Mammillaria robustispina*.

16. *Coryphantha connivens* sp. nov.

Globular or somewhat depressed, 8 to 10 cm. broad, somewhat woolly at the crown at flowering time but becoming glabrate; spines all radial but of two kinds; one kind 5 or 6, spreading or curved backward, subulate, horn-colored, the other 8 to 10, from upper part of spine-areole, clustered, erect, or toward top connivent, acicular, black at tip; flowers yellow, 6 to 7 cm. broad; perianth-segments narrowly oblong, acuminate; fruit greenish, oblong, 3 cm. long; seeds brown, oblong, 2 mm. long.

This species is common in the Valley of Mexico, especially on the pedregal. Dr. Rose collected it first in 1901 and again in 1905 and 1906; the type is his No. 8372 from near Tlalpam, collected in 1905. Dr. C. Reiche also collected it between Tacubaya and Santa Fe in 1922, and according to him the plant from this locality is the one referred to *Mammillaria pycnantha* by Schumann (Gesamtb. Kakteen 489. 1898).

The species is characterized by the peculiar clusters of spines in the upper angle of the areoles. A small plant was sent by O. Solis from Tlalpam in 1907, but it has fewer acicular spines than described above.

17. *Coryphantha pectinata* (Engelmann).

Mammillaria pectinata Engelmann, Proc. Amer. Acad. 3: 266. 1856.

Mammillaria pectinata cristata Hortus in Förster, Handb. Cact. ed. 2. 403. 1885.

Cactus pectinatus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Usually simple, globose, 3 to 6 cm. in diameter; tubercles usually arranged in 13 spirals; upper tubercles 10 to 12 mm. long, about twice as long as lower ones; areoles a little longer than broad; spines 16 to 24, all radial, those on lower areoles appressed and often a little recurved, those from upper part of upper areoles 12 to 18 mm. long, connivent over apex, yellowish white with black tips; flowers yellow, 5 cm. long; ovary 6 to 8 mm. long; fruit 12 mm. long.


Type locality: On the Pecos River in western Texas.

Distribution: Southern Texas and adjacent parts of Mexico.


Coulter and Schumann refer it to *Mammillaria radians* De Candolle, but it doubtless is a distinct species.

This plant is well illustrated by Engelmann and should be easily recognized. It appears to have been collected only rarely. The only representatives we have of it are flowers

1



2



1. *Coryphantha nickelsae*, from Monterey, Mexico.
2. *Neobesseya similis*, from Texas.

and a spine-cluster from the herbarium of J. W. Toumey, collected in his cactus garden at Tucson, June 12, 1896, and a small specimen from near the type locality obtained by Vernon Bailey, March 22, 1890, and more recently by Fisher at Langtry, Texas.

Illustrations: Cact. Journ. 1: 114; 2: 6; Dict. Gard. Nicholson Suppl. 514. f. 546; Förster, Handb. Cact. ed. 2. 402. f. 42; Rümpler, Sukkulente 204. f. 115; Journ. Hort. Home Gard. III. 46: 379; Cact. Mex. Bound. pl. 11; Watson, Cact. Cult. 169. f. 67; ed. 3. f. West Amer. Sci. 13: 40; Blanc, Cacti 73. No. 1459; Cassell's Dict. Gard. 2: 48; Remark, Kakteenfreund 15, as *Mammillaria pectinata*; Schelle, Handb. Kakteenk. 240. f. 158, as *M. radians impexicoma* [Schelle's illustration is the same as Engelmann's].

Figure 31a is from a photograph of a plant obtained by George L. Fisher near Langtry, Texas, in 1922.




FIG. 31a.—*Coryphantha pectinata*.




FIG. 31b.—*Coryphantha echinus*.

18. *Coryphantha nickelsae* (K. Brandegee).

Mammillaria nickelsae K. Brandegee, Zoe 5: 31. 1900.

Described as globular, densely cespitose, often 7 cm. high, pale green and glaucous; older plants becoming purplish; tubercles almost hidden by the overlapping spines, rather broad at base, low, not densely arranged; spines 14 to 16, all radial (a few forming a small fascicle at top of groove), slender, at first simply spreading but afterward bent back and interlaced with those of adjoining tubercles, 8 to 10 mm. long, at first yellowish at base with dark tips, but afterwards bleaching; flowers described as bright yellow, with a red center, 5 to 7 cm. broad; fruit nearly globular, 5 to 7 mm. long, green; seeds small, brown.

Type locality: Mexico, southward from Laredo, Texas.

Distribution: Northern Nuevo Leon, Mexico.

Plants collected by Robert Runyon in March 1921, on Mount La Mitra, near Monterey, which we believe should be referred here, deserve some detailed description. They grow in clusters of 4 to 12. From the axils of the lower tubercles near the surface of the ground numerous young plants or buds originate; the young spines are pale yellow, with reddish-brown tips, in age some bleaching white, others brownish to nearly black throughout; many of the first areoles have only radial spines but old plants often have one central spine 1.5 to 2 cm. long, from all the upper areoles; flowers large, light yellow; inner perianth-segments spreading, linear-lanceolate, acuminate; anthers bright yellow.

Plate III, figure 1, is from a photograph of the plant collected by Mr. Runyon, which was made at his home in Brownsville, Texas, September 15, 1921. Figure 32 is from a photograph of a specimen sent us by Dr. Richard E. Kunze in 1911.

19. *Coryphantha compacta* (Engelmann).

Mammillaria compacta Engelmann in Wislizenus, Mem. Tour North. Mex. 105. 1848.
Cactus compactus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Plants solitary, somewhat depressed, 3 to 6 cm. high, 5 to 8 cm. broad; tubercles in 13 rows, much crowded, 8 mm. long, sulcate above; radial spines 14 to 16, rigid, appressed, interwoven with adjacent ones, whitish, 10 to 20 mm. long; central spines usually wanting; flowers 2 cm. long and broad, yellow; fruit oval; seeds smooth and yellow.

Type locality: Cosihuiriachi, Chihuahua.

Distribution: Mountains of Chihuahua.

This species had long been known only from the original plant collected by Wislizenus, but in 1908 Dr. Rose visited the type locality, where he re-collected the plant, which later flowered at Washington.




FIG. 32.—*Coryphantha nickelsae*.

FIG. 33.—*Coryphantha compacta*.

The name *Coryphantha compacta* occurs in C. R. Orcutt's Circular to Cactus Fanciers, 1922.

Illustrations: Cact. Mex. Bound. pl. 74, f. 2 (seeds); Dict. Gard. Nicholson Suppl. 515. f. 548; Bull. U. S. Dept. Agr. Bur. Pl. Ind. 262: pl. 2, f. 1; Watson, Cact. Cult. ed. 2. 254. f. 95; ed. 3. 76. f. 35, as *Mammillaria compacta*.

Figure 33 is from a photograph of the plant collected by Dr. Rose.

20. *Coryphantha radians* (De Candolle).

Mammillaria radians De Candolle, Mém. Mus. list. Nat. Paris 17: 111. 1828.
Mammillaria impexicoma Lemaire, Cact. Aliq. Nov. 5. 1838.
Mammillaria daimonoceras Lemaire, Cact. Aliq. Nov. 5. 1838.
Mammillaria radians globosa Scheidweiler, Bull. Acad. Sci. Brux. 5: 494. 1838.
Mammillaria cornifera impexicoma Salm-Dyck, Cact. Hort. Dyck. 1849. 20. 1850.
Echinocactus corniferus impexicomus Poselger, Allg. Gartenz. 21: 102. 1853.
Echinocactus radicans Poselger, Allg. Gartenz. 21: 107. 1853.
Coryphantha daimonoceras Lemaire, Cactées 35. 1868.
Cactus radians Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus radians pectinoides Coulter, Contr. U. S. Nat. Herb. 3: 114. 1894.
Mammillaria radians impexicoma Schumann, Gesamtb. Kakteen 495. 1898.
Mammillaria radians daimonoceras Schumann, Gesamtb. Kakteen 496. 1898.

Solitary, globose, either obtuse or depressed at the top, 7.5 cm. in diameter; tubercles ovoid, large; axils of tubercles naked; areoles glabrate; spines all radial, 16 to 18, white or sometimes yel-

lowish, 10 to 12 mm. long, rigid, tomentose when young; flowers lemon-yellow, with outer segments tinged with red, about 10 cm. broad, the segments narrowly oblong to spatulate, acute, somewhat toothed toward the apex.

Type locality: Mexico.

Distribution: Central Mexico.

It is difficult to ascertain what the true *Mammillaria radians* of De Candolle really is. The type plant was described from specimens collected by Thomas Coulter, probably in eastern Mexico. We believe that specimens collected by Dr. Edward Palmer near San Luis Potosí, Mexico, represent the species as well as any plants we have yet seen; these, however, are cespitose as well as solitary. . The species seems nearest *Coryphantha compacta*.

Cactus radians pectinoides Coulter, based on Eschanzier's plant from San Luis Potosí (1891), we have not seen but suspect that it belongs here.




FIG. 34.—*Coryphantha radians*.




FIG. 35.—*Coryphantha sulcolanata*.

Mammillaria monoclova is only a garden name cited by Schumann (Gesamtb. Kakteen 495. 1898) as a synonym of this species.

Coryphantha impexicoma, credited to Lemaire, is given as a synonym of *Mammillaria cornifera impexicoma* Salm-Dyck by Rümpler (Förster, Handb. Cact. ed. 2. 414. 1885).

Illustrations: Blühende Kakteen 2: pl. 102; Tribune Hort. 4: pl. 139; Succulenta 5: 57, as *Mammillaria radians*; Monatsschr. Kakteenk. 15: 7, as *M. radians impexicoma*.

Figure 34 is from a photograph of the plant collected at San Rafael by Dr. Chaffey in 1910.

21. *Coryphantha sulcolanata* Lemaire, Cactées 35. 1868.

Mammillaria sulcolanata Lemaire, Cact. Aliq. Nov. 2. 1838.

Echinocactus sulcolanatus Poselger, Allg. Gartenz. 21: 102. 1853.

Mammillaria conimamma Linke, Allg. Gartenz. 25: 239. 1857.

Mammillaria cornimamma N. E. Brown, Gard. Chron. III. 2: 186. 1887.

Cactus sulcolanatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Subglobose, somewhat depressed, cespitose, , cm. high, 6 cm. thick or more; tubercles somewhat ,-angled at base, subconic above, their axils very woolly when young; spines 9 or 10, all radial, unequal, 12 to 16 mm. long, the lower and upper weaker and shorter than the lateral ones, brownish with black tips, but when young whitish yellow with purple tips; flowers large, 4 cm. long or more, widely spreading, 6 cm. broad or more; perianth-segments oblong, acute.

Type locality: Not cited, but Rümpler states that the plant was collected by Galeotti near Mineral del Monte, Hidalgo, in 1836.

Distribution: Mexico, perhaps Hidalgo, but definite range unknown.

Aulacothele sulcolanatum Monville (Lemaire, Icon. Cact. pl. 10. 1841-1847), referred here as a synonym, seems never to have been published.

Mammillaria retusa Scheidweiler is sometimes referred here also and the name has priority over *M. sulcolanata*, but we are treating it as distinct.

Echinocactus conimamma Linke was cited by Schumann (Monatsschr. Kakteenk. 5: 75. 18.) by mistake for *Mammillaria conimamma* Linke. *M. conimamma major* is listed by Haage (Cact. Kultur ed. 2. 179. 1900).

The name *Mammillaria sulcolanata macracantha* (Walpers, Repert. Bot. 2: 273. 1843) was without description.

Illustrations: Haage, Cact. Kultur ed. 2. 178, as *Mammillaria bumamma*; Blanc, Hints on Cacti 68. No. 1224, as *Mammillaria elephantidens*; Lemaire, Icon. Cact. pl. 10; Förster, Handb. Cact. ed. 2. 408. f. 45; Schelle, Handb. Kakteenk. 238. f. 154; Watson, Cact. Cult. 178. f. 72 ed. 3. f. 49; Deutsche Gärt. Zeit. 6: 65; Dict. Gard. Nicholson 4: 565. f. 40; Suppl. 518. f. 558, as *Mammillaria sulcolanata*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 2, as *Mammillaria conimamma*; Lemaire, Cactées 35. f. 2.

Figure 35 is from a photograph of the plant collected by Dr. Rose near Pachuca in 1905.




FIG. 36.—*Coryphantha retusa*.




FIG. 37.—*Coryphantha salm-dyckiana*.

22. *Coryphantha retusa* (Pfeiffer).

Mammillaria retusa Pfeiffer, Allg. Gartenz. 5: 369. 1837.
Cactus retusus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plants depressed-globose, 5 to 10 cm. in diameter, the top very woolly; tubercles rather large; areoles elliptic; spines 6 to 12, all radial, appressed, or even curved backward, yellowish to brownish, subulate, except 2 or 3 acicular ones at upper part of areoles; flowers central, yellow, about 3 cm. long; inner perianth-segments oblong, acute.

Type locality: Mexico.

Distribution: Oaxaca, Mexico.

We have referred to this species a plant common in Oaxaca, which answers the original description very well. It was collected by Pringle in 1894 (No. 5706) and by Conzatti in

1907, 1909, and 1920. It has also been sent us from the same region by O. Solis and B. P. Reko.

Figure 36 is from a photograph of a plant sent from Oaxaca by O. Solis in 1920.

23. *Coryphantha palmeri* sp. nov.

Plant-body globular; tubercles closely set in about 13 rows but not very regularly arranged, pale green, not very flaccid; radial spines 11 to 14, rather stout, spreading nearly at right angles to central one, yellowish; tips often blackish; central spine one, stout, terete, hooked at apex; young areoles very woolly; flowers central, pale yellow to nearly white, about 3 cm. long; outer perianth-segments linear-oblong, acute, brownish on broad mid-rib, entire, the inner yellow throughout, acuminate; stamens numerous; stigma-lobes, linear, cream-colored.

Collected by Dr. Edward Palmer on stony ridge near Durango, Mexico, and flowered in Washington, July 1906 (No. 557, type). Here seem to belong plants collected by Dr. Palmer at Agua Nueva, April 1905 (No. 561), and at Saltillo, October 1904 (No. 438), and July 1905 (No. 703), and also by F. E. Lloyd in Zacatecas, 1908 (No. 9).

24. *Coryphantha cornifera* (De Candolle) Lemaire, Cactées 35. 1868.

Mammillaria cornifera De Candolle, Mém. Mus. Hist. Nat. Paris 1: 112. 1828.

Mammillaria pfeifferana De Vriese, Tydschr. Nat. Geschr. 6: 51. 1839.

Mammillaria scolymoides Scheidweiler, Allg. Gartenz. 9: 44. 1841.

Mammillaria scolymoides longiseta Salm-Dyck, Cact. Hort. Dyck. 1849. 132. 1850.

Mammillaria scolymoides nigricans Salm-Dyck, Cact. Hort. Dyck. 1849. 132. 1850.

Echinocactus corniferus Poselger, Allg. Gartenz. 21: 102. 1853.

Echinocactus corniferus longisetus Poselger, Allg. Gartenz. 21: 102. 1853.

Echinocactus corniferus nigricans Poselger, Allg. Gartenz. 21: 102. 1853.

Echinocactus corniferus scolymoides Poselger, Allg. Gartenz. 21: 102. 1853.

Cactus corniferus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus pfeifferanus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus scolymoides Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plant solitary, globose, pale green; tubercles short, broad, somewhat imbricated, 12 cm. high; radial spines 16 or 17, grayish, 10 to 12 mm. long; central spine 1, stout, erect or subincurved, generally dark colored, 14 to 16 mm. long; flowers yellow, tinged with red, 7 cm. broad; inner perianth-segments oblanceolate, acuminate; fruit not seen.

Type locality: Mexico.

Distribution: Central Mexico.

We refer here a plant collected by Dr. Rose near San Juan del Rio, August 17, 1905.

Schumann referred *Mammillaria scolymoides* to *Mammillaria radians*, but its relationship is rather with *M. cornifera* as suggested by Schumann.

Mammillaria cornifera mutica Salm-Dyck (Cact. Hort. Dyck. 1849. 20. 1850), taken up afterwards as *Echinocactus corniferus muticus* by Poselger (Allg. Gartenz. 21: 102. 1853), was without description and to it was referred *Mammillaria radians* Hortus.

Illustrations: Schumann, Gesamtb. Kakteen 492. f. 81; Thomas, Zimmerkultur Kakteen 55; Bull. U. S. Dept. Agr. Bur. Pl. Ind. 262: pl. 1; Blühende Kakteen 3: pl. 125; Monatsschr. Kakteenk. 14: 73, as *Mammillaria cornifera*; Karsten and Schenck, Vegetationsbilder 2: pl. 20e, as *Mammillaria scolymoides*; Tydschr. Nat. Geschr. 6: pl. 1, f. 2, as *Mammillaria pfeifferana*.

Plate 11, figure 4, shows a plant collected by Dr. C. A. Purpus in Coahuila in 1905 which flowered in the New York Botanical Garden.

25. *Coryphantha salm-dyckiana* (Scheer).

Mammillaria salm-dyckiana Scheer in Salm-Dyck, Cact. Hort. Dyck. 1849. 134. 1850.

Mammillaria salm-dyckiana brunnea Salm-Dyck, Allg. Gartenz. 18: 394. 1850.

Echinocactus salm-dyckianus Poselger, Allg. Gartenz. 21: 102. 1853.

Cactus salm-dyckianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria delaetiana Quehl, Monatsschr. Kakteenk. 18: 59. 1908.

Plants either solitary or in clusters, nearly globular or sometimes club-shaped, 10 to 15 cm. in diameter, light green; tubercles rather short, closely set; radial spines about 15, spreading, slender, 10 to 15 mm. long, grayish or whitish; central spines 1 to 4, reddish to black, the upper ones when

present ascending and those near top of plant connivent, the lowest central stouter than others, 2 to 2.5 cm. long, porrect or curved downward; flowers large, 4 cm. long; outer perianth-segments greenish or tinged with red, the inner pale yellow; filaments greenish yellow; stigma-lobes 7.

Type locality: Near Chihuahua, Mexico.

Distribution: Common in the state of Chihuahua, Mexico.

Mammillaria salm-dyckiana was originally collected by John Potts near Chihuahua City and sent to Kew; its flowers and fruit were unknown. Schumann referred it as a synonym of *M. scheeri*, but we believe that it must be distinct and that *M. delaetiana* is the same. It was described from plants distributed by de Laet, who probably obtained them from C. R. Orcutt.

In 1908 Dr. E. Palmer collected some fine plants near Chihuahua City, from which our flower characters have been drawn.

Illustrations: Monatsschr. Kakteenk. 18: 59; 20: 92, as *Mammillaria delaetiana*.

Figure 37 is from a photograph of the plant collected by Dr. E. Palmer near Chihuahua City in 1908.

26. *Coryphantha pallida* sp. nov.

Plants either solitary or in clusters of about 10 or more, globular, 12 cm. in diameter or less, bluish green; tubercles in 13 rows, short and thick, closely set; radial spines 20 or more, white,



FIG. 38.—*Coryphantha pallida*.




FIG. 39.—*Coryphantha pycnantha*.

appressed; centrals usually , but sometimes more, the two upper more or less ascending, the lower porrect or curved downward, with tip black, or sometimes black throughout; flowers very large, often 7 cm. long and nearly as broad; outer perianth-segments narrow, greenish yellow, with a reddish stripe on back; inner perianth-segments pale lemon-yellow, broader than outermost, acuminate; ovary bearing a few narrow scales; stamens deep red, numerous; style yellow, longer than stamens; stigma-lobes; fruit greenish brown, 2 cm. long; seeds brown, shining, broader at apex than below.

Common in calcareous soil about Tehuacán, Mexico. Collected by J. N. Rose in 1901 (No. 5583, type), in '905 (Nos. 99972 and 10001), and in 1906. Living specimens were also obtained and these have flowered repeatedly in cultivation. It was also collected by C. G. Pringle in 1901 (No. 8573) and distributed as *Mammillaria pycnantha*?

In young plants the spines are not so numerous, the central spine is single, porrect, slightly curved, with black tips.

Figure 38 is from a photograph of the plant collected at the type locality in 1906.

27. *Coryphantha pycnacantha* (Martius) Lemaire, Cactées 35. 1868.

- ?*Mammillaria latimamma* De Candolle, Mém. Mus. Hist. Nat. Paris 17: 114. 1828.
Mammillaria pycnacantha Martius, Nov. Act. Nat. Cur. 16: 325. 1832.
 ?*Mammillaria acanthostephes* Lehmann, Allg. Gartenz. 3: 228. 1835.
Mammillaria arietina Lemaire * Cact. Aliq. Nov. 10. 1838.
Mammillaria scepontocentra Lemaire, Cact. Gen. Nov. Sp. 43. 1839.
Mammillaria arietina spinosior Lemaire, Cact. Gen. Nov. Sp. 94. 1839.
Mammillaria pycnacantha spinosior Monville in Salm-Dyck, Cact. Hort. Dyck. 1844. 14. 1845.
Mammillaria magnimamma arietina Salm-Dyck in Förster, Handb. Cact. 235. 1846.
Mammillaria winkleri Förster, Allg. Gartenz. 15: 50. 1847.
Mammillaria magnimamma lutescens Salm-Dyck, Cact. Hort. Dyck. 1849. 17, 121. 1850.
Echinocactus winkleri Poselger, Allg. Gartenz. 21: 102. 1853.
 ?*Echinocactus acanthostephes* Poselger, Allg. Gartenz. 21: 102. 1853.
Echinocactus pycnacanthus Poselger, Allg. Gartenz. 21: 102. 1853.
Mammillaria acanthostephes recta Hortus in Labouret, Monogr. Cact. 138. 1853.
 ? *Coryphantha acanthostephes* Lemaire, Cactées 35. 1868.
Cactus acanthostephes Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus latimamma Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus pycnacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus scepontocentrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus winkleri Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plant solitary, globular to cylindric, about 8 cm. high; tubercles broad, grooved above, glaucous-green; radial spines 10 to 12, slender, 10 to 16 mm. long; central spines about , stouter than the radials, about 25 mm. long, more or less curved backward, usually black; flowers from near center of plant, 25 mm. in diameter, yellowish; perianth-segments numerous, very narrow; stigma-lobes 5 or 6, white.

Type locality: Near the city of Oaxaca, Mexico.

Distribution: Oaxaca, Mexico.

The skeleton of the type of this species is preserved in the Munich Museum and Dr. Rose obtained a cluster of spines from this specimen in 1912.

Coryphantha pycnacantha has long been a desideratum. In September 1920 Professor Conzatti sent several small plants from near the type locality. In these, the radial spines are white, the centrals (3) are nearly black, and all more or less curved backward. In the center of the plant a quantity of white wool is developed, so abundant that it can be gathered for commercial use. With the specimens of Professor Conzatti are samples of the wool with an inquiry as to its value as a fiber.

Mammillaria magnimamma spinosior Lemaire (Salm-Dyck, Cact. Hort. Dyck. 1844. 12. 1845) was not described at the place here cited. Labouret afterwards refers it as a synonym of *M. magnimamma lutescens*.

Mammillaria cephalophora Salm-Dyck (Cact. Hort. Dyck. 1849. 137. 1850; *Echinocactus cephalophorus* Poselger, Allg. Gartenz. 21: 102. 1853; *Cactus cephalophorus* Kuntze, Rev. Gen. Pl. 1: 260. 1891) was a new name for *Melocactus mammillariaeformis* † Salm-Dyck (Allg. Gartenz. 4: 148. 1836). It was first described as a *Melocactus* (because of its woolly crown), but it seems to be more like a *Coryphantha*. Its exact origin in Mexico seems to be unknown and the flowers had not been described up to 1850. Pfeiffer stated that the seeds obtained from a dead plant were similar to those of *Mammillaria coronaria*. Schumann discussed it under *M. pycnacantha* in a note. Hemsley (Biol. Centr. Amer. Bot. 1: 502) listed it as a *Melocactus*.

Mammillaria pycnacantha scepontocentra Monville (Labouret, Monogr. Cact. 136. 1853) belongs here by implication.

Mammillaria magnimamma Otto was referred by De Candolle (Mém. Cact. 17. 1834) to his *M. latimamma*, now referred to this species.

* Schumann refers this name as a synonym of *Mammillaria centricirrha*.

† Schumann spells this name *Melocactus mamillariiformis*.

Echinocactus radiatus Hortus Belg. was referred as a synonym of *Mammillaria pycnacantha* by Pfeiffer (Enum. Cact. 180. 1837).

Illustrations: Nov. Act. Nat. Cur. 16: pl. 17; Loudon, Encycl. Pl. ed. 3. 1379. f. 19387; Abh. Bayer. Akad. Wiss. München 2: pl. 3; Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 26; Curtis's Bot. Mag. 69: pl. 3972, as *Mammillaria pycnacantha*.

Figure 39 is reproduced from the first illustration cited above; a spine-cluster is also shown.

28. *Coryphantha echinus* (Engelmann).

Mammillaria echinus Engelmann, Proc. Amer. Acad. 3: 267. 1856.

Cactus echinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Mammillaria radians echinus Schumann, Gesamtb. Kakteen 496. 1898.

Solitary, globose to subconic, 3 to 5 cm. in diameter, almost hidden under the closely appressed spines; areoles orbicular or a little longer than broad; radial spines numerous, white, so to 16 mm. long; central spines 3 or 4, the 3 upper erect or connivent over the apex, the lower one porrect on side of plant, erect near top, subulate, straight, 1.5 to 2.5 cm. long, often blackish; flowers 2.5 to 5 cm. long, yellow; outer perianth-segments linear-lanceolate; inner perianth-segments 20 to 30, narrow; stigma-lobes about 12; fruit oblong, 12 mm. long.

Type locality: On the Pecos River, Texas.

Distribution: Western Texas.

The flowers with the type plant seem to have been shriveled, for Engelmann describes them as large, apparently about 1½ or 2 inches long; in a later description he states that they are yellow. This species is very rare in collections and we have seen no flowers of it. All the illustrations cited below are based on the figure in the Mexican Boundary Survey.

The name *Coryphantha echinus* occurs in C. R. Orcutt's Circular to Cactus Fanciers, 1922.

Illustrations: Cact. Mex. Bound. pl. 10; Dict. Gard. Nicholson 4: 562. f. 32; Suppl. 515. f. Watson, Cact. Cult. 157. f. 59; ed. 3. f. 37; Förster, Handb. Cact. ed. 2. 404. f. 43; Blanc, Cacti 68. f. 1228, as *Mammillaria echinus*; Schelle, Handb. Kakteenk. 240. f. 159, as *M. radians echinus*.

Figure 31b is from a photograph of a plant obtained by George L. Fisher near Langtry, Texas, in 1922.

29. *Coryphantha durangensis* (Range).

Mammillaria durangensis Runge in Schumann, Gesamtb. Kakteen 478. 1898.

Plants solitary or in small clusters, short-cylindric, so cm. long or less, somewhat glaucous; tubercles rather prominent, in 5 or 8 series, somewhat compressed dorsally, very woolly in the axils; radial spines 6 to 8, acicular, spreading, 1 cm. long or less; central spine solitary, often erect, those of uppermost areoles connivent, black; flowers very small, about 2 cm. long, when fully expanded 2.5 to 4 cm. broad; outer perianth-segments dark purple or with only a purple stripe down center; inner perianth-segments cream-colored to pale lemon-yellow; filaments cream-colored, about length of style; style and stigma-lobes cream-colored, the latter 5, linear and curved backward; fruit globular, 5 to 8 mm. in diameter, naked, greenish; seeds brown, about 1 mm. broad.

Type locality: Villa Lerdo, Durango, Mexico.

Distribution: Northern Mexico.

Dr. E. Chaffey has collected this plant for us several times at the type locality, but it does not survive long under glass. In 1911 he found a cristate form with the lobes flattened like the joints of an *Opuntia*, bearing flowers along the edges.

This is *Mammillaria compressa* of Hildmann's Catalogue, according to Schumann (Gesamtb. Kakteen 479. 1898).

Illustration: Wiener Ill. Gart. Zeit. 29: 411. f. 105, as *Mammillaria radians*.

Plate v, figure 4, shows a plant sent by Dr. Chaffey from the type locality in 1918, which flowered in the New York Botanical Garden, April 8, 1918. Figure 40 is from a photograph of a potted plant sent by Dr. Chaffey in 1910 which flowered in Washington; figure 41 is from a photograph of another plant sent by Dr. Chaffey in 1910.

30. *Coryphantha chlorantha* (Engelmann).

Mammillaria chlorantha Engelmann in Rothrock, Rep. U. S. Geogr. Surv. 6:127. 1878.

Cactus radiosus chloranthus Coulter, Contr. U. S. Nat. Herb. 3: 121. 1894.

Mammillaria radiosus chlorantha * Schumann, Gesamtb. Kakteen 481. 1898.

Plant cylindric, sometimes 20 to 25 cm. high, 8 cm. in diameter; tubercles closely set and entirely hidden by the densely matted spines; flowers small, 35 mm. broad; outer perianth-segments ciliate; inner perianth-segments yellow or greenish yellow, linear-lanceolate, acute; stigma-lobes white; fruit central, green, 2.5 cm. long, juicy, bearing 5 or 6 scales near top; seeds brown, flattened, 1.5 mm. long, reticulated.


Type locality: Southern Utah, east of Saint George.

Distribution: Southern Utah, western Arizona, central Nevada, and eastern southern California.

Mammillaria utahensis Hildmann, cited by Schumann (Gesamtb. Kakteen 481. 1898) as a synonym of *M. radiosus*, may have been based on this plant.

Illustrations: Förster, Handb. Cact. ed. 2. 328. f. 33; Gartenflora 32: 87; Deutsche Gärt. Zeit. 7: 53, as *Mammillaria chlorantha*; Schelle, Handb. Kakteenk. 236. f. 151, as *M. radiosus chlorantha*.

Plate v, figure 7, is from a plant collected by I. Tidestrom at the type locality in 1919, which flowered in the New York Botanical Garden, May 27, 1919. Figure 42 is from a photograph of a plant collected by Major E. A. Goldman in Prospect Valley, Arizona.



FIGS. 40 and 41.—*Coryphantha durangensis*.

31. *Coryphantha vivipara* (Nuttall) Britton and Rose in Britton and Brown, Illustr. Fl. ed. 2. 2: 571. 1913.

Cactus viviparus Nuttall, Fraser's Cat. No. 22. 1813.

Mammillaria vivipara Haworth, Suppl. Pl. Succ. 72. 1819.

Mammillaria radiosus Engelmann, Bost. Journ. Nat. Hist. 6: 196. 1850.

Echinocactus radiosus Poselger, Allg. Gartenz. 21: 107. 1853.

Echinocactus viviparus Poselger, Allg. Gartenz. 21: 107. 1853.

Mammillaria vivipara vera Engelmann, Proc. Amer. Acad. 3: 269. 1856.

Mammillaria vivipara radiosus Engelmann, Proc. Amer. Acad. 3: 269. 1856.

Mammillaria vivipara radiosus Engelmann, Cact. Mex. Bound. 15. 1859, as subspecies.

Cactus radiosus Coulter, Contr. U. S. Nat. Herb. 3: 120. 1894.

Mammillaria hirschtiana Haage, Monatsschr. Kakteenk. 6: 127. 1896.

Coryphantha radiosus Rydberg, Fl. Rocky Mountains 581. 1917.

* Schumann credits this trinomial to Engelmann at the place here cited, although we believe that Engelmann never used it.

Plants solitary or in clusters forming mounds 3 to 6 dm. in diameter, globular, with prominent tubercles; areoles large, woolly; radial spines about 16, rather delicate, radiating, white; centrals to 6, divergent, much stouter, brownish, swollen at base; ovary green, naked; outer perianth-segments greenish; inner ones somewhat pinkish, long-ciliate; innermost perianth-segments pinkish purple, narrow, acuminate, entire, spreading; filaments much shorter than the segments, pinkish, but paler below; style greenish to purple above, longer than the stamens; stigma-lobes linear, purple, about 8, apiculate; fruit green when mature, juicy, nearly globular, 1.5 cm. in diameter, with several (sometimes 5 or 6) small ciliate scales scattered over its surface; seeds light brown, 1.5 mm. long.

Type locality: "Near the Mandan towns on the Missouri, lat. near 49°."

Distribution: Manitoba to Alberta, Kansas, south to northern Texas and Colorado.

The group to which *Coryphantha vivipara* belongs has always been very puzzling. Dr. Engelmann, our greatest authority on this group, was sometimes of one opinion




FIG. 42.—*Coryphantha chlorantha*.




FIG. 43.—*Coryphantha Neo-mexicana*.

and sometimes of another. Schumann rejected the specific name *vivipara* of Haworth for this plant since he thought that it was not the same as the *vivipara* of Engelmann, but in this he must be wrong, for *Mammillaria vivipara* Haworth was based upon *Cactus viviparus* Pursh, a name previously used by Nuttall, and both Pursh's and Nuttall's descriptions were based on the specimens collected by Nuttall in "Upper Louisiana" in 1812. This is undoubtedly the plant which Engelmann had in mind and which he called variety *vera*. We have not seen the type, but Pursh stated that he had seen flowers in Lambert's Garden.

Engelmann's remarks regarding the variability of the species are interesting. In the Proceedings of the American Academy (3: 269) he says:

"The extreme forms are certainly very unlike one another, but the transitions are so gradual that I can not draw strict limits between them."

Coryphantha vivipara and the three following species are closely related.

This plant is a day bloomer, and according to Engelmann the flowers become fully expanded about one o'clock in the afternoon.

Hooker in Curtis's Botanical Magazine (pl. 7718) figures and describes a plant purchased from D. M. Andrews of Boulder, Colorado, in which all the spines are brown, the flower is rose-red, and the stigma-lobes are linear and white.

Mammillaria montana is described briefly and figured (f. 1399) by Blanc in Hints on Cacti, p. 72. It is also described and figured by Darel (Illustr. Handb. Kakteen 96. f. 81), who says that it comes from Montana and Utah. It is illustrated by Haage (Cact. Kultur ed. 2. 187). It is apparently the same as *Coryphantha vivipara*.

Illustrations: Cycl. Amer. Hort. Bailey 2: f. 1356; Stand. Cycl. Hort. Bailey 4: f. 2315; Tribune Hort. 4: pl. 140; Curtis's Bot. Mag. 126: pl. 7718; De Laet, Cat. Gén. f. 43; Cact. Mex. Bound. pl. 74, f. 3 (seed); Meehan's Monthly 9: pl. 9, as *Mammillaria vivipara*; Clements, Rocky Mountain Flow. pl. 32, f. 7; Clements, Fl. Mount. Plain pl. 32, f. 7; Britton and Brown, Illustr. Fl. 2: 462. f. 2526, as *Cactus viviparus*; Monatsschr. Kakteenk. 3: 132; Schelle, Handb. Kakteenk. 236. f. 150; Floralia 42: 375, as *Mammillaria radiosa*; Cact. Mex. Bound. pl. 74, f. 4 (seed), as *Mammillaria radiosa texana*; Cact. Mex. Bound. pl. 74. f. 4 (seed), as *M. radiosa borealis*; Britton and Brown, Illustr. Fl. ed. 2. 2: f. 2985.

32. *Coryphantha neo-mexicana* (Engelmann).

Mammillaria vivipara radiosa neo-mexicana Engelmann, Proc. Amer. Acad. 3: 269. 1856.

Mammillaria radiosa neo-mexicana Engelmann, Cact. Mex. Bound. 64. 1859.

Mammillaria radiosa borealis Engelmann, Cact. Mex. Bound. 68. 1859.

Mammillaria radiosa texana Engelmann, Cact. Mex. Bound. 68. 1859.

Cactus radiosus neo-mexicanus Coulter, Contr. U. S. Nat. Herb. 3: 120. 1894.

Cactus neo-mexicanus Small, Fl. Southeast. U. S. 812. 1903.

Mammillaria neo-mexicana A. Nelson in Coulter and Nelson, Man. Bot. Rocky Mountains 327. 1909.

Plants usually solitary, globular to short-oblong, 8 to 12 cm. long, the whole body usually hidden under a mass of spines; radial spines numerous, acicular, usually white; central spines several, much stouter than the radials, pale below, brown or black towards top; flowers 4 to 5 cm. broad when fully expanded; outer perianth-segments greenish or the ones nearer center purplish, ciliate; inner perianth-segments broadly linear, acuminate and apiculate, more or less serrate above; filaments greenish, much shorter than perianth-segments; stigma-lobes extending beyond filaments, white, obtuse, not apiculate as in *Coryphantha vivipara*; fruit 2.5 cm. long, green, juicy, naked except a few hairy scales near top, capped by withered perianth, depressed at apex.

Type locality: Western Texas to New Mexico, doubtless at El Paso.

Distribution: Western Texas, New Mexico, and northern Chihuahua.

The distribution of this species can not be stated at present very definitely. It may be that some of the plants from northern New Mexico, especially those found in the mountains, may better be referred to *C. vivipara*, and the same is true of some of the plants from Texas. It is probable that the plants from central Texas and perhaps northwestern Texas may all be referred to *C. vivipara*. We have no Mexican plants before us but we have plants from El Paso, just over the Mexican Boundary line. Just how far south the species extends we do not know. We have greatly restricted the range from that given by Coulter in the Contributions from the U. S. National Herbarium (3: 120. 1894).

Illustrations: Gartenwelt 4: 159; Cact. Mex. Bound. pl. 13; Förster, Handb. Cact. ed. 2. 304. f. 30, as *Mammillaria radiosa neo-mexicana*; Watson, Cact. Cult. 181. f. 73; ed. 3. f. 50; Dict. Gard. Nicholson 4: 566. f. 41, as *Mammillaria vivipara radiosa*; Dict. Gard. Nicholson Suppl. 517. f. 554, as *Mammillaria radiosa*; Cact. Mex. Bound. pl. 74 (seed), as *Mammillaria borealis*.

Plate II, figure 1, shows a plant sent from Canutillo, Texas, by Mrs. S. L. Pattison in 1920; figure 1a shows the fruit. Figure 43 is from a photograph of a plant collected by Dr. Rose near Albuquerque, New Mexico, in 1908.

33. *Coryphantha arizonica* (Engelmann).

Mammillaria arizonica Engelmann, Bot. Calif. 1: 124. 1876.

Cactus radiosus arizonicus Coulter, Contr. U. S. Nat. Herb. 3: 121. 1894.

Mammillaria radiosa arizonica Schumann, Gesamtb. Kakteen 481. 1898.


Sometimes cespitose, forming large clumps a meter broad; each head globose to ovoid, 7.5 to 10 cm. in diameter; tubercles about 2.5 cm. long, cylindric, ascending, deeply grooved; spines

numerous, straight, rigid; radial spines 55 to 20, 10 to 30 mm. long, whitish; inner spines 3 to 6, stouter than the radial ones, deep brown above; flowers large, 5 to 7 cm. broad, rose-colored; outer perianth-segments 30 to 40, linear-subulate, with fimbriate margin; inner perianth-segments 40 to 50, lanceolate-linear, attenuate; stigma-lobes 8 to 50, white; fruit oval, green; seeds compressed, light brown, pitted.

Type locality: Northern Arizona.

Distribution: Northern Arizona, especially along the Upper River of the Grand Canyon, and perhaps also in southern Utah.*

Mammillaria arizonica Engelm., when first described, was a complex. Engelm. states that it was found "on rocky and sandy soil in northern Arizona from the Colorado eastward (Coues, Palmer, F. Bischoff) and into southern Utah (J. E. Johnson); probably in southeastern California." Engelm. afterwards described Johnson's plant from Utah as *M. chlorantha* and the California plant is doubtless his *M. deserti*. We have in the U. S. National Herbarium Palmer's specimen from Arizona but we have not seen the plant of Coues nor of Bischoff.



FIGS. 44 and 45.—*Coryphantha deserti*.

The northern range of this species is very uncertain. Engelm. extended it into southern Utah.

Plate v, figure 5, shows a plant sent by M. A. H. Spencer from the Grand Canyon, Arizona, in May 1907, which afterwards flowered in Washington.

34. *Coryphantha deserti* (Engelmann).

Mammillaria deserti Engelmann, Bot. Calif. 2: 449. 1880.

Cactus radius deserti Coulter, Contr. U. S. Nat. Herb. 3: 121. 1894.

Cactus radius alversonii Coulter, Contr. U. S. Nat. Herb. 3: 522. 1894.

Mammillaria alversonii Zeissold, Monatsschr. Kakteenk. 5: 70. 1895.

Mammillaria radiosa alversonii Schumann, Gesamtb. Kakteen 481. 1898.


Mammillaria radiosa deserti Schumann, Gesamtb. Kakteen 481. 1898.

Solitary or cespitose, usually cylindric, sometimes 2 dm. high, 6 to 9 cm. in diameter, densely covered with spines; radial spines white except



FIG. 46.—Flower of *C. deserti*.

* Our Utah reference is based on some detached flowers collected by M. E. Jones and a barren plant sent by Dr. C. D. Marsh in 1922. Both collections came from above Salina.



Coryphantha aggregata, from Arizona.

at tip, spreading; central spines several, sometimes as many as 14, much stouter than the radials, slightly spreading, those toward top of plant connivent, black or bluish black in their upper half, shading into red, nearly white at base; flowers 3 cm. long and nearly as broad when expanded, light pink, opening in bright sunlight; scales and outer perianth-segments ciliate; inner perianth-segments narrow, acute.

Type locality: Ivanpah, California.

Distribution: Deserts of southern California and southern Nevada.

This species is characterized by its stiff spines, with bluish-black tips shading into red, and is known in southern California as fox-tail cactus. The original description of *Mammillaria deserti* states that the flowers are straw-colored, tipped with pink, and this suggests *Coryphantha chlorantha* but we believe that it belongs with *Mammillaria alversonii*, which certainly has pinkish flowers, and since the name *deserti* is older than *alversonii* it is substituted for it.

Illustrations: Cact. Journ. 1: pl. for February, in part; Alverson's Cat. pl. facing 8, as *Mammillaria alversonii*; Schumann, Gesamtb. Kakteen 480. f. 79, as *M. radiosa alversonii*.

Figure 44 is from a photograph of a single plant sent by E. C. Rost; figure 45 is from a photograph of a clump photographed by E. C. Rost in its natural surroundings; figure 46 shows a flower taken from Mr. Rost's plant.




FIG. 47.—*Coryphantha aggregata*.




FIG. 48.—*Mammillaria recurvispina*.

35. *Coryphantha aggregata* (Engelmann).

Mammillaria aggregata Engelmann in Emory, Mil. Recon. 157. 1848.

Cereus aggregatus Coulter, Contr. U. S. Nat. Herb. 3: 396. 1896, as to name.

Echinocereus aggregatus Rydberg, Bull. Torr. Club 33: 146. 1906, as to name.

Plants solitary or cespitose, globular to short-oblong, very spiny; radial spines numerous, stouter than those of *Coryphantha vivipara*, white, often with brown tips, appressed; central spines several, stout, all erect and appressed or one often porrect, those towards top of plant connivent; flowers very large and showy, purplish, 5 to 7 cm. broad; outer perianth-segments ciliate; inner perianth-segments narrowly oblanceolate, often 6 mm. broad, acute, apiculate; stigma-lobes 8 to 10, elongated, white; fruit green, oblong, 2 to 2.5 cm. long, naked or occasionally bearing a small scale on the side, juicy; seeds dark brown, 2 mm. long.

Type locality: Head waters of the Gila.

Distribution: Western New Mexico, southeastern Arizona, and northern Sonora.

Mammillaria aggregata came from the headwaters of the Gila. The type was not preserved and is known only from a drawing reproduced in Emory's report. There has

been much discussion about the identity of the plant; Coulter transferred it to *Cereus*, referring to it *Cereus coccineus* and *C. phoeniceus* and assigning to it a wide range, Colorado to San Luis Potosí. Rydberg transferred the name to *Echinocereus* but applied it to the same group of plants described by Coulter. A careful restudy of the original illustration and Engelmann's description and a restudy of all the cacti of similar habit in the southwest leads us to a different conclusion from that reached by Dr. Coulter and Dr. Rydberg. Engelmann, who described it as a *Mammillaria*, says that it appears to be allied to *M. vivipara*, and this we believe is its true relationship. A *Mammillaria* from the region about Flagstaff often forms the great clusters mentioned by Engelmann, and while we believe that it differs from the one found in northern Arizona it is certainly a near ally, probably representing the closely related species from southeastern Arizona and southwestern New Mexico which has often passed as *M. arizonica*.

Engelmann referred a specimen which he had from Sonora to his variety *Mammillaria vivipara neo-mexicana* with the remark that it was "a form with more spines than any other."

Plate IV shows a clump sent by Mrs. Ruth C. Ross from near Aravaipa, Arizona, in July 1922. Figure 47 is from a photograph of a single plant obtained by Dr. Rose near Benson May 1, 1908, which afterwards flowered in Washington.

36. *Coryphantha cubensis* Britton and Rose, Torreyia **12**: 15. 1912.

Mammillaria urbaniana Vaupel, Monatsschr. Kakteenk. **22**: 65. 1912.

Plants depressed-globose, tufted, 2 to 3 cm. broad, pale green; tubercles numerous, vertically compressed, 6 to 7 mm. long, 4 to 5 mm. wide, about 3 mm. thick, grooved on upper side from apex to below middle, the groove very distinct; spines about 10, whitish, radiating, acicular but weak, 3 to 4 mm. long, those of young tubercles subtended by a tuft of silvery white hairs, 1.5 mm. long; flowers pale yellowish green, 16 mm. high, the segments acute; filaments, style, and stigma-lobes yellowish; fruit red, less than 1 cm. long, naked; seeds black, somewhat angled.

Type locality: Among stones in barren savanna, southeast of Holguin, Oriente, Cuba.

Distribution: Type locality and vicinity.

This species is very inconspicuous and perhaps for that reason is rare in collections. It has only twice, to our knowledge, been collected, both times by Dr. J. A. Shafer, once in 1909 (No. 2946) and again in 1912 (No. 12432), who gave a short account of its discovery in the journal of the New York Botanical Garden (No. 155). He states that it barely protrudes through the layer of broken stones that filled the interstices between the larger rocks; that the largest plants were scarcely an inch in diameter, one of them bearing a small yellowish flower. It lives only a short time in greenhouse cultivation.

On account of the name *Mammillaria cubensis* Zuccarini (Labouret, Monogr. Cact. 59. 1853) Vaupel gave a new specific name to the plant when he transferred it from *Coryphantha*.

Plate V, figure 1, shows the plant collected by Dr. Shafer in 1912 which flowered in the New York Botanical Garden in July of the same year; figure 1a shows the fruit and figure 1b shows a tubercle from the same plant.

37. *Coryphantha sulcata* (Engelmann).

*Mammillaria sulcata** Engelmann, Bost. Journ. Nat. Hist. **5**: 246. 5845.

Mammillaria strobiliformis Mühlenpfordt, Allg. Gartenz. **16**: 19. 1848. Not Engelmann, 1848.

Mammillaria calcarata Engelmann, Bost. Journ. Nat. Hist. **6**: 195. 1850.

Coryphantha calcarata Lemaire, Cactées 35. 1868.

Cactus calcaratus Kuntze, Rev. Gen. Pl. **1**: 259. 1891.

Cactus scolymoides sulcatus Coulter, Contr. U. S. Nat. Herb. **3**: 116. 1894.

Mammillaria radians sulcata Schumann, Gesamtb. Kakteen 496. 1898.

Cactus sulcatus Small, Fl. Southeast. U. S. **812**. 1903.

Cespitose, 8 to 12 cm. in diameter; tubercles rather large, 10 to 12 mm. long, somewhat flattened, soft; radial spines acicular, straight, white; central spines several, one somewhat stouter than

* Förster (Handb. Cact. 255. 5846) credits such a name to Pfeiffer but it is without description.



M. E. Eaton del. 1 to 4, 6, 7

A. A. Newton del. 5

A. Hoar & Co. Baltimore

1. Flowering plant of *Coryphantha cubensis*.
- 1a. Fruit of same.
- 1b. Tubercle of same.
2. Flowering plant of *Neomammillaria confusa*.
3. Flowering plant of *Neomammillaria geminispina*.

4. Top of flowering plant of *Neomammillaria confusa*.
5. Flowering plant of *Coryphantha arizonica*.
6. Flowering plant of *Coryphantha bumamma*.
7. Flowering plant of *Coryphantha chlorantha*.

the others, porrect or slightly curved outward, others erect; flowers several, from near center of plant, 5 cm. in diameter or more, yellow, with a red center; inner perianth-segments lanceolate, apiculate; filaments reddish; style greenish yellow, exserted beyond stamens; stigma-lobes 7 to 10, yellow, notched at apex;* fruit oblong, greenish; seeds oblong, shining, dark brown.

Type locality: Industry, Texas.

Distribution: Southern Texas.

The herbarium sheets of this plant, sent us from the Missouri Botanical Garden, contain seeds, fruit, and style. Dr. Coulter speaks of seeing the spines of the type.

The name *Mammillaria sulcata*, first given by Engelmann, was changed by him to *M. calcarata* on account of *M. sulcata* Pfeiffer, but this was a later name and hence can not replace Engelmann's first one.

This species was collected by Lindheimer at Industry, Texas, growing with *Mammillaria similis*, but while the two are similar in habit, this plant differs from *M. similis* in having green fruit and brown oblong seeds instead of red fruit and black globose seeds, as well as in other ways. It has not been collected much in recent years and its characters and




FIG. 49.—*Coryphantha sulcata*.

range have been involved with other species. Miss Ellen D. Schulz sent us plants from San Antonio, Texas, in June 1921, and Robert Runyon sent us plants and photographs in 1922, which have enabled us to restudy the species in connection with its type now kept in the Engelmann Herbarium in the Missouri Botanical Garden.

Mammillaria goernгии was given by Haage (Cact. Kultur ed. 2. 183. 1900) as a new name for *M. calcarata*.

Illustrations: Cact. Mex. Bound. pl. 74. f. 1, as *Mammillaria calcarata*; Monatsschr. Kakteenk. 27: 65, as *Mammillaria radians sulcata*.

Plate x, figure 1, shows a plant photographed by Robert Runyon at Sabinal, Texas, April 28, 1922. Figure 49 is from a photograph of four fruits sent by Professor Albert Ruth, of Polytechnic, Texas, in 1922.

* Whether this is a constant character we do not know, but we have observed it in three flowers, all from the same plant. It has not been noted before in any other species of *Coryphantha*.

PUBLISHED SPECIES, PERHAPS OF THIS GENUS.

MAMMILLARIA CALOCHLORA Hortus, Monatsschr. Kakteenk. **26**: 167. 1916; **2**: 133. 1917.

This seems undoubtedly a species of *Coryphantha*, but we have not been able to identify it. There is considerable confusion regarding this plant, as the following note from Meyer would indicate:

"I have gotten Mr. Quehl to send me the flower of *Mammillaria calochlora* Hort. and I see that this also agrees exactly with the flower of Grässner's *M. delaetiana*. As third and last I have now gotten Mr. de Laet to send me also a little plant of equal size of his genuine *M. delaetiana* Quehl and this one is entirely different from the two others in form and color of the body, areoles, and spines."

We have a small specimen and a photograph sent us by L. Quehl in 1921.

MAMMILLARIA CORDIGERA Heese, Gartenflora **59**: 445. 1910.

Short-cylindric, 6 cm. high, 4.5 cm. in diameter; tubercles 4-angled, broader than long, grooved above; spine-areoles longer than broad; radial spines 4 to 15, white, spreading; central spines 4, erect, curved if not hooked at apex, 15 mm. long; flowers and fruit unknown.

Type locality: Not cited.

Distribution: Doubtless Mexico.

This species we know only from descriptions and illustration. The illustration is so much like that of *Mammillaria bombycina* that we at first were inclined to combine them. From the observations of others there seem to be important technical differences which separate them, not only specifically but also generically. It may prove to be a synonym of *C. sulcolanata*, for we have recently examined a skeleton sent us by Bödeker which resembles very much the plants collected by Rose in Hidalgo, Mexico, which we have already referred to that species.

Illustration: Gartenflora **59**: f. 50, as *Mammillaria cordigera*.

MAMMILLARIA CORNUTA Hildmann in Schumann, Gesamtb. Kakteen **496**. 1898.

Simple, grayish green, somewhat depressed, 4 to 5 cm. high, 6 to 8 cm. in diameter; tubercles spiraled, in 5 to 8 series; radial spines 5 to 7, subulate, straight or somewhat curved, white, 4 to 8 mm. long; central spine solitary, horn-colored; flowers said to be rose-colored; fruit unknown.

Type locality: Mexico.

From the description it is difficult to identify this species; its rose-colored flowers suggest a relationship with *Coryphantha elephantidens* but its spine-clusters are differently described.

MAMMILLARIA POTOSIANA Jacobi, Allg. Gartenz. **24**: 92. 1856.

Erect, cylindric, light green; tubercles conical, triangular at base, bearing 2 yellow glands in their axils; radial spines 15 or 16, subulate, equal or nearly so, 6 mm. long; central spine solitary, porrect but somewhat incurved at apex, subulate, 10 to 12 mm. long; flowers yellow.

Type locality: San Luis Potosí, Mexico.

Jacobi comments on the species as follows:

"Comptroller Shafer in Münster received this beautiful plant in a shipment of plants from San Luis Potosí in Mexico, under the name of *Mammillaria raphidacantha*. From the given description it is adequately clear that the plant considered is another and undescribed one. The form of the tubercles as well as the number and form of the spines is other evidence, also the grooves upon the upper sides of the tubercles which are always present in the case of *M. raphidacantha* are here lacking throughout.

"The stem of the plant is cylindrical, dark green, finely punctate with white dots; tubercles conical, 3-angled at the base, gradually flattened above; axils sinuate with 2 yellow glands, inclosed by a ring of yellowish-white tomentum; areoles terminal, oval, the younger ones whitish tomentose, later naked; radial spines 15 or 16, radiating, somewhat recurved, needle-formed, two-colored. In older plants there appears here and there a longer and stronger central spine with the tip slightly bent downward. All the spines are awl-shaped and stiff.

"The radial spines when young are white with brownish (burnt) tips, later amber-colored above and below, grayish in the middle. The plant described is 3" high and a little more than an inch in diameter; radials 3, centrals 5 or 6 lines long. The plant in my possession did indeed bloom last summer but I was hindered unfortunately in describing the flowers in detail. They are smaller than those of *M. raphidacantha*, very similar in form, but the petals are yellow with saffron-yellow central stripes on the outer side."

Although Jacobi states definitely that the tubercles are not grooved on the upper surface, yet the presence of glands would indicate that the plant is not a *Mammillaria* but, more likely, a *Coryphantha* of the Series *Recurvatae* and perhaps one of the species already described. We have never seen glands in the axils of tubercles, except in genera having grooved tubercles. In cultivated specimens growing under abnormal conditions tubercles are sometimes produced without a groove and with glands in their axils.

MAMMILLARIA RAMOSISSIMA Quehl, Monatsschr. Kakteenk. 18: 127. 1908.

Globose to short-cylindric, dull grayish green; radial spines about 12, about 1 cm. long; central spines usually 1, sometimes 2 or 3; flowers and fruit unknown.

Type locality: Not cited.

Illustration: Monatsschr. Kakteenk. 18: 127.

MAMMILLARIA RECURVISPINA De Vriese, Tijdschr. Nat. Geschr. 6: 53. 1839.

Cactus recurvispinus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Solitary, somewhat depressed, about 16 cm. in diameter, glaucous; tubercles few, large, somewhat compressed, obtuse; areoles and axils of tubercles described as naked; spines all radial, 8, subulate, more or less incurved; flowers and fruit unknown.

Type locality: Mexico.

This plant was referred by Labouret to *Mammillaria sulcolanata* but was discussed by Schumann under *M. scheeri*; judging from the illustration, it is not close to either of these species but it is much nearer *Coryphantha bumamma*.

Illustration: Tijdschr. Nat. Geschr. 6: pl. 1. f. 1.

Figure 48 is a reproduction of the illustration cited above.

Mammillaria speciosa De Vriese (Tijdschr. Nat. Geschr. 6: 52. 1839. Not Don, 1830) is listed by Schumann among the species not known to him. It probably belongs to some species of *Coryphantha*.

The following names are without descriptions and can not be referred to any known species: *Coryphantha conspicua* Lemaire, Cactées 34. 1868; *Coryphantha engelmannii* Lemaire, Cactées 34. 1868; *Coryphantha hookeri* Lemaire, Cactées 34. 1868; *Coryphantha sublanata* Lemaire, Cactées 35. 1868.

7. NEOBESSEYA gen. nov.

Simple or tufted cacti, globose or somewhat depressed; tubercles irregular or somewhat spiraled, most of them grooved on upper side; flowers borne near top of plant, large, yellow or pink, probably always day-blooming; fruit globose, bright red, indehiscent; seeds black, globose, pitted, with a prominent white aril.

Type species: *Mammillaria missouriensis* Sweet.

Four species are recognized, all from the Great Plains of the United States.

The generic name commemorates Dr. Charles Edwin Bessey (1845-1915), professor in the University of Nebraska and for many years one of our eminent botanical teachers.

The genus is nearest *Coryphantha*, but it has very different fruit and seeds.

KEY TO SPECIES.

Flowers yellow.

Outer perianth-segments naked 1. *N. wismannii*

Outer perianth-segments ciliate.

Inner perianth-segments long-acuminate 2. *N. similis*

Inner perianth-segments at most acute 3. *N. missouriensis*

Flowers grayish pink 4. *N. notesteinii*

1. *Neobesseyia wissmannii* (Hildmann).

Mammillaria similis robustior Engelm., Bost. Journ. Nat. Hist. 6: 200. 1850.

Mammillaria nuttallii robustior Engelm., Proc. Amer. Acad. 3: 265. 1856.

Mammillaria missouriensis robustior S. Watson, Bibl.

Index 1: 403. 1878.

Cactus missouriensis robustior Coulter, Contr. U. S. Nat. Herb. 3: 111. 1894.

Mammillaria wissmannii Hildmann in Schumann, Gesamtb. Kakteen 498. 1898.

Cactus robustior Small, Fl. Southeast. U. S. 812. 1903.

Plant solitary, or forming mounds 2 to 3 dm. in diameter and 1 dm. high with 25 heads or more; areoles elliptic when young, conspicuously white-woolly, the head usually globose, tubercles rather large, spreading, somewhat narrowed towards apex; Spines 7 to 14, when young white to brownish, in age gray with yellow swollen base, acicular, 1.5 to 2 cm. long, sometimes all radial and spreading, rarely 1 or 2 centrals and these porrect; flowers large, 4 to 5 cm. long, dark yellow; scales on flower-tube strongly nerved; margin of perianth-segments naked; inner segments abruptly long-apiculate; fruit globose, 8 mm. in diameter.

Type locality: Not cited, presumably Texas.

Distribution: Central Texas.

Illustration: Blühende Kakteen 1: pl. 5, as *Mammillaria wissmannii*.

Figure 50 is a reproduction of the illustration cited above.




FIG. 50.—*Neobesseyia wissmannii*.

2. *Neobesseyia similis* (Engelmann).

Mammillaria similis Engelmann, Bost. Journ. Nat. Hist. 5: 246. 1845.

Mammillaria similis caespitosa Engelmann, Bost. Journ. Nat. Hist. 6: 200. 1850.

Echinocactus similis Poselger, Allg. Gartenz. 21: 107. 1853.

Mammillaria nuttallii caespitosa Engelmann, Proc. Amer. Acad. 3: 265. 1856.

Mammillaria missouriensis caespitosa S. Watson, Bibl. Index 1: 403. 1878.

Cactus missouriensis similis Coulter, Contr. U. S. Nat. Herb. 3: 111. 1894.

Mammillaria missouriensis similis Schumann, Gesamtb. Kakteen 498. 1898.

Cactus similis Small, Fl. Southeast. U. S. 812. 1903.

Coryphantha similis Britton and Rose in Britton and Brown, Illustr. Fl. ed. 2. 2: 571. 1913.

Plants sometimes growing in large clumps, to 1.5 dm. high by 2 to 3 dm. in diameter, containing 25 individuals or more; larger plants 6 to 10 cm. in diameter; tubercles deep green, cylindric, sometimes 2 cm. long, when young the groove filled with white wool; spines all puberulent; radial spines 12 to , spreading, dirty white with brownish tips; central spine solitary or often wanting, similar to but stouter and longer than the radials; flowers , to 6 cm. long, light yellow, the outer lobes tinged with brown and green; inner perianth-segments long, narrow, acuminate; flower-tube definite, covered nearly to its base with short greenish stamens; style green; stigma-lobes 4 to 6, linear; fruit globular or short-oblong, 10 to 20 mm. in diameter; seeds large, globose, 2 mm. in diameter.

Type locality: Near Industry, Texas.

Distribution: Eastern Texas.

Engelmann says that the flowers and fruits are larger than in *Mammillaria nuttallii*.

The inner perianth-segments gradually taper to the apex.

S. Watson and others refer here *Mammillaria caespitosa* Gray (Struct. Bot. 421. f. 838), but the plant illustrated by Gray is *Echinocereus reichenbachii*. The Index Kewensis refers *Mammillaria caespitosa* Gray, as they also do *Mammillaria similis*, to *Mammillaria missouriensis*. (See Cactaceae 3: 26).

Illustration: Cact. Mex. Bound. pl. 74, f. 7, as *Mammillaria nuttallii caespitosa* (seed).

Plate in, figure 2, shows a plant collected by F. E. Upham at Fort Worth, Texas, which flowered in Washington.

3. Neobesseyia missouriensis (Sweet).

Cactus mammillaris Nuttall, Gen. Pl. 1: 295. 1818. Not Linnaeus, 1753.

Mammillaria missouriensis Sweet, Hort. Brit. 171. 1826.

Mammillaria simplex Torrey and Gray, Fl. N. Amer. 1: 553. 1840.

Mammillaria nuttallii Engelm., Pl. Fendl. 49. 1849.

Mammillaria nuttallii borealis Engelm., Proc. Amer. Acad. 3: 264. 1856.

Cactus missouriensis Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Mammillaria missouriensis nuttallii Schelle, Handb. Kakteenk. 241. 1907.

Coryphantha missouriensis Britton and Rose in Britton and Brown, Illustr. Fl. ed. 2. 2: 570. 1913.

Plants solitary or cespitose, globose, 2.5 to 5 cm. in diameter; tubercles more or less spiraled, 10 to 15 mm. long; spines 10 to 20, acicular, gray, pubescent, all radial or sometimes 1 central; flowers greenish yellow; outer perianth-segments narrowly oblong, gradually tapering to an acute apex, ciliate; inner segments linear-lanceolate, attenuate; fruit globose, scarlet, about 1 cm. in diameter; seeds mm. in diameter.

Type locality: On the high hills of the Missouri, probably to the mountains.

Distribution: North Dakota to Montana, Colorado to Kansas, Oklahoma, and perhaps northern Texas.

This little cactus has a wide distribution on the Great Plains; both its conspicuous yellow flowers and its round red fruits are very attractive.

Coryphantha nuttallii, credited to Engelm., is cited as a synonym of *Mammillaria nuttallii* by Rümpler (Förster, Handb. Cact. ed. 2. 407. 1885).

Illustrations: Meehan's Monthly 10: pl. 3; Gartenwelt 1: 85, as *Mammillaria missouriensis*; Gartenwelt 1: 89, as *M. missouriensis viridescens*; Britton and Brown, Illustr. Fl. 2: f. 2525, as *Cactus missouriensis*; Schelle, Handb. Kakteenk. 241. f. 160, as *M. missouriensis nuttallii*; Cact. Mex. Bound. pl. 74, f. 6, as *M. nuttallii borealis*; Blanc, Cacti 72. No. 1426; Blühende Kakteen 3: pl. 145, as *M. nuttallii*; Britton and Brown, Illustr. Fl. ed. 2. 2: f. 2984, as *Coryphantha missouriensis*.

Plate XI, figure 4, shows a plant from a large clump sent by Professor C. O. Chambers in 1921 from Stillwater, Oklahoma.

4. Neobesseyia notesteinii (Britton).

Mammillaria notesteinii Britton, Bull. Torr. Club, 8: 367. 1891.

Cactus notesteinii Rydberg, Mem. N. V. Bot. Gard. 1: 272. 1900.

Oval, solitary or cespitose, about 3 cm. in diameter; tubercles nearly terete, about 6 mm. high; spines 12 to 18, white, turning gray, weak, slender, 8 to 12 mm. long, pubescent throughout, a central one usually present and frequently pink-tipped; flowers 15 to 25 mm. broad, ash-gray, tinged and penciled with pink, the segments broadly linear-oblong, mucronate; fruit obovoid; seeds black, globose, pitted.

Type locality: Near Deer Lodge, Montana.

Distribution: Known only from the type locality.

Professor F. N. Notestein, who first collected and observed this little cactus, found it in gravelly soil near a small creek; it differs from the other species of the genus in the color of the flowers and the more pubescent spines.

8. ESCOBARIA gen. nov.

Globose or cylindric, usually cespitose cacti, never milky; tubercles grooved above, persisting as knobs at the base of old plants after the spines have fallen; spines both central and radial, never hooked; flowers small, regular, appearing from top of plant at bottom of groove of young tubercles; stamens and style included; fruit red, naked (or with one scale), indehiscent, globular to oblong, crowned by the withering perianth; seeds brown to black; aril basal or subventral, oval.

Type species: *Mammillaria tuberculosa* Engelm.

The two species of this genus known to Schumann were placed by him in the subgenus *Coryphantha* of *Mammillaria*; they are like the *Coryphanthae* in having grooved flower-bearing tubercles, but are otherwise different, especially in the flowers, fruit, and seeds.

Eight species are known from northern Mexico and southern Texas.

The genus commemorates the work of two distinguished Mexicans, the Escobar brothers, Rómulo and Numa, of Mexico City and Juárez.

KEY TO SPECIES.

- Outer perianth-segments ciliate.
 Groove of tubercles without glands.
 Flowers large for the genus, 2 to 2.5 cm. long.
 Plants elongated; seeds very small, brown, with ventral hilum 1. *E. tuberculosa*
 Plants usually globose; seeds larger than in *E. tuberculosa*, black, with a sub-basal hilum 2. *E. dasyacantha*
 Flowers small, about 1.5 cm. long.
 Plants globose to stout-cylindric.
 Inner perianth-segments pointed.
 Inner perianth-segments broad 3. *E. chihuahuensis*
 Inner perianth-segments narrow 4. *E. runyonii*
 Inner perianth-segments obtuse 5. *E. chaffeyi*
 Plants slender-cylindric 6. *E. sneedii*
 Groove of tubercles with glands 7. *E. bella*
 Outer perianth-segments eciliate 8. *E. lloydii*

1. *Escobaria tuberculosa* (Engelmann).

Mammillaria strobiliformis Scheer in Salm-Dyck, Cact. Hort. Dyck. 1849. 504. 1850. Not Engelmann, 1848.

Echinocactus strobiliformis Poselger, Allg. Gartenz. 21: 107. 1853.

Mammillaria tuberculosa Engelmann, Proc. Amer. Acad. 3: 268. 1856.

Cactus tuberculosus Kuntze, Rev. Gen. Pl. 1: 261. 1891.



Cactus strobiliformis Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria strobiliformis pubescens Quehl, Monatsschr. Kakteenk. 17: 87. 1907.

Mammillaria strobiliformis durispina Quehl, Monatsschr. Kakteenk. 17: 87. 1907.

Mammillaria strobiliformis rufispina Quehl, Monatsschr. Kakteenk. 17: 87. 1907.

Mammillaria strobiliformis caespititia Quehl, Monatsschr. Kakteenk. 19: 173. 1909.

FIG. 51.—*Escobaria tuberculosa*.FIG. 52.—*Escobaria dasyacantha*.

Usually growing in clumps, cylindric or becoming so, 5 to 18 cm. high, 2 to 6 cm. in diameter; tubercles more or less regularly arranged in spirals, 6 mm. long; radial spines numerous, white, sometimes as many as 30, acicular, 4 to 15 mm. long; central spines several, stouter than radials, brown to blackish or colored only at tips, one of them usually porrect; flowers 2.5 cm. in diameter when fully expanded, light pink; outer perianth-segments acute, ciliate; inner perianth-segments narrowly pointed; fruit oblong, up to 20 mm. long, red; seeds pitted, with a small ventral hilum.

Type locality: Mountains near El Paso and eastward.

Distribution: Southwestern Texas, southern New Mexico, and adjacent Mexico.

Flowers appear in the afternoon and last for two days at least.

The name *Coryphantha tuberculosa* occurs in C. R. Orcutt's Circular to Cactus Fanciers, 1922.

Illustrations: Cact. Mex. Bound. pl. 12, f. 1 to 16, as *Mammillaria tuberculosa*; Förster, Handb. Cact. ed. 2. 417. f. 46; Schelle, Handb. Kakteenk. 235. f. 149, as *M. strobiliformis*.

Figure 51 is from a photograph of the plant sent by Dr. Shreve from near El Paso, Texas, in 1920.

2. *Escobaria dasyacantha* (Engelmann).

Mammillaria dasyacantha Engelmann, Proc. Amer. Acad. 3: 268. 1856.

Cactus dasyacanthus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Globose to short-oblong, usually 4 to 7 cm. in diameter but sometimes 20 cm. long; radial spines 20 or more, white, bristle-like; central spines about 9, stouter and longer than the radials, upper half usually reddish or brownish, often 2 cm. long; flowers pinkish; perianth-segments narrowly oblong, ciliate, apiculate; stigma-lobes green; fruit clavate, scarlet, 15 to 20 mm. long; seeds black, 1 mm. in diameter, slightly flattened, pitted, with a narrow white subbasal hilum.

Type locality: El Paso and eastward.

Distribution: Western Texas, southern New Mexico, and northern Chihuahua.

We have examined the type of this species which was collected by Charles Wright at El Paso in 1852.

Escobaria dasyacantha is sometimes mistaken for *Escobaria tuberculosa*, but the stems are usually globose and the seeds larger and of a different shape. Engelmann speaks of its resemblance to *Echinocactus intertextus dasyacanthus*, now *Echinomastus dasyacanthus*, but this is only superficial, for the flowers, fruit, and seeds of the two species are very different. The name *Coryphantha dasyacantha* occurs in C. R. Orcutt's Circular to Cactus Fanciers, 1922. We had never seen this plant in cultivation until it was recently sent by Mrs. S. L. Pattison from western Texas.

Illustrations: Cact. Mex. Bound. pl. 12, f. 17 to 22, as *Mammillaria dasyacantha*.

Plate VII, figure 1, shows a plant sent by Mrs. S. L. Pattison from near El Paso, Texas, in 1921 which flowered in the New York Botanical Garden. Figure 52 is from a photograph of another plant sent by Mrs. Pattison from the same region.

3. *Escobaria chihuahuensis* sp. nov.

Plants often solitary, perhaps also cespitose, globose to short-cylindric, very spiny; tubercles short, usually hidden by the spines; radial spines numerous, spreading; central spines several, longer than radials, usually brown or black in upper part; flowers small, 1 to 1.5 cm. long, purple; outer perianth-segments broad, often rounded at apex with ciliate margins; inner perianth-segments pointed.

Common in the mountains near Chihuahua, where it was collected by Palmer (No. 72, type) in 1908 and by Pringle (Nos. 250, 251) in 1885.

This plant should be compared with *Mammillaria grusonii* Runge (Gartenflora 38: 105. f. 20. 1889). L. Quehl believed that *M. grusonii* was closely related to *M. scheeri*, but he apparently knew it only from the original illustration and description. It does not suggest any of the species of *Coryphantha* to us.

4. *Escobaria runyonii* sp. nov.

Cespitose, with numerous (sometimes 100) globose to short-oblong heads, grayish green, 3 to cm. long with fibrous roots; tubercles 5 mm. long, terete in section with very narrow groove above; groove at first white-woolly, not glandular; radial spines numerous, acicular, white, 4 to mm. long; central spines stouter than radials, to 7, slightly spreading with brown or black tips, 6 to 8 mm. long; flowers 1.5 cm. long, pale purple; segments with a dark purple stripe down the middle and pale margins; outer perianth-segments narrow-oblong, with thin ciliate margins; inner perianth-segments narrower than the outer, with margins entire, acute; filaments purplish; style very pale; stigma-lobes 6, green; fruit scarlet, globose to short-oblong, 6 to 9 mm. long, juicy.

Collected by Robert Runyon in July 1921 and again in October of the same year near Reynosa, Mexico, about 75 miles up the Rio Grande from Brownsville, Texas, and on

August 10, 1921, near Rio Grande, Starr County, Texas. The plant flowered in Washington March 13, 1922.

Plate vi, figure 1, is from a photograph of the type plant taken by Robert Runyon. Figure 53 is from a photograph taken by Robert Runyon.

5. *Escobaria chaffeyi* sp. nov.

Short-cylindric, 6 to 12 cm. long by 5 to 6 cm. in diameter, almost covered by the numerous white spines; tubercles rather short, light green, with a narrow groove above; radial spines numerous, spreading, bristly; central spines several, a little shorter than the radials and brown or black-tipped; flowers 15 mm. long, cream-colored or sometimes purplish; outer perianth-segments ciliate; inner perianth-segments oblong, obtuse, entire; style white; stigma-lobes very short, yellowish green; fruit crimson, 2 cm. long.

Collected by Dr. Elswood Chaffey near Cedros, Zacatecas, Mexico, in June 1910 (No. 5, type), and by F. E. Lloyd near the same locality in 1908 (No. 29).




FIG. 53.—*Escobaria runyonii*.



FIG. 54.—*Escobaria sneedii*.

6. *Escobaria sneedii* sp. nov.

Densely cespitose, sometimes with as many as 50 joints, creeping or spreading; joints cylindric, up to 6 cm. long, 1 to 2 cm. in diameter; tubercles numerous, hidden under the many spines, terete, 2 to 3 mm. long, in age naked; groove narrow, hairy throughout its length; axils of tubercles not setose; spines 20 in a cluster or more, nearly white, or the larger ones brown at tip, longest one 6 mm. long, all usually appressed, but the longer ones near top, connivent; flowers small, 10 mm. long or less when dry, the outer segments long-ciliate; fruit (immature) a little longer than thick, 5 to 7 cm. long, green (?), at first juicy, naked; seeds globose, brown, nearly 1 mm. in diameter, pitted.


This curious little plant was sent us in February 1921 by Mrs. S. L. Pattison from southwestern Texas; it was collected by J. R. Sneed, who at first found only three clumps, but afterwards a fourth clump was discovered and again it was found in June 1921 just after it had flowered. It is known from a single station on the Franklin Mountains, Texas. According to Mrs. Slater the flowers are pink to saffron.

Figure 54 is from a photograph of a single plant sent by Mrs. Pattison in 1921.


7. *Escobaria bella* sp. nov.

Cespitose, cylindric, 6 to 8 cm. long; tubercles nearly terete, 1.5 to 2 cm. long, the groove white-hairy, with a narrow brownish gland near center; radial spines several, whitish, 1 cm. long or less; central spines 3 to 5, brown, unequal, the largest 2 cm. long or more, ascending; flowers

1



2



1. *Escobaria runyonii*, from Texas.
2. *Neomammillaria hemisphaerica*, from Texas.

central, small, rotate, nearly 2 cm. broad; perianth-segments pinkish with pale margins, linear-oblong, acute, the outer ones ciliate; filaments reddish; upper part of style and stigma-lobes green.

Collected by J. N. Rose and Wm. R. Fitch on hills of Devil's River, Texas (No. 17991).

Plate VII, figure 4, shows the type, which flowered in the New York Botanical Garden, March 31, 1914; figure a shows a tubercle with its gland-bearing groove.

8. *Escobaria lloydii* sp. nov.

Plant growing in clumps and resembling a small species of *Echinocereus*; old plants bearing naked corky tubercles; radial spines about 20, spreading, slender, white; central spines several, stout, with black or with brownish tips, 2 cm. long; flowers greenish with a central stripe on outside, 2.5 cm. long; filaments, style, and stigma-lobes green; fruit red, globose to short-oblong, 6 to 12 mm. long; seeds black, pitted, globose, 1 mm. in diameter.

Collected by F. E. Lloyd in foothills of Sierra Zuluaga, Zacatecas, Mexico, March 29, 1908 (No. 5).

This species is near *Escobaria tuberculosa*, but it has much stouter central spines and greenish white, eciliate inner perianth-segments.

SPECIES PERHAPS OF THIS RELATIONSHIP.

MAMMILLARIA EMSKOETTERIANA Quehl, Monatsschr. Kakteenk. 20: 139. 1910.

Cespitose, globose to short-cylindric, 5 cm. high; tubercles conic, their axils naked; radial spines 20 to 25; central spines 6 to 8, setaceous, white with black tips; flowers brownish yellow, 3 cm. long.

Type locality: Not cited.

We obtained a specimen of this plant from Quehl in 1913, but it has not done well nor has it flowered and we have not been able to refer it to any described species, but believe that it may be near *Escobaria tuberculosa*. Mr. Quehl believed that it was near *Mammillaria dasyacantha*, but if it came from San Luis Potosí, as Mr. Quehl supposed, it is doubtless specifically distinct from both. The following note is a translation of some remarks by Mr. Quehl:

"Our illustration shows a grafted specimen which has naturally grown more corpulent and consequently permits one to see better its general structure and the arrangement of the spines. Ungrafted specimens are thicker, lower, and, without other characteristics, can not be distinguished from a red-spined *Mammillaria pusilla* var. *multiceps*. Only a closer inspection reveals the wartfurrows and consequently the *Coryphantha*. The similarity is so great that I suspect that the new species is already more disseminated though not correctly recognized and the plants are either set aside or ignored as a form of *Mammillaria pusilla*. The plants before me were raised by Mr. Robert Emskötter, fancy and commercial gardener, of Magdeburg, after whom I have named the species, from mixed seed which he received from San Luis Potosí, so that Mexico may be regarded as its home."

Illustration: Monatsschr. Kakteenk. 20: 139.

9. BARTSCHELLA gen. nov.

Usually cespitose, globose to short-oblong cactus; tubercles large, somewhat united with the adjacent ones as in certain species of *Echinocactanae*, terete, not grooved, juicy, not milky; spines both radial and central, the latter usually hooked; flowers borne near top of plant, large, light purple or lavender; fruit short, hidden among the tubercles, seemingly dry, circumscissile; seeds dull black, pitted, with a narrow cylindric base, slightly constricted above; hilum large, slightly depressed, triangular.

Type species: *Mammillaria schumannii* Hildmann.

While this genus is probably to be referred to the *Coryphanthanae*, it possesses some characters of certain species of *Echinocactanae*, but the origin of the flower is quite different from any of them. The flower is large, like that of some species of *Coryphantha*, but the tubercles are not grooved and the seeds are not brown and reticulated. It differs from the

typical species of the so-called *Mammillaria* in its large flowers and black seeds, while from all of these genera it differs in its circumscissile fruit.

This monotypic genus is named for Dr. Paul Bartsch, curator in the United States National Museum, distinguished in conchology, who has sent us cacti from many out of the way places.

1. *Bartschella schumannii* (Hildmann).

Mammillaria schumannii Hildmann, Monatsschr. Kakteenk. 1: 125. 1891.

Mammillaria venusta K. Brandegee, Zoe 5: 8. 1900.

More or less cespitose (as many as 40 stems have been reported in a single cluster), 6 cm. high or less; axils slightly woolly, without bristles; radial spines 9 to 15, stout, 6 to 12 mm, long, brownish above, glabrous; central spines usually 1, sometimes 2 or 3, one of these usually hooked; in seedlings 10 or 11 radial spines developing, these spreading, feather-like with long spreading hairs; in one-year-old plants the spines simply puberulent, all white with brown tips and one central much longer than the others and strongly hooked; flower 3 to 4 cm. in diameter, the segments about 10, lanceolate, acuminate; stamens numerous, erect, shorter than the style; style slender, erect, pale; stigma-lobes 6, linear, green; fruit short, dull in color; seeds usually found in a cup between the tubercles, less than 1 mm. long.




FIG. 55.—*Bartschella schumannii*.




FIG. 55.—*Pelecyphora aselliformis*.

Type locality: Not cited.*

Distribution: Southern Lower California.

This species has been rare in collections, but considerable material was collected by Dr. Rose at Cape San Lucas, Lower California, in March 1911 (No. 16375). Living specimens were sent us from Lower California by Ivan M. Johnston in 1921.

Dr. C. H. Thompson writes under date of September 15, 1911, as follows:

"Your No. 16375, *Mammillaria venusta*, puzzles me. We received three plants from the New York Botanical Garden. Two are considerably shriveled but are reviving. The third is more plump and shows the vegetation characters better. In these it would readily be taken for *Mammillaria*, yet there are some appearances of the mamillate *Echinocacti*. You will observe how commonly adjacent tubercles cohere as in that group of *Echinocactus*, quite distinct from any *Mammillaria* that I know. Yet the position of the flower excludes it from *Echinocactus*. With flower and fruit characters you have observed it strikes me as being distinct from either genus."

* *Mammillaria schumannii* was described from a cultivated plant, but *M. venusta* came from San José del Cabo, Lower California.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering plant of *Escobaria dasyacantha*.
2. Fruit of *Dolicothele sphaerica*.
- 2a. Seeds of same.
3. Flowering plant of *Neomammillaria arida*.
4. Flowering plant of *Escobaria bella*.

- 4a. Tubercle of the same.
5. Flowering plant of *Neomammillaria crocidata*.
6. Flowering plant of *Bartschella schumannii*.
7. Flowering plant of *Neomammillaria carnea*.

Mammillaria schumanniana (Monatsschr. Kakteenk. 12: 178. 1902) was evidently intended for *M. schumannii*.

Illustrations: Monatsschr. Kakteenk. 1: facing 89; Thomas, Zimmerkultur Kakteen 51, as *Mammillaria schumannii*.

Plate VII, figure 6, shows a plant collected by Dr. Rose at Cape San Lucas, Lower California, in March 1911 (No. 16375), while a member of the scientific staff of the U. S. Steamer Albatross. Figure 55 is from a photograph of another plant from the same collection.

10. PELECYPHORA Ehrenberg, Bot. Zeit. 1: 737. 1843.

Plants small, cespitose, cylindric or globose, tuberculate, watery tubercles not arranged on ribs, strongly flattened, crowned with an elliptic areole bearing a pectinate spine, never grooved; flowers borne near center, broad, campanulate, purplish, the segments in definite series; flower-tube very short, slender; stamens short; fruit small, naked; seeds black, smooth.

Only one species, native of Mexico, is here recognized, *Pelecyphora aselliformis* Ehrenberg, the type. A second species has generally been referred here but it differs so widely from the other that we have no hesitancy in segregating it generically (see genus No. 13, p. 64).

The generic name is from *πέλεκυς* hatchet, and *φόρος* bearing, referring to the shape of the tubercles.

The plant has usually been regarded as a near relative of *Mammillaria*, but it has little in common with that genus. The flowers are central, borne in a mass of wool or hairs; the tubercles are not grooved and the seeds are black and smooth. It has been difficult for us, with the material at hand, to make out definitely the origin and position of the flower, but it seems to originate on the central sunken disk. This disk at first bears only clusters of hairs in the center of which the flower is produced. In time the flower opens and the tubercle, with its peculiar spiny crown, is developed, leaving in its axil the tuft of hairs about the flower.

1. *Pelecyphora aselliformis* Ehrenberg, Bot. Zeit. 1: 737. 1843.

Pelecyphora aselliformis concolor Hooker in Curtis Bot. Mag. 99: pl. 6061. 1873

Pelecyphora aselliformis grandiflora Haage jr., Cact. Kultur ed. 2. 206. 1900.

Tufted, cylindric, 5 to 10 cm. high, 2.5 to 5 cm. in diameter, covered with tubercles arranged in spirals; tubercles strongly flattened laterally, somewhat stalked at base; areoles at top of tubercles very bug and narrow, crowned by an elongated, scale-like spine with numerous lateral ridges, usually free at tip, giving a peculiar pectinate appearance; flowers cm. broad or more, campanulate; perianth-segments in 4 rows, the outer ones sometimes white, oblong, acute; stamens borne at top of flower-tube, much shorter than perianth-segments; stigma-lobes 4, erect; seeds 1 mm. broad, kidney-shaped.

Type locality: Mexico.

Distribution: About San Luis Potosí, Mexico.

This plant does not do well in cultivation. It is known generally as the hatchet cactus, and is also called peote and peyote, also peyotillo and peotillo; it is said by the Mexicans to possess medicinal properties.

Mammillaria aselliformis is, according to Watson (Cact. Cult. 188. 1889), was described in 1843, but we have found no other reference to it, except that Dr. A. Weber gives it as a synonym, crediting it to Monville. The name *Anhalonium asselliforme* Weber and *Ariocarpus aselliformis* Weber (Dict. Hort. Bois 931. 1898), quoted by Schumann as synonyms, were not formally published. *Pelecyphora fimbriata* Hildmann (Monatsschr. Kakteenk. 3: 68. 1893), simply a name, may or may not belong here.

Illustrations: Haage, Cact. Kultur ed. 2. 206, as *Pelecyphora aselliformis grandiflora*; Amer. Gard. 11: 474; Curtis's Bot. Mag. 99 pl. 6061, as *Pelecyphora aselliformis concolor*; Rümpler,

Sukkulenten 208. f. 118; Gartenflora 34: 25; Watson, Cact. Cult. 189. f. 75; ed. 3. f. 52; Cycl. Amer. Hort. Bailey 1: 203. f. 303; Stand. Cycl. Hort. Bailey 2: f. 718; Illustr. Hort. 5: pl. 186; Förster, Handb. Cact. ed. 2. 237 f. 21; Cact. Journ. 1: 107, 149; Krook, Handb. Cact. 34; Ann. Rep. Smiths. Inst. 1908: pl. 14, f. 6; Palmer, Cult. Cact. 117; Schelle, Handb. Kakteenk. 275. f. 197; Monatsschr. Kakteenk. 29: 81; Weinberg, Cacti 23; Knippel, Kakteen pl. 28; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 3; Garten-Zeitung 4: 218. f. 50; Blanc, Cacti 78. No. 1710; West Amer. Sci. 11: 8; Balt. Cact. Journ. 1: 89; 2: 164; Floralia 42: 369; Remark, Kakteenfreund 22; Haage, Cact. Kultur ed. 2. 206.

Figure 56 is reproduced from a painting made by Miss E. I. Schutt in 1907, of a plant sent from San Luis Potosí in 1905 by Dr. E. Palmer.

11. PHELLOSPERMA gen. nov.

A globular to cylindric, usually cespitose cactus with a large, fleshy, branched root; tubercles not grooved above, not milky; flowers borne in axils of old tubercles, funnel-shaped; fruit globular to cylindric, red, depressed at apex; seeds large (for this group), dull black, not pitted but rugose, with a thick corky base nearly as large as the body.

Type species: *Mammillaria tetrancistra* Engelm.

This genus differs from all its relatives in its very peculiar seeds. The flower, in its shape and origin, suggests the following genus, but in its color and size resembles *Coryphantha radiosa*. A single species is known, native of the western United States.

The generic name is from *φέλλος* cork, and *σπέρμα* seed, referring to the corky base of the seed.

1. *Phellosperma tetrancistra* (Engelmann).

Mammillaria tetrancistra Engelmann, Amer. Journ. Sci. II. 14: 337. 1852.

Mammillaria phellosperma Engelmann, Proc. Amer. Acad. 3: 262. 1856.

Cactus phellospermus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus tetrancistrus Coulter, Contr. U. S. Nat. Herb. 3: 104. 1894.

Solitary or cespitose, cylindric; sometimes becoming very large and then 3 dm. long, usually very spiny; root elongated, carrot-shaped or sometimes branched; tubercles terete, often elongated, their axils naked; radial spines numerous, acicular, white or sometimes with a brown tip, not pungent; central spines 1 to 4, stouter and longer than the radials, often brown or black, one or all strongly hooked; flower 3.5 to 4 cm. long, purple; base of tube slender, greenish, naked; scales and outer perianth-segments ciliate; style and stigma-lobes cream-colored; fruit rather variable in size, sometimes 3.7 cm. long, becoming dry in age, with a depressed umbilicus; seeds black, dull, 2 mm. long.

Type locality: San Felipe, California.

Distribution: Western Arizona, southeastern California, southern Utah, and southern Nevada; probably northern Lower California.

Mr. C. R. Orcutt, under date of March 5, 1922, comments on the distribution of this plant as follows:

"It reaches its greatest development on sandy and gravelly slopes near the White Water River east of Banning, California. It no doubt enters Lower California, for I believe that I have found it within a mile of the boundary line. It is comparatively rare in Arizona."

We have seen no specimens from Utah, but suspect that the plants from that state which have been referred to *Mammillaria grahamii* probably belong here. The species should be looked for in northern Lower California and Sonora.

Illustrations: Cact. Mex. Bound. pl. 7; Engler and Prantl, Pflanzenfam. 3^{6a}: 162. f. 56, B; Cact. Journ. 1: pl. for February; Bol. Direccion Estudios Biol. 2: f. 3; Monatsschr. Kakteenk. 20: 167, as *Mammillaria phellosperma*.

Figure 58 is from a photograph of a plant sent from California in 1921 by E. C. Rost; figure 57 shows a seed taken from a plant sent by Loren G. Polhamus in 1921 from Bard, California.



FIG. 57.—Seed of *Phellosperma tetrancistra*.

12. DOLICHOTHELE (Schumann) gen. nov.

Plant-body globose, more or less cespitose, soft in texture, never milky; tubercles elongated, not grooved above; flowers borne in axils of old tubercles, very large, with a definite funnel-shaped tube; inner perianth-segments yellow, spatulate, tapering into a claw and borne on top of tube; stamens forming a spiral about style and borne on whole face of throat, but forming a definite ring at top of throat; style slender; stigma-lobes linear; ovary exserted, naked; fruit smooth, greenish, purplish, or red, globose, ellipsoid or short-oblong; seeds black or brownish.

Type species: *Mammillaria longimamma* De Candolle.

The generic name is from *δολιχός* long, and *θηλή* nipple, referring to the elongated tubercles.

The fruit is not often collected and is not well known. Dr. Rose obtained a single fruit of one of the species, the only one we had then seen, in a private collection in Rome in 1915; this is nearly globular, red, thin-walled, many-seeded; the seeds are brownish, pitted, slightly flattened, pointed at base, with a small sub-basal hilum. In October 1921, Robert Runyon sent us a number of fruits which were greenish white to purplish, with black seeds, these somewhat flattened and pitted.




FIG. 58.—*Phellosperma tetrancistra*.




FIG. 59.—*Dolichothele longimamma*.

Mammillaria camptotricha Dams (Gartenwelt 10: 14. 1905) is usually considered as a close relative of this group, but it differs widely from it in the flowers as well as in other ways, and we believe that it is not congeneric with it (see page 126).

Three species, natives of southern Texas and northern and central Mexico, are recognized.

KEY TO SPECIES.

- Spines glabrous, even when very young; species of Texas and northern Mexico 1. *D. sphaerica*
 Spines puberulent; species of central Mexico.
 Tubercles very long (sometimes cm. long), pale green, glaucous; radials 6 or more;
 central spines usually present 2. *D. longimamma*
 Tubercles much shorter, bright green; radial spines 4 or 5; central spines none 3. *D. uberiformis*

1. *Dolichothele sphaerica* (Dietrich).

Mammillaria sphaerica Dietrich in Poselger, Allg. Gartenz. 21: 94. 1853.

Cactus sphaericus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria longimamma sphaerica K. Brandegee, Cycl. Amer. Hort. Bailey 2: 975. 1900.

Low and depressed, often growing in large cespitose masses 2 dm. in diameter, with a large thickened root; tubercles soft and turgid, resembling those of the following species (*D. longimamma*) but shorter, 12 to 16 mm. long; areoles small, circular, at first short-lanate; spines 12 to 15, glabrous, generally pale yellow, a little darker at base at first, in age darker, often reddish, 7 to 9 mm. long, spreading or a little curved backward; central spine 1, straight; flowers appearing toward top of plant but not from axils of younger tubercles, with a rotate limb 6 to 7 cm. broad; inner perianth-segments widely spreading, oblanceolate, acute to apiculate, tapering at base into a slender claw; stigma-lobes 8, yellow, narrow; fruit greenish white to purplish, short-oblong, 10 to 15 mm. long, juicy, very fragrant; seeds black, flattened, with a straight ventral face, rounded on the back, pitted; hilum subventral.

Type locality: Near Corpus Christi, Texas.

Distribution: Southern Texas and northern Mexico, especially along the Rio Grande from Eagle Pass to the sea.

Mr. R. D. Camp and Mr. Robert Runyon have recently found this species in abundance about Brownsville. With the aid of their material and the excellent photograph made by Mr. Runyon we have been able to present a detailed description of this plant.



FIG. 60.—*Dolichothele sphaerica*.




FIG. 61.—*Dolichotliele longimamma*.

According to Mr. Runyon, the flowers are very large and handsome. The fruit does not ripen until about the middle of October, and in one plant a single fruit continued to grow until the 27th of March and had a pronounced pleasing odor. This is the first case which has come under our notice in which any of the *Coryphanthanae* develop any odor in the fruits.

Illustration: Haage and Schmidt, Haupt-Verz. 1912: 36, as *Mammillaria sphaerica*.

Plate 1, figure 2, is from a photograph sent us by Robert Runyon from Brownsville, Texas; plate VII, figure 2, shows a fruit and figure a shows a seed from a plant collected by Mr. Runyon at Brownsville in 1921. Figure 60 is from a photograph of a flowering plant made by Mr. Runyon at Brownsville in 1921.

2. *Dolichothele longimamma* (De Candolle).

Mammillaria longimamma De Candolle, Mém. Mus. Hist. Nat. Paris 17: 113. 1828.

Mammillaria longimamma hexacentra Berg, Allg. Gartenz. 8: 130. 1840.

Mammillaria longimamma gigantothele Berg in Förster, Handb. Cact. 183. 1846.

Mammillaria longimamma congesta Hortus in Förster, Handb. Cact. 183. 1846.

Mammillaria uberiformis hexacentra Salm-Dyck, Cact. Hort. Dyck. 1849. 6. 1850.

- Mammillaria melaleuca* Karwinsky in Salm-Dyck, Cact. Hort. Dyck. 1849. 108. 1850.
Mammillaria globosa Link, Allg. Gartenz. 25: 240. 1857.
Mammillaria uberiformis gracilior Meinshausen, Wochenschr. Gärt. Pflanz. 1: 26. 1858.
Mammillaria longimamma luteola Hortus in Förster, Handb. Cact. ed. 2. 246. 1885.
Cactus longimamma Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus melaleucus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Mammillaria longimamma globosa Schumann, Gesamtb. Kakteen 508. 1898.

Solitary or cespitose, about 10 cm. high; tubercles elongated, 5 cm. long, somewhat glaucous, their axils hairy or naked; spine-areoles with white hairs when young, in age naked; radial spines 6 to 12, widely spreading, acicular, 2.5 mm. long, white to pale yellow, swollen and darker at base, puberulent; central spines 1 to 3, usually solitary, porrect, similar to the radials but usually darker with a blackish tip; flowers citron-yellow, 4 to 6 cm. long.

Type locality: Mexico.

Distribution: Central Mexico.

F. Haage jr. in his Choice Cacti lists ten varieties under this species; those not accounted for elsewhere are *ludwigii* and *melaleuca*.

Grässner in his Kakteen 1912 and also 1914 listed *Mammillaria longimamma* var. *ludwigii*. This may be a printer's error.

Mammillaria longimamma melaleuca is in the trade (Grässner). *Mammillaria longimamma pseudo-melaleuca* is advertised by Haage and Schmidt in their 1922 Catalogue.

Mammillaria longimamma spinosior (Wochenschr. Gärt. Pflanz. 1: 26. 1858), credited Link's Catalogue, but without description, is of this relationship.

Mammillaria hexacentra Otto and *Mammillaria gigantothele* (Förster, Handb. Cact. 183. 1846) were never described.

Krook (Handb. Cact. 41. 1855) mentions the variety *congesta* Hortus but gives no description. Several varieties of *Mammillaria longimamma* are in gardens; the following are mentioned by Schelle: *cristata*, *compacta* (the name cited by Rümpler in 1885), *major*, *laeta*, and *malaena*.

Mammillaria centricirrha flaviflora is referred by Schumann as a synonym of *M. melaleuca* which we have listed among the synonyms of *Dolichothele longimamma*. *M. alpina* Martius, mentioned elsewhere, may be of this relationship.

Illustrations: Monatsschr. Kakteenk. 29: 81, as *Mammillaria longimamma globosa*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 23, as *M. longimamma gigantothele*; Blühende Kakteen 2: pl. 73; De Candolle, Mém. Cact. pl. 5; Schumann, Gesamtb. Kakteen 792. f. 114; Monatsschr. Kakteenk. 8: 149; Schelle, Handb. Kakteenk. 244. f. 162; Förster, Handb. Cact. ed. 2. f. 22, a and b; Ann. Rep. Smiths. Inst. 1908: pl. 14, f. 2; Watson, Cact. Cult. 164. f. 63; ed. 3. f. 40; Dict. Gard. Nicholson 4: 564. f. 35; Suppl. 516. f. 551; De Laet, Cat. Gén. f. 89, as *Mammillaria longimamma*.

Figure 61 is from a photograph obtained from L. Quehl; figure 59 is from a photograph of the plant collected by Dr. E. Palmer near Victoria, Mexico, in 1907.

3. *Dolichothele uberiformis* (Zuccarini).

- Mammillaria uberiformis* Zuccarini in Pfeiffer, Enum. Cact. 23. 1837.
Mammillaria uberiformis major Hortus in Förster, Handb. Cact. ed. 2. 244. 1885.
Mammillaria uberiformis variegata Hortus in Förster, Handb. Cact. ed. 2. 244. 1885.
Mammillaria laeta Rümpler in Förster, Handb. Cact. ed. 2. 247. 1885.
Cactus uberiformis Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Mammillaria longimamma uberiformis Schumann, Gesamtb. Kakteen 508. 1898.

Globose, about 7.5 cm. high and 10 cm. in diameter; tubercles elongated, 2.5 to 3 cm. long, 12 to 15 mm. in diameter, bright green, shining, their axils naked; spine-areoles nearly naked; spines 4 or 5, all radial, puberulent, horn-colored to reddish, nearly equal; flowers yellow, 3 cm. broad; outer perianth-segments reddish; inner perianth-segments in 2 series, oblong, acute, acuminate; filaments white; style yellow; stigma-lobes 5 or 6, reflexed.

Type locality: Near Pachuca, Mexico.

Distribution: Central Mexico.

Illustrations: Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 13; Abh. Bayer. Akad. Wiss. München 2: pl. 1, VII. f. 6; Rümpler, Sukkulente 196. f. 109, as *Mammillaria uberiformis*.

Figure 62 is reproduced from the first illustration cited above.

13. SOLISIA gen. nov.

Plants very small, solitary, globular, tuberculate, milky; tubercles not arranged in ribs, small covered by broad pectinate spines; areoles very narrow and long; flowers lateral, yellow, small borne in axils of old tubercles; axils of tubercles neither hairy nor woolly; fruit naked, small, oblong; seeds black, smooth, dome-shaped with a broad basal hilum.

The type species, *Pelecypora pectinata* B. Stein, is here segregated from *Pelecypora*, with which it has little in common; it differs in being solitary, not cespitose, and in having the juice milky, not watery; the flowers small, lateral and yellow, not large, central and purple; the axils of the tubercles naked, not woolly; and the hilum of the seed broad and large, not small.




FIG. 62.—*Dolichothele uberiformis*.




FIG. 63.—*Solisia pectinata*.

The genus is named in honor of Octavio Solís of the City of Mexico, an earnest student of the cacti. Only one species is known.

1. *Solisia pectinata* (B. Stein).

Pelecypora pectinata B. Stein, Gartenflora 34: 25. 1885.

Pelecypora aselliformis pectinifera Rümpler in Förster, Handb. Cact. ed. 2. 238. 1885.

Pelecypora aselliformis pectinata Nicholson,* Dict. Gard. 4: 585. 1888.

Pelecypora aselliformis cristata Watson, Cact. Cult. 190. 1899.

Mammillaria pectinifera Weber, Dict. Hort. Bois 804. 1898.

Plants 1 to 3 cm. in diameter, fibrous-rooted, entirely hidden by the large overlapping spine-clusters; areoles narrow and long; spines 20 to 40, all radial, 5.5 to 2 mm. long, white, appressed; flowers small; fruit 6 mm. long; seed 1 mm. long.

Type locality: Mexico.

Distribution: Tehuacán, Mexico.

The cristate form of this species, when grown as a graft on some of the *Cereus* allies, becomes much larger than the normal form.

* Haage (Cact. Kultur ed. 2. 206. 1900) credits the variety to Ehrenberg.

This plant is very rare in living collections and is known only from a few localities near Tehuacán; one of these is near El Riego Hotel, where Dr. Rose obtained some 50 plants in 1905 but all have since died. We have been endeavoring since to obtain additional plants but Dr. Reko reports that this hill has been burned over and that no plants can now be found. Dr. Rose found it scattered over the top and side of a rounded hill, growing here and there among the stones and stunted plant life, looking not unlike the dull earth and pebbles.

Illustrations: Gartenflora **34**: 25; Garten-Zeitung **4**: 182. f. 42, No. 14; 217. f. 48; Grässner, Kakteen **1912**: 29; Ann. Rep. Smiths. Inst. **1908**: pl. 14, f. 5; Möllers Deutsche Gärt. Zeit. **25**: 477. f. 11, No. 4; **29**: 88. f. 10 (abnormal form), as *Pelecypora pectinata*; Monatsschr. Kakteenk. **29**: 81; Schelle, Handb. Kakteenk. 275. f. 198; Garten-Zeitung **4**: 217. f. 49, as *P. pectinata cristata*; Monatsschr. Kakteenk. **3**: 172. f. 5, as *P. aselliformis pectinata*.

Figure 63 is from an enlarged photograph showing the top of a plant, collected by Dr. Rose at Tehuacán in 1905.

14. NEOMAMMILLARIA nom. nov.

*Mammillaria** Haworth, Syn. Pl. Succ. 177. 1812. Not Stackhouse, 1809.

Plants globose, depressed-globose, or short-cylindric, occasionally much elongated, some with milky, others with watery juice; tubercles arranged in more or less spiraled rows, never on vertical ribs, terete, angled or sometimes flattened, never grooved on upper surface, usually bearing wool or hairs and sometimes bristles, but without glands in their axils and crowned by the spine-areoles; spines in clusters on top of tubercles, sometimes all alike, sometimes with central ones very different from the radial, all straight or sometimes one or more of central spines hooked; flowers, so far as known, diurnal, all from axils of old tubercles, much alike as to size and shape, more or less campanulate, comparatively small, variously colored, commonly red, yellowish or white to pinkish; perianth-segments rather narrow, spreading; stamens numerous, borne on base of perianth-tube, short, included; style about length of stamens; stigma-lobes linear; fruit usually clavate, rarely if ever globose, usually ripening rapidly, naked, scarlet (*Mammillaria brandegeei* with some scales and white fruit, according to Schumann) or white or greenish in a few species; seeds brown in some species, black in others.

The type is *Mammillaria simplex* Haworth, based on *Cactus mammillaris* Linnaeus.

We have given much time in attempting to group the species into definite series but have not succeeded, since many of the species are little known and incompletely described.

The name, *Neomammillaria*, as here used, replaces the name *Mammillaria* of Haworth (1812), which is a homonym of the *Mammillaria* of Stackhouse (1809), a genus of Algae.

The genus, as here treated, differs from Schumann's treatment (Gesamtb. Kakteen 472-601, 1898) in that we exclude three of his four subgenera, *Coryphantha*, *Dolichothele*, and *Cochemiea*, giving them generic rank. From his fourth subgenus we have excluded *Mammillaria micromeris* as the type of the genus *Epithelantha* † and *M. phellosperma* to the genus *Phellosperma* (see page 60).

The species, of which we recognize 150, are native chiefly of Mexico, extending northward into the southwestern United States; one species is reported as far north as Utah and Nevada. Two species are known from the West Indies (none is found in Jamaica or in the Lesser Antilles south of Antigua). Several species are known from Central America (none has been reported from Costa Rica, El Salvador, or Panama). One species is found in Venezuela and neighboring islands and one is described from Colombia, perhaps in error.

During the period of our investigation political conditions in Mexico have prevented our obtaining much original information concerning many of the species and have made it necessary for us to depend largely upon published descriptions and illustrations.

* The name was also spelled *Mammilaria* by Torrey and Gray (Flora 1: 553) and *Mamillar* a by Reichenbach (Mössl, Handb. ed. 2. 1: 1. 1827) and by Schumann (Gesamtb. Kakteen 472 and elsewhere).

† See Cactaceae, 3: 92. 1922.

KEY TO SPECIES.

- A. None of spines hooked (1-104, 150).
 B. Seeds brown (1-80).
 C. Tubercles giving off milk freely when pricked or cut (1-53).
 D. Axils of tubercles without bristles (1-33).
 E. Tubercles more or less elongated.
 F. Tubercles terete throughout.
 Spines yellow or reddish.
 Spines red1. *N. mammillaris*
 Spines yellow2. *N. nivosa*
 Spines usually white except tips, at least not red or yellow.
 Central spines 1 or 2.
 Central spines about length of radials3. *N. gaumeri*
 Central spines much longer than radials.4. *N. petrophila*
 Central spines 4 to 7.
 Outer perianth-segments entire; central spines long, slender .5. *N. arida*
 Outer perianth-segments erose; central spines not elongated, stouter than in preceding species...6. *N. brandegeei*
 FF. Tubercles more or less angled.
 G. Tubercles nearly terete towards apex.
 Outer perianth-segments and scales more or less fimbriate.
 Flowers reddish7. *N. gummifera*
 Flowers light yellow8. *N. macdougalii*
 Outer perianth-segments and scales entire.
 Radial spines white; flowers pinkish9. *N. beyderi*
 Radial spines brownish; flowers white to cream-colored.
 Plant hemispheric; radial spines 9 to 13; perianth-segments acute.....10. *N. hemisphaerica*
 Plant much flattened; radial spines 18 or fewer; perianth-segments acuminate11. *N. applanata*
 GG. Tubercles angled to top.
 H. Spines very unequal, some much elongated.
 Spines whitish12. *N. phymatothele*
 Spines horn-colored, reddish or black.
 No definite central spine.
 Spines horn-colored, short, curved.....13. *N. magnimamma*
 Spines reddish, long.....14. *N. macracantha*
 Central spines definite.
 Central spines 2.....15. *N. johnstonii*
 Central spines solitary.
 Central spine 2 to 3 cm. long16. *N. melanocentra*
 Central spine cm. long; oblong.....17. *N. runyonii*
 HH. Spines nearly equal, at least none much elongated.
 Flowers red to pinkish.
 Outer perianth-segments ciliate18. *N. sartorii*
 Outer perianth-segments not ciliate (so far as known).
 Central spines none.
 Spines pinkish with black tips19. *N. seitziana*
 Spines straw-colored throughout20. *N. ortegae*
 Central spines 1 or more.
 Central spines solitary; radial spines nearly equal .21. *N. meiacantha*
 Central spines; some of radials very short22. *N. scrippsiana*
 Flowers yellowish.
 Central spines 4 to 623. *N. gigantea*
 Central spines usually wanting24. *N. peninsularis*
 EE. Tubercles very short, symmetric.
 F. Plants globose or depressed.
 G. Axils of tubercles naked.....25. *N. flavovirens*
 GG. Axils of tubercles laniferous.
 Some of spines deciduous26. *N. sempervivi*
 None of spines deciduous.
 Central spines present.....27. *N. obscura*
 Central spines wanting.....28. *N. crocidata*
 FF. Stems cylindric or ovoid.
 Central spines wanting.
 Tubercles nearly terete.....29. *N. polythele*
 Tubercles 4-angled.
 Tubercles pointed; axils very woolly.....30. *N. carnea*
 Tubercles not pointed; axils not very woolly31. *N. lloydii*
 Central spines several.
 Radial spines reduced to short bristles32. *N. zuccariniana*
 Radial spines more elongated than in last species33. *N. formosa*

KEY TO SPECIES—continued.

- DD. Axils of tubercles with bristles as well as wool
 E. Some of spines much elongated, curved, and flexuous.
 Definite central spines wanting34. *N. compressa*
 Central spines present.
 Central spines weak35. *N. mystax*
 Central spines stiff36. *N. petterssonii*
 EE. None of spines elongated, or if elongated, not flexuous.
 F. Tubercles terete or nearly so.
 G. Axils of tubercles bearing yellow wool37. *N. eichlamii*
 GG. Axils of tubercles bearing white wool.
 H. Spines all radial.
 Spines or 6, in young plants sometimes only 438. *N. karwinskiana*
 Spines always 439. *N. praelii*
 HH. Spines both radial and central.
 I. Radial spines numerous, 52 or more.
 Central spines reddish, not much longer than the
 radials.
 Outer perianth-segments ciliate40. *N. standleyi*
 Outer perianth-segments setose41. *N. evermanniana*
 Central spines elongated, usually white except at tip.
 Flowers yellow42. *N. parkinsonii*
 Flowers dark red43. *N. geminispina*
 II. Radial spines few, to 9.
 Spines black when young44. *N. pyrrhocephala*
 Spines at most brownish.
 Flowers yellow45. *N. woburnensis*
 Flowers pinkish46. *N. collinsii*
 FF. Tubercles strongly angled.
 G. Spines both radial and central.
 Radial spines numerous47. *N. chinocephala*
 Radial spines few, bristle-like.
 Central spines 4 to 648. *N. tenampensis*
 Central spines 249. *N. polygona*
 GG. Spines few, all of one kind.
 Flowers yellow50. *N. confusa*
 Flowers rose-colored or white.
 Flowers rose-colored.
 Plants globose; stigma-lobes 4 or 551. *N. villifera*
 Plants cylindric; stigma-lobes 852. *N. polyedra*
 Flowers white53. *N. conzattii*
 CC. Milk-tubes developed, if at all, only in stem; tubercles not milky (54-80).
 D. Central spines wanting.
 Spines subulate; areoles elliptic54. *N. napina*
 Spines mostly acicular; areoles circular.
 Spines numerous55. *N. lanata*
 Spines few (4 to 6).
 Spines 5 or 6, short, straight56. *N. kewensis*
 Spines 4, elongated, curved.
 Flowers large (2.5 cm. broad)57. *N. subpolyedra*
 Flowers small.
 Spines long and weak58. *N. galeottii*
 Spines subulate59. *N. tetracantha*
 DD. Central spines present.
 E. Central spines usually 2, sometimes solitary.
 F. Radial spines 20 or more.
 Central spines stout and not very long; stigma-lobes white.
 Plant round or nearly so at apex; central spine often 160. *N. elegans*
 Plant strongly umbilicate; central spines always 261. *N. pseudoperbella*
 Central spines long62. *N. dealbata*
 FF. Radial spines 20 or fewer.
 Radial spines white, bristle-like.
 Stigma-lobes red.
 Globose or somewhat elongated63. *N. haageana*
 Depressed-globose64. *N. perbella*
 Stigma-lobes white.
 Radial spines appressed65. *N. collina*
 Radial spines not appressed66. *N. donatii*
 Radial spines brownish when young, stouter than in the last67. *N. mundtii*
 EE. Central spines usually 4, sometimes more.
 F. Central spines white or yellow.
 Radial spines white.
 Plant globose.
 Axils of tubercles not setose; central spines usually 468. *N. celsiana*
 Axils of tubercles setose; central spines usually 969. *N. aureiceps*
 Plant cylindric.
 Plants from Yucatan70. *N. yucatanensis*
 Plants from Central America71. *N. ruettii*

KEY TO SPECIES—continued.

- Radial spines yellow.
 Plants globular. 72. *N. pringlei*
 Plants slender-cylindric. 73. *N. cerralboa*
- FF. Central spines brown or black.
 Central spines black. 74. *N. phaeacantha*
 Central spines brown.
 Axils of tubercles not setose. 75. *N. graessneriana*
 Axils of tubercles setose.
 Tubercles closely set.
 Central spines not very different from radial.
 Plant body elongated; spines brownish or reddish. 76. *N. spinosissima*
 Plant body globose; radial spines whitish. 77. *N. densispina*
 Central spines very different from the radial. 78. *N. nunezii*
 Tubercles spreading.
 Central spines unequal; stigma-lobes green. 79. *N. amoena*
 Central spines nearly equal; stigma-lobes rose-colored. 80. *N. rhodantha*
- BB. Seeds black; neither tubercles nor stems milky (81-104).
 C. Spines plumose. 81. *N. plumosa*
- CC. Spines not plumose.
 D. Radial spines weak and hair-like.
 Central spines with yellow tips. 82. *N. prolifera*
 Central spines with brown tips. 83. *N. multiceps*
- DD. Radial spines not hair-like.
 E. Spines yellow.
 Spines 2 to 8, glabrous, more or less twisted or bent. 84. *N. camptotricha*
 Spines about 20, pubescent, straight. 85. *N. eriacantha*
- EE. Spines not yellow.
 F. Spines 25 to 80.
 Spines pubescent or lanate.
 Spines lanate, 25 to 30. 86. *N. schiedeana*
 Spines pubescent or puberulent. 87. *N. lasiacantha*
 Spines not pubescent.
 Spines all very much alike.
 Perianth-segments obtuse. 88. *N. denudata*
 Perianth-segments pointed.
 Flowers about 7 mm. long. 89. *N. lenta*
 Flowers about 2 cm. long. 90. *N. candida*
 Central spines 1 to 6, very unlike others. 91. *N. vetula*
- FF. Spines 20 or fewer but sometimes more in *N. oliviae* and *N. pottsii*.
 Plant globose.
 Flowers red. 92. *N. fertilis*
 Flowers white.
 Central spines solitary; radials 7 to 9. 93. *N. decipiens*
 Central spines to 8; radials 16 to 20. 94. *N. discolor*
- Plant cylindric.
 Joints very fragile, breaking loose when touched or jarred. 95. *N. fragilis*
 Joints not fragile.
 Spines all radial, recurved, sometimes with one central. 96. *N. elongata*
 Spines both radial and central.
 Plants globose to short-cylindric. 97. *N. oliviae*
 Plants slender-cylindric.
 Axils of tubercles not bristly.
 Spines all yellow. 98. *N. echinaria*
 Spines not yellow.
 Upper central spines more or less connivent
 over top of plant. 99. *N. pottsii*
 Upper central spines not connivent. 100. *N. mazatlanensis*
 Axils of tubercles bristly.
 Stems slender-cylindric; central Mexican species. 101. *N. sphacelata*
 Stems short-cylindric or globose (sometimes globose
 in *N. palmeri*); Lower Californian species.
 Spines nearly white or at least becoming so; seeds
 minute.
 Spines all white or nearly so; spine-areoles at
 first lanate. 102. *N. albicans*
 Spines tan with dark tips; spine-areoles not
 lanate. 103. *N. slevinii*
 Spines not white; seeds 3 mm. long. 104. *N. palmeri*
- AA. Some of central spines hooked; radial spines never hooked (105-149).
 B. Tubercles milky; seeds brown.
 Plants globose. 105. *N. uncinata*
 Plants cylindric. 106. *N. hamata*
- BB. Tubercles not milky except sometimes in *N. rekoii*; seeds mostly black.
 C. Seeds brown.
 Fruit red; flowers from side of plant. 107. *N. rekoii*
 Fruit green; flowers from near base of plant. 108. *N. solisii*



KEY TO SPECIES—continued.

- CC. Seeds black.
- D. Fruit elongated, clavate, ripening quickly.
- E. Seeds not rugose.
- F. Plants usually small; spines setaceous to delicately acicular.
- Central spines yellow.
- Central spines glabrous 109. *N. pygmaea*
- Central spines pubescent.
- Flowers white 110. *N. wildii*
- Flowers yellowish 111. *N. seideliana*
- Central spines red to brown.
- Outer perianth-segments ciliate.
- Central spines shorter than flower; perianth-segments acute 112. *N. barbata*
- Central spines longer than flower; perianth-segments obtuse 113. *N. mercadensis*
- Outer perianth-segments entire.
- Axils of tubercles setose.
- Inner perianth-segments white to yellowish.
- Central spines 3 or 4.
- Radial spines about 25; flowers 2 cm. long. 114. *N. kunzeana*
- Radial spines about 20; flowers 1 cm. long. 115. *N. hirsuta*
- Central spines 7 to 9 116. *N. multihamata*
- Inner perianth-segments red or reddish.
- Radial spines weak and hair-like.
- Central spines several 117. *N. longicoma*
- Central spine solitary 118. *N. bocasana*
- Radial spines stiff.
- Radial spines glabrous 119. *N. multiformis*
- Radial spines pubescent 120. *N. scheidweileri*
- Axils of tubercles not setose.
- Flowers 2.2 cm. long or more.
- Central spines solitary 121. *N. saffordii*
- Central spines 3 122. *N. schelbasi*
- Flowers 1 to 1.5 cm. long.
- Plants caespitose 123. *N. glochidiata*
- Plants solitary.
- Inner perianth-segments acuminate. 124. *N. trichacantha*
- Inner perianth-segments acute only 125. *N. painteri*
- FF. Plants stout; central spines at least stout-acicular to subulate.
- G. Outer perianth-segments ciliate.
- Fruit purple, ovoid to globular.
- Radial spines 15 or less; fruit large (25 mm. long) . . . 126. *N. wrightii*
- Radial spines 20 to 30; fruit 10 to 15 mm. long . . . 127. *N. viridiflora*
- Fruit scarlet, clavate.
- Outer perianth-segments long-ciliate 128. *N. wilcoxii*
- Other perianth-segments and upper scales short-ciliate.
- Perianth rotate; stigma-lobes red 129. *N. mainae*
- Perianth campanulate; stigma-lobes green.
- Flowers white 130. *N. boedekeriana*
- Flowers purple to pinkish.
- Radial spines often as many as 30.
- Inner perianth-segments acuminate 131. *N. microcarpa*
- Inner perianth-segments usually obtuse or rounded. 132. *N. milieri*
- Radial spines often as few as 3; inner perianth-segments acute 133. *N. sheldonii*
- GG. Perianth-segments not ciliate.
- H. Setae in axils of tubercles.
- I. Seeds constricted above base.
- J. Flowers greenish or pink, and small.
- K. Flowers greenish, 10 to 12 mm. long; central spines yellowish to reddish. 134. *N. armillata*
- KK. Flowers pink, 20 mm. long or more; central spines dark brown 135. *N. fraileana*
- JJ. Flowers nearly white, seeds much larger. 136. *N. swinglei*
- II. Seeds not constricted above base.
- Central spines several; flowers yellowish 137. *N. dioica*
- Central spines usually solitary; flowers rose-colored. . 138. *N. goodridgei*
- HH. Setae wanting in axils of tubercles.
- I. Flowers rotate 139. *N. zephyranthoides*
- II. Flowers campanulate.
- Plants globose.
- Flowers white 140. *N. carretii*
- Flowers pink to purplish.
- Inner perianth-segments obtuse 141. *N. jaliscana*
- Inner perianth-segments acute to acuminate . . . 142. *N. bombycina*
- Plants slender, elongated and cylindric.
- Flowers small, pinkish 143. *N. occidentalis*
- Flowers large, purplish. 144. *N. fasciculata*
- EE. Seeds rugose (perhaps a generic type) 145. *N. nelsoni*,
- DD. Fruit depressed, long-persisting (perhaps a generic type) 146. *N. longiflora*
- AAA. Species not grouped: 147, *N. tacubayensis*; 148, *N. umbrina*; 149, *N. verhaertiana*; 150, *N. xanthina*.

1. *Neomammillaria mammillaris* (Linnaeus).

- Cactus mammillaris* Linnaeus, Sp. Pl. 1: 466. 1753.
Cactus mammillaris glaber De Candolle, Pl. Succ. 137. 1799.
Mammillaria simplex Haworth, Syn. Pl. Succ. 177. 1812.
 ? *Mammillaria conica* * Haworth, Suppl. Pl. Succ. 71. 1819.
Mammillaria parvimamma Haworth, Suppl. Pl. Succ. 72. 1819.
Cactus microthele Sprengel, Syst. 2: 494. 1825.
Mammillaria simplex parvimamma Lemaire, Cact. Gen. Nov. Sp. 98. 1839.
Mammillaria caracasana † Otto in Salm-Dyck, Cact. Hort. Dyck. 1849. 107. 1850.
Mammillaria mammillaris Karsten, Deutsche Pl. 888. 1882.
 ? *Cactus conicus* Kuntze, Rev. Gen. Pl. 1: 259. 1891.
Cactus parvimammus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Globose to short-cylindric, 4 to 6 cm. high; tubercles short, 5 to 7 mm. long, conic, nearly terete, pale green, only slightly woolly in their axils; spine-areoles bearing a dense mass of white wool when young; spines reddish brown, acicular; radial spines 10 to 12, spreading, 5 to 7 mm. long; central spines 3 or 4, stouter and a little longer than the radials; flowers 8 to 10 mm. long, cream-colored; outer perianth-segments narrow, bearing long mucronate tips; fruit 15 to 20 mm. long, red; seeds minute, brown.

FIG. 64.—*Neomammillaria mammillaris*.FIG. 65.—*Neomammillaria macdougallii*.

Type locality: Tropical America.

Distribution: Northern Venezuela and neighboring Dutch Islands.

This plant was the first-known species of the genus and the only one known to Linnaeus; it was described and illustrated by Commelin in 1697 and by Hermann in 1698. It was one of the first cacti discovered; Aiton states that it was cultivated by Bishop Compton before 1688. The cited distribution of the species has usually been inexact or erroneous; Linnaeus gave no definite locality but restricted it to the warm parts of America.

Nuttall assigns it also to the hills of the Missouri River, and De Candolle's range covers that of both Linnaeus and of Nuttall. Nuttall's plant was subsequently found to be different from the one of the Caribbean region. Schumann gives the range as the West Indies but his description covers two or three species. A number of his references are erroneous, for neither Wright's plant (No. 2619, as *Mammillaria pusilla*) from Cuba nor Haworth's plant (Syn. Pl. Succ. 177, as *Mammillaria prolifera*) from the West Indies belongs

* Tubercles large, conic; spines less than 10, all radial, red, but paler at base; flowers and fruit unknown. Neither Pfeiffer nor Schumann knew this species or its origin. The Index Kewensis refers it to South America. If from that region it might be a species of *Discocactus*, near *D. placitifolius*, but it may belong here.

† This is the original spelling, but Schumann wrote it *M. caracasana*.

here. The name *Cactus prolifer* Willdenow (Pfeiffer, Enum. Cact. 9. 1837) is doubtless to be referred here. Fawcett lists the plant from Jamaica (as *Mammillaria simplex*) but no specimens are known to us from that island, which was searched by Dr. Britton and the late Mr. William Harris.

De Tussac (Fl. Antill. 2: 216, pl. 32) refers it to Santo Domingo and he describes and figures it, mentioning a locality in the desert near Gonaives which, however, is in Haiti; his illustration, while undoubtedly of this species, is not an original but copied from that of De Candolle (Pl. Succ. pl. 111). The only similar plant we know from his locality is *Mammillaria pusilla*, described as *M. pusilla haitiensis* by Schumann, which has been collected by Buch at this locality, and we have specimens from other collectors. We now believe that *Neomammillaria mammillaris* is confined to the coast of Venezuela and the adjacent islands, among which is Curaçao. In 1913 Dr. Britton and Dr. Shafer found it common on the top of a limestone hill in Curaçao (No. 3085) and in the same year Mr. Pittier obtained living plants near Cabo Blanco, Venezuela (No. 6471). These two are the only collections which have been made in recent years.

Steudel (1821), under *Mammillaria simplex*, compares this species with *Cereus flavescens* and *C. lanuginosus*, but he must have meant *Cactus* instead of *Cereus*.

Mammillaria microthele Monville and *M. micrantha* Hortus are names which Rümpler (Förster, Handb. Cact. ed. 2. 335. 1885) refers to *M. caracassana*; Salm-Dyck (Cact. Hort. Dyck. 1844. 9. 1845) also referred to it *M. micracantha* Monville.

Mammillaria simplex affinis Otto is mentioned by Förster (Handb. Cact. 217. 1846), but is not described.

Mammillaria karstenii Poselger (Allg. Gartenz. 21: 95. 1853) is listed by Schumann among his little-known species. The Index Kewensis states that it comes from Argentina, which is doubtless a mistake. The type locality is given as "La Canada," a common Spanish locality name. If collected by Karsten, it probably was obtained in Venezuela, in which case it would probably be referable to *Neomammillaria mammillaris*.

Mammillaria fuliginosa Salm-Dyck (Cact. Hort. Dyck. 1849. 93. 1850) we do not know, but if it came from Venezuela, where it is referred doubtfully by the Index Kewensis, it would belong here.

Illustrations: Hermann, Parad. 132. pl. 137, as *Echinomelocactus minor*, etc.; Commelin, Hort. Amst. 1: 105. f. 55; Plukenet, Opera Bot. 1: 148. pl. 29, f. 1, as *Ficoides*, etc.; Bradley, Hist. Pl. Succ. 3: 11. pl. 29, as melon-thistle; Loudon, Encycl. Pl. ed. 2 and 3. 410. f. 6839; De Candolle, Pl. Succ. 137. pl. 111; Fl. Antill. 2: pl. 32, as *Cactus mammillaris*; De Candolle, Mém. Cact. pl. 7, as *Mammillaria simplex*.

Figure 64 is reproduced from a colored drawing by Miss M. E. Eaton of a plant obtained by Dr. Britton and Dr. Shafer on Curaçao in 1913, which fruited the same year in the New York Botanical Garden.

2. *Neomammillaria nivosa* (Link).

Mammillaria nivosa Link in Pfeiffer, Enum. Cact. II. 1837.

Cactus nivosus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Coryphantha nivosa Britton, Ann. Mo. Bot. Gard. 2: 45. 1915.

Often forming large clusters 8 dm. in diameter, of 25 heads or more; separate specimens usually globose but sometimes cylindric, the largest ones 18 cm. in diameter, very spiny; tubercles milky, 10 mm. long, their axils filled with white wool; spines usually 14, bright yellow, acicular, the longer ones 1.5 cm. long; spine-areoles when young woolly, in age naked; flowers cream-colored, 1.5 cm. long; fruit clavate, 12 mm. long, red; seeds brown.

Type locality: Tortola Island, Virgin Islands.

Distribution: Southern Bahamas, Mona, Desecheo, Culebra, Buck Island, St. Thomas, Little St. James Island, Tortola, and Antigua.

Known as the snowy cactus in the Virgin Islands and as the woolly nipple-cactus in the Bahamas.

The plant inhabits crevices of rocks and locally is very abundant. On Mona Island, between Porto Rico and Santo Domingo in the Mona Passage, it exists in immense numbers on the limestone plateau.

Mammillaria tortolensis (Pfeiffer, Enum. Cact. 11. 1837) was published by Pfeiffer as a synonym of *M. nivosa*. The same or similar plant was briefly described by Forbes (Journ. Hort. Tour 148, 1837).

Illustrations: Förster, Handb. Cact. ed. 2. 331. f. 34; Schelle, Handb. Kakteenk. 264. f. 186; De Laet, Cat. Gén. f. 46; Blühende Kakteen 3: pl. 165, as *Mammillaria nivosa*.

Figure 66 is from a photograph of a plant collected on Turks Island, British West Indies, in July 1916 and sent us by the Director of the New York Aquarium; figure 243 (Britton and Rose, Cactaceae 3: p. 231) shows the plant on Mona Island, Porto Rico.




FIG. 66.—*Neomammillaria nivosa*.

3. *Neomammillaria gaumeri* sp. nov.

Cespitose, the branches short, globose to short-cylindric, up to 15 cm. long, growing half hidden in the sand; tubercles dark green, short, nearly terete, obtuse, 5 to 7 mm. long, very milky; axils naked even when young; spine-areoles conspicuously white-woolly at first, soon naked; radial spines 10 to 12, spreading, acicular, white with brown tips or lower ones in cluster darker, 5 to 7 mm. long; central spine solitary, porrect, usually brown; flowers very abundant from near top of plant but not from axils of young areoles, creamy white, small, 10 to 14 mm. long; outer perianth-segments greenish, brown-tipped; scales on flower-tube broadly ovate, scarious; fruit crimson, clavate, 18 to 20 mm. long, naked.

Common in the sand dunes of Progreso, Yucatan; collected first by George P. Gaumer and sons, April 1916 (No. 23349, type); re-collected in 1918 and again in 1921.

This species is remarkable for its unusual habitat and was the first of the genus reported from Yucatan. A second species has since been collected by Dr. Gaumer (see p. 114).



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Fruiting plant of *Neomammillaria gaumeri*.
2. Flowering plant of *Neomammillaria heyderi*.
- 2a. Fruit of same.
3. Flowering and fruiting plant of *Neomammillaria hemisphaerica*.
4. Flowering plant of *Neomammillaria compressa*.
5. Flowering plant of *Neomammillaria geminispina*.
6. Flowering plant of *Neomammillaria hemisphaerica*.

It is perhaps nearest some of the species from Texas, such as *N. hemisphaerica* and *N. heyderi*, but when growing it is easily distinguished by the peculiar white mats of wool on the young spine-areoles.

The following interesting note has been contributed by Dr. Gaumer, in whose honor the plant is named:

"The flowers begin to open at 8 a. m., are fully open at noon, close at dawn, and shrink the next morning, leaving the ovary wholly imbedded in the mass of the plant at the base of the tubercles; it remains dormant from 3 to 6 months, then suddenly develops to an inch in length in 48 hours. If put away in a dry place the bright crimson berries last from 3 to 6 months without decaying or changing their color. When thoroughly ripe they have a rather pleasant sweetish taste and are said to be edible.

"The plant multiplies by seed and by segmentation; this latter is accomplished by the plant putting out numerous shoots from its upper surface; these send out roots; the old plant decays and the little ones are often rolled about by the cattle or by the winds, and later send out stronger roots that finally anchor them to the sand, generally under a clump of brush."

Plate VIII, figure 1, shows the type plant which flowered in the New York Botanical Garden, July 24, 1918, soon after its arrival from Yucatan; plate XIII, figure 2, is from a photograph of the plant showing the large masses of white wool at the young spine-areoles.

4. *Neomammillaria petrophila* (Brandege).

Mammillaria petrophila Brandege, *Zoe* 5: 193. 1904.

Sometimes cespitose, milky, globular, 15 cm. in diameter or less; tubercles short, broad at base; spines at first chestnut-colored, becoming pale in age; radial spines 10, about 1 cm. long, a little spreading; central spine 1 (rarely 2), 2 cm. long, darker and stouter than the radials; flowers bright greenish yellow, 18 to 20 mm. long; perianth-segments hardly acute, sometimes slightly erose; stamens and style yellow; stigma-lobes 6; fruit small, roundish; seeds reddish brown, smooth, less than 1 mm. long.

Type locality: Sierra de la Laguna, Lower California.

Distribution: Mountains of southern Lower California.

We know this species only from description and illustration.

Illustration: Monatsschr. Kakteenk. 17: 57, as *Mammillaria petrophila*.

5. *Neomammillaria arida* (Rose).

Mammillaria arida Rose in Quehl, Monatsschr. Kakteenk. 23: 181. 1913.

Plants usually single, globular, deeply seated in the ground, 3 to 6 cm. in diameter, containing much milk and giving it off freely when injured; tubercles nearly terete; radial spines about 15, pale, ascending, the bases sometimes yellowish and the tips dark; central spines 4 to 7, 12 to 16 mm. long, much longer than the radials, dark brown, erect; flowers 1 cm. long; outer perianth-segments dark purple with lighter margins, entire; inner perianth-segments cream-colored to almost pale yellow; stamens pale; stigma-lobes green; fruit clavate, red, 15 cm. long; seeds brown.

Type locality: Hills near Pichilique Island near La Paz, Lower California.

Distribution: Known only from the type locality.

Plate VII, figure 3, shows one of the plants collected by Dr. Rose in 1911 which flowered in the New York Botanical Garden, July 2, 1912.

6. *Neomammillaria brandegeei* (Coulter).

Cactus brandegeei Coulter, Contr. U. S. Nat. Herb. 3: 96. 1894.

Cactus gabbii Coulter, Contr. U. S. Nat. Herb. 3: 109. 1894.

Mammillaria gabbii Engelm. in K. Brandege, *Erythea* 5: 116. 1897.

Mammillaria brandegeei K. Brandege, *Erythea* 5: 116. 1897.

Cylindric to globular, flattened, solitary or in clusters of 2 to 8; tubercles angled; axils woolly; radial spines 9 to 16, 8 to 10 mm. long, yellowish brown; central spines 3 to 6, a little longer and darker than the radials; flowers 15 mm. long; outer perianth-segments ovate, striate, ciliate; inner perianth-segments greenish yellow, narrower than the outer, entire; fruit white (according to Schumann), bearing a few narrow scales.

Type locality: San Jorge, Lower California.

Distribution: Lower California, San Quintin, and southward.

If we are right in referring *Mammillaria gabbii* here, this species was first collected by W. M. Gabb in southern Lower California in 1867 and was described by Dr. Engelmann as a new species but was not published. In 1894 Dr. Coulter published Engelmann's description, but used the name of *Cactus gabbii*. On a previous page, however, he published *Cactus brandegeei* which, if the same, takes precedence.

We have placed this species next to *Neomammillaria arida*, which is known to have nearly terete tubercles, while *N. brandegeei* is described as having angled tubercles, as they certainly are in herbarium specimens; whether this species has angled or terete tubercles in life we are in doubt.

We have not seen fresh fruit of this plant but Schumann describes it as white, which is unusual in this genus; it is also peculiar in bearing several small scales.

Illustrations: Blühende Kakteen 2: pl. 119; Schumann, Gesamtb. Kakteen Nachtr. 137. f. 34; Monatsschr. Kakteenk. 11: 153, as *Mammillaria brandegeei*.

7. *Neomammillaria gummifera* (Engelmann).

Mammillaria gummifera Engelmann in Wislizenus, Mem. Tour North. Mex. 105. 1848.

*Cactus gummifer** Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Depressed-globose, 8 to 12 cm. in diameter; tubercles light green, milky, somewhat 4-angled; axils of tubercles and spine-areoles white-tomentose when young; radial spines 10 to 12, ascending, white with brownish or even blackish tips, the lower ones stouter and longer than the others, often 2 to 2.5 cm. long and somewhat recurved; central spines 1 or 2, sometimes 4; flowers 3 cm. long, 52 to 25 mm. wide when fully open, brownish red outside; inner perianth-segments reddish white with dark red band in middle.

Type locality: Cosihuiriachi, Chihuahua.

Distribution: Northern Mexico.

This species was collected by Dr. A. Wislizenus in the state of Chihuahua, Mexico, about 1846. Specimens were sent to Dr. Engelmann at St. Louis, who described it in 1848 but without seeing flowers or fruit; two years afterward he described the flowers but the fruit is yet unknown. In 1894 Dr. J. M. Coulter redescribed the species, stating that it had never been re-collected. Professor Schumann in his Monograph does not recognize it, but refers it to his list of doubtful species. In 1908 Dr. Rose visited the type locality and obtained a single living specimen.

Illustrations: Cact. Mex. Bound. pl. 9, f. 18 to 20, as *Mammillaria gummifera*.

8. *Neomammillaria macdougalii* (Rose).

Mammillaria macdougalii Rose, Stand. Cycl. Hort. Bailey 4: 1982. 1916.

Usually low and flattened on top, but very old plants sometimes nearly globular and then 12 to 15 cm. in diameter with a carrot-shaped root; tubercles flattened dorsally, strongly angled, deep green; young areoles bearing white wool, but becoming naked in age; axils of tubercles often bearing long white wool; radial spines 10 to 12, white or somewhat yellowish, the lower ones a little stouter, brown or black at top or sometimes throughout; central spines 1 or 2, stout, yellowish, brown-tipped, similar to the radials; flowers 3.5 cm. long, cream-colored; outer perianth-segments short-fimbriate; fruit red, clavate, 3 cm. long.

Type locality: Near Tucson, Arizona.

Distribution: Southeastern Arizona.

Figure 65 is from a photograph of a plant collected by Dr. MacDougal in the Santa Catalina Mountains; figure 67 is from a photograph of another plant sent by Dr. MacDougal from the same region in November 1909.

* Coulter writes this name *Cactus gummiferus* (Contr. U. S. Nat. Herb. 3: 98.1894).

9. Neomammillaria heyderi (Mühlenpfordt).*Mammillaria heyderi* Mühlenpfordt, Allg. Gartenz. 16: 20. 1848.*Cactus heyderi* Kuntze, Rev. Gen. Pl. 1: 260. 1891.? *Mammillaria buchheimeana* Quehl, Monatsschr. Kakteenk. 2: 97. 1917.

Plant globose or somewhat flattened at apex; tubercles conic, 12 mm. long, when young bearing wool in their axils; young spine-areoles white-woolly; radial spines 20 to 22, white, setaceous, the lower ones stouter and longer; central spine solitary, brown at base and apex, 5 to 6 mm. long; flowers pinkish, the segments linear-oblong; fruit oblong, red.

Type locality: Not cited.

Distribution: Texas and northern Mexico.

Illustration: Schulz, Wild Fl. San Antonio pl. 13 in part, as *M. heyderi*.

Plate VIII, figure 2, shows a plant sent to Dr. Rose by Mrs. S. L. Pattison in 1921 which flowered in the New York Botanical Garden on April 21 of that year; figure a shows the fruit.




FIG. 67.—*Neomammillaria macdougallii*.




FIG. 68.—*Neomammillaria phymatothele*.

10. Neomammillaria hemisphaerica (Engelmann).*Mammillaria hemisphaerica* Engelmann in Wislizenus, Mem. Tour North. Mex. 105. 1848.*Mammillaria heyderi hemisphaerica* Engelmann, Proc. Amer. Acad. 3: 263. 1856.*Cactus heyderi hemisphaericus* Coulter, Contr. U. S. Nat. Herb. 3: 97. 1894.*Cactus hemisphaericus* Small, Fl. Southeast. U. S. 811. 1903.

Deep-seated in the soil, hemispheric, 8 to 12 cm. broad, dark green; tubercles only slightly angled, not very closely set, 1 to 1.5 cm. long, somewhat pointed, their axils nearly naked in the dormant stages; spine-areoles woolly when young, becoming glabrate in age; radial spines 9 to 13, widely spreading, acicular, the upper ones more delicate, 4 to 8 mm. long, brownish or smoky, often with black tips; central spine solitary, porrect, brown; flowers small, cream-colored, 1 to 1.5 cm. long; inner perianth-segments acute; filaments pinkish; style pinkish; stigma-lobes 6 to 10, greenish yellow; fruit slender, clavate, red, 1 to 1.5 cm. long.

Type locality: Below Matamoros on the Rio Grande.

Distribution: Southeastern Texas and northeastern Mexico.

This species was collected in 1846 by the St. Louis Volunteers in the Mexican War and taken back to Dr. George Engelmann; it flowered and he described it briefly in 1848 and in more detail in 1850. It was recently re-collected near Brownsville, Texas, just across the river from Matamoros by Robert Runyon and sent to us with a photograph taken in situ, here reproduced (plate VI, figure 2).

This species differs from *Neomammillaria applanata* in being less flattened and in having fewer spines and white flowers.

Cactus heyderi hemisphaericus, as treated by Coulter, must be a composite, the western and southern forms probably representing different species.

Illustrations: Cact. Mex. Bound. pl. 9, f. 15 to 17, as *Mammillaria heyderi hemisphaerica*.

Plate VIII, figure 6, shows a flowering plant from near Brownsville, Texas, collected by Robert Runyon; figure 3 shows a flowering and fruiting plant obtained by Dr. Rose at Laredo, Texas, in 1913, which flowered in the New York Botanical Garden, March 23, 1914; plate VI, figure 2, is from a photograph taken near Brownsville, Texas, by Robert Runyon in 1920.

11. *Neomammillaria applanata* (Engelmann).

Mammillaria applanata Engelmann in Wislizenus, Mem. Tour North. Mex. 105. 1848.

Mammillaria declivis Dietrich, Allg. Gartenz. 18: 235. 1850.

Mammillaria texensis Labouret, Monogr. Cact. 89. 1853.

Mammillaria heyderi applanata Engelmann, Proc. Amer. Acad. 3: 263. 1856.

Cactus texensis Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plants much flattened; tubercles somewhat angled, their axils naked; radial spines 10 to 18, the radials widely spreading, lower ones darker brown than upper; central spine one, porrect, dark brown; young spine-areoles very woolly; flower-buds pointed, greenish; outer perianth-segments greenish, lanceolate, acuminate margins not ciliate; inner segments 2.5 cm. long, cream-colored, lanceolate, acuminate, with a broad green stripe down the middle; filaments white, shorter than the style; stigma-lobes green; fruit scarlet, naked, 2.5 to 3.5 cm. long; seeds brown.

Type locality: Rocky plains on the Pieddenales, Texas.

Distribution: Central and southern Texas.

The description is based on plants flowering in cultivation. It is one of the earliest species to flower in the spring, beginning soon after the first of March; the fruit requires a full year to mature.

Mammillaria lindheimeri Engelmann, given by Hemsley (Biol. Centr. Amer. Bot. 1: 525. 1880) and by the Index Kewensis as a synonym of *M. texensis*, belongs here.

Neomammillaria applanata, *N. heyderi*, and *N. hemisphaerica* are closely related and may represent races of the same species.

Illustrations: Blanc, Cacti 66. No. 1116; Gartenflora 30: 412; Cact. Journ. 1: pl. for March; Meehan's Monthly 1: 4; Balt. Cact. Journ. 1: 138; 2: 259; Förster, Handb. Cact. ed. 2. 333. f. 35, as *Mammillaria applanata*; Ann. Rep. Smiths. Inst. 1908: pl. 9, f. 1; Gartenflora 29: 52, as *Mammillaria heyderi*; Schelle, Handb. Kakteenk. 263. f. 185; Blühende Kakteen 1: pl. 43; Cact. Mex. Bound. pl. 9, f. 4 to 14, as *Mammillaria heyderi applanata*.

Plate IX, figure 1, shows a plant in flower and fruit, collected by Dr. Rose on hills above Devil's River, Texas, in 1913, which flowered in the New York Botanical Garden, February 2, 1914.

12. *Neomammillaria phymatothele* (Berg).

Mammillaria phymatothele Berg, Allg. Gartenz. 8: 129. 1840.

Mammillaria ludwigii Ehrenberg, Linnaea 14: 376. 1840.

Cactus ludwigii Kuntze, Rev. Gen. Pl. 1: 260. 1891.


Cactus phymatothele Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Simple, subglobose, glaucous-green; axils of young tubercles bearing white wool, becoming naked; tubercles large, 4-sided; areoles when young white-woolly, in age naked; radial spines 7 to 20, grayish white, the three upper smaller, the central (Schumann says 1 or 2) recurved; flowers described by Schumann as carmine-colored.

Type locality: Mexico.

Distribution: Central Mexico.

We know this species only from the description and illustration.




M. E. Eaton del. 1 to 4
D. G. Passmore del. 5

A. Hoen & Co. Baltimore

1. Flowering and fruiting plant of *Neomammillaria applanata*.
2. Top of fruiting plant of *Neomammillaria karwinskiana*.
3. Top of flowering plant of *Neomammillaria karwinskiana*.
4. Flowering plant of *Neomammillaria macrantha*.
5. Flowering plant of *Neomammillaria mystax*.

Illustration. Blühende Kakteen 1: pl. 32, as *Mammillaria centricirrha* var.

Figure 69 is from a photograph sent us by L. Quehl; figure 68 is a reproduction of the illustration cited above; figure 70 shows a plant grown in the Missouri Botanical Garden in 1905 as *Cactus neumannianus*.





FIGS. 69 and 70.—*Neomammillaria phymatothele*.

13. *Neomammillaria magnimamma* (Haworth).

- Mammillaria magnimamma* Haworth, Phil. Mag. **63**: 41. 1824.
Mammillaria divergens De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 113. 1828.
Mammillaria gladiata Martius, Nov. Act. Nat. Cur. **16**: 336. 1832.
Mammillaria ceratophora Lehmann, Allg. Gartenz. **3**: 228. 1835.
Mammillaria recurva Lehmann in Pfeiffer, Enum. Cact. **15**. 1837.
Mammillaria hystrix Martius in Pfeiffer, Enum. Cact. **25**. 1837.
Mammillaria ebrenbergii Pfeiffer, Allg. Gartenz. **6**: 274. 1838.
Mammillaria microceras Lemaire, Cact. Aliq. Nov. **6**. 1838.
Mammillaria deflexispina Lemaire, Cact. Aliq. Nov. **6**. 1838.
Mammillaria versicolor Scheidweiler, Bull. Acad. Sci. Brux. **5**: 494. 1838.
? *Mammillaria conopsea* Scheidweiler, Bull. Acad. Sci. Brux. **5**: 496. 1838.
Mammillaria centricirrha Lemaire, Cact. Gen. Nov. Sp. **42**. 1839.
Mammillaria centricirrha macrothela Lemaire, Cact. Gen. Nov. Sp. **42**. 1839.
Mammillaria neumanniana Lemaire, Cact. Gen. Nov. Sp. **53**. 1839.
Mammillaria conopsea longispina Scheidweiler, Bull. Acad. Sci. Brux. **6**: 92. 1839.
Mammillaria pentacantha Pfeiffer, Allg. Gartenz. **8**: 406. 1840.
Cactus magnimamma Salm-Dyck in Steudel, Nom. ed. 2. **1**: 246. 1840.
Mammillaria subcurvata Dietrich, Allg. Gartenz. **12**: 232. 1844.
Mammillaria diadema Mühlenpfordt, Allg. Gartenz. **13**: 346. 1845.
Mammillaria krameri Mühlenpfordt, Allg. Gartenz. **13**: 347. 1845.
Mammillaria foersteri Mühlenpfordt, Allg. Gartenz. **14**: 371. 1846.
? *Mammillaria tetracentra* Otto in Förster, Handb. Cact. **214**. 1846.
Mammillaria bockii Förster, Allg. Gartenz. **15**: 50. 1847.
Mammillaria pazzanii Stieber, Bot. Zeit. **5**: 491. 1847.
Mammillaria divaricata Dietrich, Allg. Gartenz. **16**: 210. 1848.
Mammillaria hopferiana Linke, Allg. Gartenz. **16**: 329. 1848.
Mammillaria glauca Dietrich in Linke, Allg. Gartenz. **16**: 330. 1848.
Mammillaria centricirrha hopferiana Salm-Dyck, Cact. Hort. Dyck. 1849. 17, 123. 1850.
Mammillaria megacantha Salm-Dyck, Cact. Hort. Dyck. 1849. 123. 1850.
Mammillaria megacantha rigidior Salm-Dyck, Cact. Hort. Dyck. 1849. 18, 124. 1850.

- Mammillaria uberimamma* Monville in Labouret, Monogr. Cact. 520. 1853.
 ? *Mammillaria cirrosa** Poselger, Allg. Gartenz. 21: 94. 1853.
Mammillaria pachytele Poselger, Allg. Gartenz. 23: 17. 1855.
Mammillaria lactescens† Meinshausen, Wöchenschr. Gärt. Pflanz. 2: 117. 1859.
Mammillaria falcata Hortus in Förster, Handb. Cact. ed. 2. 345. 1885.
Mammillaria gebweileriana Haage in Förster, Handb. Cact. ed. 2. 358. 1885.
Mammillaria schmidtii Sencke in Förster, Handb. Cact. ed. 2. 376. 1885.
Mammillaria krameri viridis Haage in Förster, Handb. Cact. ed. 2. 372. 1885.
Cactus bockii Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus centricirrhus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus conopseus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus diadema Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus divergens Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus ebrenbergii Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus foersteri Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus gladiatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus glaucus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus kranzeri Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus lactescens Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus megacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus microceras Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus hystrix Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus divaricatus Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Lamarck, 1783.
Cactus neumannianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pazzanii Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pentacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus recurvus Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Miller, 5768.
Cactus versicolor Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus tetracentrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus subcurvatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Mammillaria centricirrha magnimamma Schumann, Gesamtb. Kakteen 582. 1898.
Mammillaria centricirrha divergens Schumann, Gesamtb. Kakteen 582. 1898.
Mammillaria centricirrha bockii Schumann, Gesamtb. Kakteen 582. 1898.
Mammillaria centricirrha recurva Schumann, Gesamtb. Kakteen 582. 1898.
Mammillaria centricirrha krameri Schumann, Gesamtb. Kakteen 582. 1898.

FIG. 71.—*Neomammillaria magnimamma*.FIG. 72.—*Neomammillaria macracantha*.

Globose, the larger plants 10 cm. in diameter, sometimes solitary but oftener cespitose with 25 in a cluster or more, very milky throughout; tubercles conic or somewhat flattened or faintly 4-angled, 1 cm. long, the axils when young densely woolly; spines 3 to 5, very unequal in length, the upper ones short and straight, the lower one or two 1.5 to 4.5 cm. long, recurved or incurved, all horn-colored, with black tips; flowers cream-colored; fruit clavate, 2 cm. long, crimson; seeds brownish.

* Schumann refers *Mammillaria cirrosa* (he spells it *M. cirrhosa*) doubtfully to *M. centricirrha*, but judging from the description it may belong elsewhere.

† Here was referred *M. neumanni glabrescens* Regel (Förster, Handb. Cact. ed. 2. 370. 1885).

Type locality: Not cited.

Distribution: Central Mexico.

This plant is very common in central Mexico, especially in the Valley of Mexico, about Tula, farther north, and also east of the City of Mexico. It makes large cespitose mounds, sometimes with many-headed branches, and has peculiar incurved spines and small flowers. It is frequently collected and has been shipped abundantly to Europe, where it has been much named, often from single joints. Our synonymy shows 34 specific names under *Mammillaria* and nearly as many under *Cactus*. Some writers have given these names varietal rank, so that this species now has about 100 names. It is a very characteristic plant and, while it may easily be confused with other species, yet, when clearly understood, its distinctness is evident.

Mammillaria zooderi was referred by Schumann (Gesamtb. Kakteen 582. 1898) as a synonym of *M. centricirrha* but the Index Kewensis Suppl. 5. cites Schelle (Handb. Kakteenk. 268. 1907), who gives it as a synonym of *M. centricirrha zooderi*. Neither the specific nor the varietal name can be considered published.

Schelle (Handb. Kakteenk. 266 to 268. 1907) lists 62 varietal names of *Mammillaria centricirrha*, all but one or two of which are based on species of the same name. Some of these perhaps are to be referred elsewhere, but we have listed them here as follows:

| | | | |
|--------------|----------------------|-------------|--------------|
| amoena | falcata | lactescens | posteriana |
| arietina | foersteri | lehmannii | pulchra |
| bockii | gebweileri | longispina | recurva |
| boucheana | gladiata | macracantha | schiedeana |
| ceratophora | glauca | magnimamma | schmidtii |
| cirrha | globosa | megacantha | spinosior |
| conopsea | grandidens | microceras | subcurvata |
| cristata | guilleminiana | montsii | tetracantha |
| deflexispina | hopfferiana | moritziana | uberimamma |
| destorum | hystrix | neumanniana | valida |
| de tampico | hystrix grandicornis | nordmannii | versicolor |
| diacantha | hystrix longispina | obconella | viridis |
| diadema | jorderi | pachythele | zooderi |
| clivaticata | krameri | pazzanii | zuccariniana |
| divergens | krameri longispina | pentacantha | |
| ehrenbergii | krausei | polygoria | |

The following garden names are listed by Schumann (Gesamtb. Kakteen 582. 1898) as belonging to this species:

| | | | |
|--------------|------------|------------|-------------|
| boucheana | hystrix | moritziana | tetracantha |
| destorum | jorderi | nordmannii | viridis |
| de tampico | lehmannii | obconella | zooderi |
| grandicornis | longispina | posteriana | |
| grandidens | montsii | spinosior | |

Illustrations: Hort. Belge 5: pl. 6, as *Mammillaria conopsea*; Reiche, Elem. Bot. f. 166, as *M. centricirrha*. Schelle's figure (Handb. Kakteenk. 268. f. 189) we are not able to place. The illustration in Blühende Kakteen (1: pl. 32) as *M. centricirrha* var. does not seem to be of this relationship.

Plate XI, figure 1, shows a small potted plant which flowered in the New York Botanical Garden, May 6, 1913. Figure 71 is from a photograph of a plant obtained on the pedregal near San Angel, Valley of Mexico, by O. Solis in 1919.

14. *Neomammillaria macracantha* (De Candolle).

Mammillaria macracantha De Candolle, Mém. Mus. Hist. Nat. Paris 17: 113. 1828.

Cactus macracanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus alternatus Coulter, Contr. U. S. Nat. Herb. 3: 95. 1894.

Mammillaria centricirrha macracantha Schumann, Gesamtb. Kakteen 582. 1898.

Depressed-globose, 2 to 3 cm. high, 6 to 15 cm. in diameter; axils of old tubercles naked, of young ones densely lanate; tubercles ovoid, somewhat 4-sided; young spine-areoles somewhat tomentose; spines 1 or 2, somewhat angled, elongated, the longest 5 cm. long (but not elongated in

greenhouse specimens), porrect or more or less reflexed, reddish in age; flowers dark pink, a little longer than the tubercles; perianth-segments linear, spreading; stigma-lobes 5 to 7, rose-colored.

Type locality: Mexico.

Distribution: San Luis Potosí.

Our description is based on plants from San Luis Potosí, Mexico, especially those collected by Mrs. Vera in 1912.

Schumann refers *Mammillaria macracantha* to *M. centricirrha* but it must be different. Rümpler refers to it also *M. zuccarinii*, but this has different flowers and we have recognized it as a species. *M. macrantha* (Förster, Handb. Cact. 189. 1894) is referred here.

Illustrations: De Candolle, Mém. Cact. pl. 9; Förster, Handb. Cact. ed. 2. 378. f. 38, as *Mammillaria macracantha*; Schumann, Gesamtb. Kakteen f. 93; Thomas, Zimmerkultur Kakteen 57; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 13, as *M. centricirrha macracantha*; (?) Engler and Prantl, Pflanzenfam. 3^{6a}: 170. f. 7, E, as *M. centricirrha*.




FIG. 72a.—*Neomammillaria macracantha*.

Plate ix, figure 4, shows a plant received from Kew in 1902, which flowered in the New York Botanical Garden on April 27, 1912. Figure 72 is a reproduction of the first illustration cited above; figure 72 is from a photograph of the plant distributed by the Kew Gardens in 1902 which flowered in the New York Botanical Garden in 1905.

15. *Neomammillaria johnstonii* sp. nov.

Plants large for the genus, globular to short-oblong, 15 to 20 cm. high, slightly depressed at apex; tubercles 1 to 1.5 cm. long, 4-angled throughout, somewhat bluish, naked in their axils, milky; spine-areoles when young short-floccose, in age glabrate, circular; radial spines 10 to 14, white, but with brown tips, somewhat spreading, stiff acicular; central spines 2, much longer and stouter than the radials, slightly diverging, bluish brown; flowers from near top of plant but from axils of old tubercles, campanulate, 2 cm. long; outer perianth-segments ovate-lanceolate, greenish white with a reddish-brown mid-rib; inner perianth-segments narrow, acuminate, white; filaments short, pinkish; style pinkish; stigma-lobes linear, 6 or 7, green.

1



2



1. *Coryphantha sulcata*, from Sabinas Texas.
2. *Neomammillaria runyonii*, from Monterey, Mexico.

Collected at San Carlos Bay, Sonora, Mexico, by Ivan M. Johnston ill 1921 (No. 4373) and flowered in Washington in April 1922 and April 1923.

Figure 72*b* is from a photograph of the type specimen.

16. *Neomammillaria melanocentra* (Poselger).

Mammillaria melanocentra Poselger, Allg. Gartenz. 23: 17. 1855.

Mammillaria erinacea Poselger, Allg. Gartenz. 23: 18. 1855.

Mammillaria valida Weber, Dict. Hort. Bois 806. 1898.

Short-cylindric, glaucous-green; tubercles in 8 and 13 spirals, strongly angled; radial spines 6, stout-subulate, 1.5 to 2 cm. long, brownish; central spines solitary, black, 2 to 3 cm. long, greatly overtopping the stem; flowers pinkish red, the segments linear, acute.




FIG. 72*b*.—*Neomammillaria johnstonii*.

Type locality: Near Monterey, Mexico.

Distribution: Mexico, but range unknown.

Illustration: Blühende Kakteen 3: pl. 129, as *Mammillaria melanocentra*. Figure 73 is a reproduction of the illustration cited above.

17. *Neomammillaria runyonii* sp. nov.

Plants deep-seated, depressed; tubercles milky, elongated, 1.5 cm. long, strongly 4-angled, their tips widely separated from each other, their axils long-woolly (never setose), especially when young, sometimes permanently so; young spine-areoles long-woolly, but in age glabrate; radial spines 6 to 8, slightly ascending, the outer ones stouter and often dark brown in color, the inner ones about half the length of the outer and nearly white; central spine solitary, brown to black, erect, 10 to 14 mm. long; flowers about 2 cm. long, purple; perianth-segments oblong; fruit red, clavate, 12 to 16 mm. long; seeds brown.

Collected on El Mirador, near Monterey, Mexico, by Robert Runyon in 1921.

Plate x, figure 2, is from a photograph of one of the plants Mr. Runyon originally brought from El Mirador.



18. *Neomammillaria sartorii* (J. A. Purpus).*Mammillaria sartorii* J. A. Purpus, Monatsschr. Kakteenk. 21: 50. 1911.

Globose to short-cylindric, 5 to 13 cm. in diameter, cespitose, very milky, bluish green; tubercles strongly 4-angled, pointed, 8 to 12 mm. long, their axils without bristles and in time without wool; spine-areoles circular when young, densely white-woolly but in age glabrate; spines 4 to 6, very unequal, 5 to 8 mm. long, whitish or sometimes brownish, the central spine solitary; flowers 1.5 to about 2 cm. long, deep carmine; perianth-segments oblong, apiculate, the tip dry, the outer ciliate, the inner serrulate; stamens and style purplish above; stigma-lobes 4, purple, short; fruit carmine; seeds brown.

Type locality: Barranca de Panoaya, Vera Cruz, Mexico.

Distribution: Mountains of Vera Cruz, 300 to 600 meters altitude.

Our description of this interesting and variable little plant is drawn from specimens sent to us by Dr. C. A. Purpus in 1920, collected at the type locality. There the plant grows

FIG. 73.—*Neomammillaria melanocentra*.FIG. 74.—*Neomammillaria seitziana*.

among rocks in rich humus of the decaying leaves in half shade or in the sun. It is very different from any other *Neomammillaria* which we have seen; the tubercles are copiously milky and the slightest bruise causes the white milk to ooze out. It flowered in Washington in April 1923.

Dr. C. A. Purpus writes that this species is common in many of the barrancas of Vera Cruz and that it is very variable. When first described two forms (*brevispina* and *longispina*) were characterized.

The species was named for Florantino Sartorius (1837-1908) who assisted Dr. Purpus for many years in his botanical expeditions. He was a son of Carlos Sartorius (1795-1872), a distinguished scientist who went to Mexico about 1825, where he made large collections of plants. Mr. W. Botting Hemsley (Biol. Centr. Amer. Bot. 4: 123) states that his herbarium was left to the Smithsonian Institution, but no record of this gift can now be found nor can any of his plants be found in the U. S. National Herbarium.

Here may or may not belong *Mammillaria rebsamiana* (Cact. Journ. 2: 176), advertised as a new discovery by Louis Murillo, who lived at Jalapa, Mexico.

Illustration: Monatsschr. Kakteenk. 21: 51, as *Mammillaria sartorii*.

Figure 75 is from a photograph showing two plants sent from the type locality of *Mammillaria sartorii* by Dr. Purpus in 1920.

19. Neomammillaria seitziana (Martius).

Mammillaria seitziana Martius in Pfeiffer, Enum. Cact. 18. 1837.
Mammillaria foveolata Mühlenpfordt, Allg. Gartenz. 14: 372. 1846.
Cactus foveolatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus seitzianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Solitary or somewhat proliferous at base, cylindric, 12 cm. high; tubercles green, conic, somewhat angled; axils of tubercles woolly; areoles at first white-woolly, becoming glabrate; spines 4,* the upper and lower longer than the lateral; flowers rose-colored, about 25 mm. long; outer perianth-segments olive colored; inner perianth-segments linear, lanceolate, white, nerved with red; stamens white; stigma-lobes 6.

Type locality: Ixmiquilpan, Mexico.

Distribution: State of Hidalgo.

We have not seen this species and hence our description is compiled.




FIG. 75.—*Neomammillaria sartorii*.

Mammillaria senckena and *M. senckei* are two names listed as synonyms of this species, but we do not find that they have ever been published.

Illustration: Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 8, as *Mammillaria seitziana*. Figure 74 is reproduced from the illustration cited above.

20. Neomammillaria ortegae sp. nov.

Simple to short-clavate, 5 to 8 cm. in diameter, light green, lactiferous; tubercles rather short (8 to 10 mm. long), broader at base, obscurely 4-angled, somewhat pointed, very woolly but not setose in their axils; spines all radial, 3 or 4, more commonly (sometimes with 1 or 2 small additional spines or bristles, perhaps deciduous), spreading, straw-colored, 6 to 10 mm. long; flowers small; fruit clavate, 1 cm. long; seeds numerous, small, angled, brown.

Collected by J. G. Ortega in Sinaloa, Mexico, in 1921 and 1922.

Figure 76 shows the type specimens as photographed in the U. S. National Museum under the direction of A. J. Olmstead.

* Schumann says central spines yellow.

21. *Neomammillaria meiacantha* (Engelmann).

Mammillaria meiacantha Engelmann, Proc. Amer. Acad. 3: 263. 1856.
Cactus meiacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Somewhat depressed, 12 cm. broad or more; tubercles milky, bluish green, more or less angled, somewhat flattened dorsally, their axils naked; spines 5 to 9, ascending, pale flesh-colored, the tips darker, the lower a little stouter than the upper; central spines porrect, similar to but a little stouter than radials and often subradial; spine-areoles short-woolly at first; flowers not very abundant, at least on cultivated plants; inner perianth-segments white with a pink stripe along inside of midrib, one-fourth its width, greenish brown on outside; filaments white; style pink; stigma-lobes yellow; fruit scarlet, 22 mm. long; seeds brownish.

Type locality: Western Texas and New Mexico.

Distribution: Texas, New Mexico, and northern Mexico.

According to Dr. Engelmann, this species was first obtained in New Mexico by the Missouri Volunteers in 1847 and it has frequently been collected since that time. In Mexico it extends as far south as Zacatecas, but develops into some unusual forms. It was repeatedly collected in Zacatecas by F. E. Lloyd in 1908.




FIG. 76.—*Neomammillaria ortegae*.

Illustrations: Blühende Kakteen 1: pl. 47*; Blanc, Cacti 71. No. 1388; Cycl. Amer. Hort. Bailey 2: f. 1357. Stand. Cycl. Hort. Bailey 4: f. 2316; West Amer. Sci. 13: 39; Schelle, Handb. Kakteenk. 258. f. 190; Cact. Mex. Bound. pl. 9, f. 1 to 3; Cact. Journ. 1: pl. for October, as *Mammillaria meiacantha*.

Figure 77 shows the plant illustrated in the Mexican Boundary Report as cited above.

22. *Neomammillaria scrippsiana* sp. nov.

Globose or becoming short-cylindric, 6 cm. high; tubercles milky, in 26 rows, bluish green, very woolly in axils when young; spine-areoles very woolly at first; radial spines 8 to 20, slender, pale with reddish tips; central spines generally 2, a little longer than radials, brown throughout, slightly divergent; flowers borne near top of plant but not in axils of youngest tubercles, about 1 cm. long, pinkish, with margins of perianth-segments paler; anthers pinkish; stigma-lobes about 6, recurved, cream-colored.

Collected by Dr. Rose in the barranca of Guadalajara, Jalisco, in September 1903 (No. 871, type). The plant has flowered repeatedly in Washington since April 1906. Specimens were afterward collected near the same place by C. R. Orcutt. It is named in honor of E. W. Scripps, the founder of Science Service and The Scripps Institution for Biological Research of the University of California.

Figure 78 is from a photograph of the type specimen.

* This plate is labeled *Mammillaria meionacantha*, but described under *M. meonacantha*.

23. *Neomammillaria gigantea* (Hildmann).

Mammillaria gigantea Hildmann in Schumann, Gesamtb. Kakteen 578. 1898.

Solitary or cespitose, depressed-globose, 10 cm. high, 15 to 17 cm. in diameter; axils of tubercles lanate; radial spines 12, subulate, white, 3 mm. long; central spines 4 to 6, stout, 2 cm. long, curved, yellowish brown; flowers yellowish green.

Type locality: Guanajuato, Mexico.

Distribution: Known only from the type locality.

Mammillaria macdowellii Heese and *M. guanajuatensis* Runge are two names referred here by Schumann (Gesamtb. Kakteen 578. 1898), but they were not published.

Plate XI, figure 3, shows a plant in fruit, collected by Dr. Safford at the type locality.

24. *Neomammillaria peninsularis* sp. nov.

Plants solitary or in clusters, deeply seated in the ground, more or less flat-topped, bluish green, the stems and tubercles very milky; tubercles erect, pointed, 4-angled, pale green; radial




FIG. 77.—*Neomammillaria meiacantha*.




FIG. 78.—*Neomammillaria scripsiana*.

spines 4 to 8, nearly erect, short and pale with brown tips, one sometimes nearly central; axils of tubercles bearing long wool but in age naked; flowers 1.5 cm. long, arising from old tubercles but near the center; outer perianth-segments narrow, reddish; inner perianth-segments narrow, acuminate, green or light yellow with erose margins; stamens pale; style longer than stamens; stigma-lobes green, linear.

Collected by Dr. Rose at Cape San Lucas, Lower California, March 23, 19 " (No. 16377).

25. *Neomammillaria flavovirens* (Salm-Dyck).

Mammillaria flavovirens Salm-Dyck, Cact. Hort. Dyck. 1849. 117. 1850.

Either solitary or somewhat cespitose, globose or short-cylindric, 6 to 8 cm. high, light or yellowish green; tubercles somewhat 4-angled; axils naked; radial spines 5, slender, subulate; central spines solitary, porrect; flowers white, streaked with rose.

Type locality: Not cited.

Distribution: Mexico.

The above description is compiled, since the species is not otherwise known to us.

Mammillaria flavovirens cristata Salm-Dyck (Cact. Hort. Dyck. 1849. 16. 1850) is only a name.

The name *Mammillaria daedalea viridis* Fennel is given by Labouret (Monogr. Cact. 100. 1853) as a synonym of *M. flavovirens*.

26. *Neomammillaria sempervivi* (De Candolle).

Mammillaria sempervivi De Candolle, Mém. Mus. Hist. Nat. Paris 17: 114. 1828.

Mammillaria sempervivi tetracantha De Candolle, Mém. Mus. Hist. Nat. Paris 17: 114. 1828.

Mammillaria caput-medusae Otto in Pfeiffer, Enum. Cact. 22. 1837.

Mammillaria diacantha Lemaire, Cact. Aliq. Nov. 2. 1838.

Mammillaria sempervivi laeteviridis Salm-Dyck, Cact. Hort. Dyck. 1849. 113. 1850.

Mammillaria caput-medusae centrispina Salm-Dyck in Labouret, Monogr. Cact. 91. 1853.

Mammillaria caput-medusae crassior Salm-Dyck in Labouret, Monogr. Cact. 91. 1853.

Mammillaria caput-medusae tetracantha Salm-Dyck in Labouret, Monogr. Cact. 91. 1853.

Cactus sempervivi Kuntze, Rev. Gen. Pl. 1: 261. 1891.




FIG. 79.—*Neomammillaria sempervivi*.




FIG. 80.—*Neomammillaria polythele*.

Solitary or somewhat cespitose, flattened above, narrowed below; axils of tubercles very woolly; tubercles short, milky, angled; spine-areoles very woolly when young, but glabrate in age; radial spines 3 to 7, short, white, caducous; central spines 2, ascending, brownish, stoutish; flowers dull white with reddish lines; inner perianth-segments acute, spreading.

Type locality: Mexico.

Distribution: Central Mexico.

Dr. Rose collected what he took to be this species in the Barranca Sierra de la Mesa, Hidalgo, Mexico, in 1905, but this plant differs somewhat from De Candolle's illustration. The central spines, while generally 2, are sometimes 3 and are not so stout; the radial spines are deciduous, as they should be in this species. It flowered once at Washington.

An examination of the original description of *Mammillaria caput-medusae* suggests the probability that this species is identical with *Mammillaria sempervivi*. The two names appeared in collections in 1829 and may have come from a common source. Indeed, Schumann credits T. Coulter with having obtained *M. caput-medusae*, while we know that *M. sempervivi* was based on Coulter's plant and, then, too, Pfeiffer refers *M. sempervivi* as a synonym of *M. caput-medusae*. Knippel's illustration of *M. caput-medusae* (pl. 19) seems to be referable here. Nicholson states that *M. caput-medusae* is only a form of this species.

Mammillaria staurotypa (Förster, Handb. Cact. 221. 1846), credited to Scheidweiler by Schumann and referred by him as a synonym of *M. caput-medusae*, seems never to have been described but may belong here.

The two varieties of *Mammillaria caput-medusae*, *tetracantha* and *hexacantha*, given by Salm-Dyck (Cact. Hort. Dyck. 1844. 10. 1845) are without description. The first was afterwards described by Labouret.

Illustrations: De Candolle, Mém. Cact. pl. 8; Förster, Handb. Cact. ed. 2. 344. f. 36; Schumann, Gesamtb. Kakteen 589. f. 95; Dict. Gard. Nicholson 4: 565. f. 38; Suppl. 518. f. 556; Watson, Cact. Cult. 175. f. 70, as *Mammillaria semper-vivi*; Schelle, Handb. Kakteenk. 270. f. 192; Succulenta 5: 51, as *M. caput-medusae*.

Figure 79 is a reproduction of the first illustration cited above.

27. *Neomammillaria obscura* (Hildmann).

Mammillaria obscura Hildmann, Monatsschr. Kakteenk. 1: 52. 1891.

Solitary, depressed-globose, blackish green; axils woolly; tubercles arranged in 13 and 2! spirals, angled, stout, woolly in their axils but not setose; radial spines 6 to 8, subulate, white, unequal, the upper ones shorter than the lower; central spines 2 to 4, the lower one slightly curved, black; flowers small, yellowish white.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

The plant is known to us only from description and illustration.

Seeds of this species were introduced into Germany from Mexico about 1885 by Mr. Droege and flowers were obtained in 1891.

The earlier name, *Mammillaria obscura* Scheidweiler (Förster, Handb. Cact. 213. 1846), but used only as a synonym and for some other plant, does not interfere with our present use of the name.

Illustration: Monatsschr. Kakteenk. 1: facing 52, as *Mammillaria obscura*.

28. *Neomammillaria crocidata* (Lemaire).

Mammillaria crocidata Lemaire, Cact. Aliq. Nov. 9. 1838.

Mammillaria webbiana Lemaire, Cact. Gen. Nov. Sp. 45. 1839.

Cactus crocidatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus webbianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plant globose or a little depressed, 5 to 6 cm. in diameter; radial spines 6 or 7, dark brown or nearly black; central spines none; axils of tubercles in young plant densely woolly; flowers from axils of old tubercles near the top of plant, small, reddish purple, 12 to 14 mm. long; outer perianth-segments ciliate; inner perianth-segments acuminate; filaments, style, and stigma-lobes reddish; stigma-lobes 3 or 4; fruit not seen.

Type locality: Mexico.

Distribution: Central Mexico.

Described here from plants collected by Dr. Rose near Queretaro, Mexico, in 1906, which flowered in August and September 1908, and again in April 1909 (No. 1072). Our specimen has more spines than the original *M. crocidata*; it is also near *M. carnea* but with different colored stigma-lobes; its tubercles are about 6 mm. high.

Schumann places this species near *M. carnea* and among the cylindric species, but it was originally described as depressed.

Mammillaria crocidata quadrispina Pfeiffer and Salm-Dyck, mentioned by Förster (Handb. Cact. 220. 1846) as a rare form and afterwards briefly described by Labouret (Monogr. Cact. 93. 1853), may or may not belong here.

Plate VII, figure 5, shows a flowering plant collected by Dr. Rose in Queretaro in 1906 and painted in the New York Botanical Garden, September 5, 1911.

29. Neomammillaria polythele (Martius).

- Mammillaria polythele* Martius, Nov. Act. Nat. Cur. **16**: 328. 1832.
Mammillaria quadrispina Martius, Nov. Act. Nat. Cur. **16**: 329. 1832.
Mammillaria columnaris Martius, Nov. Act. Nat. Cur. **16**: 330. 1832.
Mammillaria affinis De Candolle, Mém. Cact. **11**. 1834.
Mammillaria setosa Pfeiffer, Allg. Gartenz. **3**: 379. 1835.
Mammillaria polythele quadrispina Salm-Dyck in Walpers, Repert. Bot. **2**: 271. 1843.
Mammillaria polythele columnaris Salm-Dyck in Walpers, Repert. Bot. **2**: 271. 1843.
Mammillaria polythele setosa Salm-Dyck, Cact. Hort. Dyck. 1844. 9. 1845.
Mammillaria polythele hexacantha Salm-Dyck, Cact. Hort. Dyck. 1849. 15. 1850.
Mammillaria polythele latimamma Salm-Dyck, Cact. Hort. Dyck. 1849. 112. 1850.
Cactus affinis Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
Cactus quadrispinus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Cactus setosus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Cactus polythele Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
? *Mammillaria hidalgensis* Purpus, Monatsschr. Kakteenk. **17**: 118. 1907.

Elongated, cylindric, often 3 to 5 dm. high, 7 to 10 cm. in diameter; tubercles milky, in about 21 spirals, 10 to 12 mm. long, nearly terete, somewhat narrowed toward apex, dull green; axils of young tubercles densely long-woolly, the wool nearly covering the top of the plant, in age becoming naked; spines 2 to 4, sometimes 6, all radial, somewhat spreading, 1 to 2.5 cm. long, reddish, straight or a little curved; flowers from near top of plant, reddish, 8 to 10 mm. long; perianth-segments narrow, acuminate; fruit red, clavate; seeds small, brownish.

Type locality: Mexico.

Distribution: State of Hidalgo.

In 1905 Dr. Rose collected living plants of this species near Ixmiquilpan. It is a rather striking plant, growing very tall and flowering near the top.

Schumann places this species in the Section *Hydrochylus*, in which the sap is watery, but Martius in his original description says definitely that it is milky.

Mammillaria aciculata Otto (Pfeiffer, Enum. Cact. 29. 1837; *M. polythele aciculata* Salm-Dyck, Cact. Hort. Dyck. 1844. 9. 1845) is referred here by Schumann but should be excluded; it came from the cold regions of Mexico and was described as having 20 white slender radial spines.

Mammillaria columnaris minor Martius and *M. quadrispina major*, mentioned by Förster (Handb. Cact. 214, 215. 1846), probably belong here.

Mammillaria cataphracta Martius was given by Pfeiffer (Enum. Cact. 11. 1837) as a synonym of *M. affinis* and by Salm-Dyck (Hort. Dyck. 155. 1834) as a synonym of *M. angularis*.

Illustrations: Nov. Act. Nat. Cur. **16**: pl. 19, as *Mammillaria polythele*; Monatsschr. Kakteenk. **17**: 119; Möllers Deutsche Gärt. Zeit. **25**: 47. f. 8, No. 10, as *M. hidalgensis*; De Candolle, Mém. Cact. pl. 6, as *M. affinis*; Abh. Bayer. Akad. Wiss. München **2**: pl. 1, I. f. 2, as *M. columnaris*.

Figure 80 is from a photograph of a plant collected in the state of Hidalgo in 1905 which has heretofore passed as *Mammillaria hidalgensis*.

30. Neomammillaria carnea (Zuccarini).

- Mammillaria carnea* Zuccarini in Pfeiffer, Enum. Cact. 19. 1837.
Mammillaria subtetragona Dietrich, Allg. Gartenz. **8**: 169. 1840.
Mammillaria aeruginosa Scheidweiler, Allg. Gartenz. **8**: 338. 1840.
Mammillaria pallescens Scheidweiler, Allg. Gartenz. **9**: 42. 1841.
Mammillaria villifera carnea Salm-Dyck, Cact. Hort. Dyck. 1849. 16. 1850.
Mammillaria villifera aeruginosa Salm-Dyck, Cact. Hort. Dyck. 1849. 16. 1850.
*Mammillaria villifera cirrosa** Salm-Dyck, Cact. Hort. Dyck. 1849. 115. 1850.
Cactus aeruginosus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
Cactus carneus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
Cactus pallescens Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Cactus subtetragonus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Mammillaria carnea cirrosa Gürke, Blühende Kakteen **1**: under pl. 60. 1905.
Mammillaria carnea aeruginosa Gürke, Blühende Kakteen **1**: under pl. 60. 1905.

Plants solitary, cylindric, 8 to 9 cm. high; tubercles 4-angled, milky, their axils woolly, the upper ones erect; spines 4, straight, reddish, the lower one 10 mm. long, twice as long as the other; flowers

* Förster (Handb. Cact. ed. 2. 342. 1885) spells this name, *cirrrosa*.

borne in the old axils; outer perianth-segments nearly 2 cm. long, nearly erect, flesh-colored; fruit pear-shaped, obtuse, bright red.

Type locality: Ixmiquilpan, Mexico.

Distribution: Central and southern Mexico.

Mammillaria villifera Otto, referred here by Schumann, must belong elsewhere, since the axils of the tubercles bear setae.

Illustrations: Blühende Kakteen 1: pl. 60; Monatsschr. Kakteenk. 28: 59; Schelle, Handb. Kakteenk. 271. f. 193, as *Mammillaria carnea*.

Plate VII, figure 7, shows a plant collected by Dr. Rose at Tehuacán in 1906, which flowered in the New York Botanical Garden, May 4, 1912. Figure 81 is from a photograph of a plant collected by Dr. Rose at Tehuacán in 1905.




FIG. 81.—*Neomammillaria carnea*.




FIG. 82.—*Neomammillaria lloydii*.

31. *Neomammillaria lloydii* sp. nov.

Plant-body at first flattened but in cultivation becoming elongated, sometimes 10 cm. long, 6 to 7 cm. in diameter; axils of young tubercles only slightly woolly; tubercles milky, small, numerous, 4-angled, woolly when quite young; radial spines 3 or 4, ascending, glabrous, the uppermost one red or dark brown, the others whitish, 2 to 5 mm. long; central spines none; flowers in a ring near center of plant; outer perianth-segments dark red with light or colored margins; inner perianth-segments white with a tinge of red, and dark-red central stripes, not ciliate, apiculate, spreading above; filaments pale below, pinkish above; style pinkish above.

Collected by F. E. Lloyd in the State of Zacatecas, Mexico, in '909, and flowered in Washington in 1911, 1912 (March), and 1915 (April).

Figure 82 is from a photograph of the type plant (Lloyd, No. 55).

32. *Neomammillaria zuccariniana* (Martius).

Mammillaria zuccariniana Martius, Nov. Act. Nat. Cur. 16: 331. 1832.

Globose to elongated-cylindric, 8 to 20 cm. long, bluish green, milky; areoles and axils of young tubercles filled with white wool; radial spines wanting or represented by very stout bristles; central spines 2 to 4, black, unequal, 2 to 12 mm. long, spreading; flowers about x cm. long, with a broad open throat; outer perianth-segments brownish, acute; inner perianth-segments lanceolate, acute, entire, magenta-colored; filaments purplish; stigma-lobes 3 or 4, purplish, broad, truncate; fruit red, 10 mm. long; seeds brownish.

Type locality: Mexico.

Distribution: San Luis Potosí, Mexico.

We have had this plant in cultivation for a number of years; Dr. E. Palmer obtained it near San Luis Potosí in 1905 (No. 590); it was also collected by Mrs. Vera from the same locality in 1912.

Illustration: Martius, Nov. Act. Nat. Cur. **16**: pl. 20, as *Mammillaria zuccariniana*.

Figure 83 is from a photograph of a plant collected by Dr. E. Palmer near Alvarez, San Luis Potosí, May 1905, which afterwards flowered in Washington, D. C.

33. *Neomammillaria formosa* (Galeotti).

Mammillaria formosa Galeotti in Scheidweiler, Bull. Acad. Sci. Brux. **5**: 497. 1838.

Mammillaria formosa microthele Salm-Dyck, Cact. Hort. Dyck. 1849. 87. 1850.

Mammillaria formosa dispicula Monville in Labouret, Monogr. Cact. 60. 1853.

Mammillaria formosa gracilispina Monville in Labouret, Monogr. Cact. 60. 1853.

Mammillaria formosa laevior Monville in Labouret, Monogr. Cact. 60. 1853.

Cactus formosus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.

Somewhat clavate, sunken at the apex; axils lanate; tubercles spirally arranged, obtusely 4-angled, light green; areoles naked; radial spines 20 to 22, white, rigid, radiating; central spines 6, spreading, thickened at base, at first flesh-colored at base, black at tip, becoming black throughout or grayish; flowers red.

Type locality: Near San Felipe.

Distribution: San Luis Potosí, Mexico, according to Hemsley.

Dr. Safford has referred here a plant collected by Dr. E. Palmer at San Luis Potosí which may be the plant which is passing under this name, but it does not seem to answer the original descriptions.

Illustration: Garten-Zeitung **4**: 182. f. 42, No. 11, as *Mammillaria formosa*.




FIG. 83.—*Neomammillaria zuccariniana*.




FIG. 84.—*Neomammillaria compressa*.

34. *Neomammillaria compressa* (De Candolle).

Mammillaria compressa De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 112. 1828.

Mammillaria subangularis De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 112. 1828.

Mammillaria triacantha De Candolle, Mém. Mus. Hist. Nat. Paris **17**: 113. 1828.

Mammillaria cirrhifera Martius, Nov. Act. Nat. Cur. **16**: 334. 1832.


Mammillaria angularis Link and Otto in Pfeiffer, Enum. Cact. **12**. 1837.

Mammillaria cirrhifera angulosior Lemaire, Cact. Gen. Nov. Sp. 95. 1839.

Mammillaria longiseta Mühlenpfordt, Allg. Gartenz. **53**: 346. 1845.

- Mammillaria cirrifer longiseta* Salm-Dyck, Cact. Hort. Dyck. 1849. 18. 1850.
Mammillaria squarrosa Meinshausen Wöchenschr. Gärten. Pflanz. 2: 116. 1859.
Cactus cirrhifer Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus compressus Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Salisbury, 1796.
Cactus longisetus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus squarrosus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus subangularis Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus triacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
 ? *Mammillaria angularis fulvispina* Schumann, Gesamtb. Kakteen 576. 1898.
Mammillaria angularis longiseta Salm-Dyck in Schumann, Gesamtb. Kakteen 576. 1898.
Mammillaria angularis triacantha Salm-Dyck in Schumann, Gesamtb. Kakteen 576. 1898.
Mammillaria angularis compressa Schumann, Gesamtb. Kakteen 577. 1898.
Mammillaria oettingenii Zeissold, Monatsschr. Kakteenk. 8: 10. 1898.
Mammillaria kleinschmidtiana Zeissold, Monatsschr. Kakteenk. 8: 21. 1898.

Growing in large clumps, cylindric, pale bluish green; axils of tubercles white-woolly, setose; tubercles short, compressed laterally, keeled below, more rounded above; young spine-areoles white-woolly; principal spines 4, sometimes with 1 to 3 very short accessory ones from the lower part of the areole; lower spine much longer, spreading or recurved, 5 to 6 cm. long, somewhat angled; all spines pale, more or less tinged with brown, with dark tips; flower small, pinkish, so to 52 mm. long; outer perianth-segments acute, somewhat ciliate; inner perianth-segments narrow, acuminate, with spreading tips; stamens and style pale; stigma-lobes 5, linear; fruit clavate, red; seeds brown.



FIGS. 85 and 86.—*Neomammillaria compressa*.

Type locality: Mexico.

Distribution: Central Mexico.

Our description is drawn largely from specimens which flowered in March 1908 and which were collected by Dr. Rose at Higuerillas, Queretaro, in 5905. Dr. Rose also found this species very abundant in the deserts of Queretaro and living specimens brought back by him have frequently flowered both in New York and Washington. These are identical with plants sent from Berlin, labeled *Mammillaria angularis longiseta*. The species as here treated is variable and more exhaustive field work might require some modifications in the description.

The varieties *Mammillaria cirrifer major* and *M. cirrifer fulvispina* (Salm-Dyck, Cact. Hort. Dyck. 1844. 11. 1845) are without descriptions. The two varieties *M. cirrifer albispinga* and *M. centricirra macrothela* were listed as synonyms of *M. subangularis* by Walpers (Repert. Bot. 2: 272. 1843). To *M. subangularis* is also referred *M. subcirrifer* by Förster (Handb. Cact. 234. 1846).

Here doubtless belongs *Mammillaria angularis fulvescens* (Salm-Dyck, Cact. Hort. Dyck. 1849. 18. 1850).

Illustrations: Schelle, Handb. Kakteenk. 264. f. 187 (?); Möllers Deutsche Gärt. Zeit. 25: 486. 1. 20; Krook, Handb. Cact. 38 (?); Gartenwelt 15: 410; Contr. U. S. Nat. Herb. 10: pl. 16, f. B, as *Mammillaria angularis*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 8, as *M. angularis compressa*; Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 7, as *M. cirrhifera*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 28, as *M. angularis rufispina*; Blanc, Cacti 67. f. 1170; Cact. Journ. 1: pl. for March; 2: 7, 93, as *M. cirrhifera longispina*.

Plate VIII, figure 4, shows a plant collected by Dr. Rose in Queretaro in 1905, which flowered in the New York Botanical Garden, March 17, 1913. Figure 84 is from a photograph of a plant sent by Dr. E. Palmer from San Luis Potosí in 1905; figure 8 is from a photograph of a plant collected by Dr. Rose near Higuierillas, Mexico, in 1905; figure 86 is from a photograph of this plant growing in the open in the Huntington Collection, southern California; figure 88 is from a photograph of a plant sent by Dr. C. A. Purpus from Minas de San Rafael in 1910; figure 87 shows spines from the plant collected by Dr. Rose at Ixmiquilpan in 1905.




FIG. 87.—Spine-cluster of *N. compressa*.




FIG. 88.—*Neomammillaria compressa*.

35. *Neomammillaria mystax* (Martius).

- Mammillaria mystax* Martius, Nov. Act. Nat. Cur. 16: 332. 1832.
Mammillaria leucotricha Scheidweiler, Allg. Gartenz. 8: 338. 1840.
Mammillaria xanthotricha Scheidweiler, Allg. Gartenz. 8: 338. 1840.
Mammillaria mutabilis Scheidweiler, Allg. Gartenz. 9: 43. 1841.
Mammillaria funkii Scheidweiler, Allg. Gartenz. 9: 43. 1841.
Mammillaria autumnalis Dietrich, Allg. Gartenz. 16: 297. 1848.
Mammillaria mutabilis xanthotricha Salm-Dyck, Cact. Hort. Dyck. 1849. 17, 120. 1850.
Mammillaria maschalacantha Monville in Labouret, Monogr. Cact. 106. 1853.
Mammillaria maschalacantha leucotricha Monville in Labouret, Monogr. Cact. 106. 1853.

Mammillaria maschalacantha xantotricha Monville in Labouret, Monogr. Cact. 106. 1853.

Cactus funckii Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus maschalacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus leucotrichus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus mutabilis Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus mystax Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus xanthotrichus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Globose to short-cylindric, 7 to 15 cm. high, flat-topped; tubercles in as many as 34 rows, thickly set, full of milk which freely flows when pricked or cut; radial spines 8 to 10, small, white; central spines 4, 3 about twice as long as the radial ones, the other much elongated, 6 to 7 cm. long; flowers 1.5 to 2 cm. long, appearing in 2 or 3 rows, very abundant; inner perianth-segments dark red, 12 mm. long; stigma-lobes 4 or 5, greenish; fruit red, 2 to 2.5 cm. long.

Type locality: Mexico. According to Hemsley, Karwinsky's plant, which is the type, came from Ixmiquilpan and San Pedro Nolasco at about 6,000 feet altitude.

Distribution: Highlands of southern central Mexico.

This species is characterized by the long, erect, central spines which overtop the plant in the wild state; in cultivation these elongated spines do not always occur. The species is common in cultivation; in collections it is usually known as *Mammillaria mutabilis*.



FIG. 89.—*Neomammillaria mystax*.



FIG. 90.—*Neomammillaria petterssonii*.

Mammillaria krauseana, a name from Gruson's Catalogue, is cited by Schumann (Gesamtb. Kakteen 595. 1898) as a synonym of *M. mutabilis*.

Mammillaria meschalacantha Hortus (Salm-Dyck, Cact. Hort. Dyck. 1844. 10. 1845), according to the Index Kewensis, is a misspelling for *M. maschalacantha*.

Mammillaria maschalacantha dolichacantha Monville was given as a doubtful synonym of *M. maschalacantha* by Labouret (Monogr. Cact. 106. 1853).

Mammillaria mutabilis autumnalis (Monatsschr. Kakteenk. 30: February 1920) is offered for sale by Grässner.

Mammillaria mutabilis laevior Salm-Dyck (Cact. Hort. Dyck. 1849. 17, 120. 1850), with *M. leucocarpa* Scheidweiler as a synonym, was given as a variety of *M. mutabilis*, but it was not described. *M. xanthotricha laevior* Salm-Dyck (Cact. Hort. Dyck. 1844. 11. 1845), also undescribed, seems to be the same.

Schumann refers here *Mammillaria cirrhifera*, but certainly Pfeiffer's illustration (Abbild. Beschr. Cact. 1: pl. 7) with its long, curved, radial spines and no centrals is very different; we have referred it to *Neomammillaria magnimamma*.

Illustrations: Nov. Act. Nat. Cur. 16: pl. 21; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 17, as *Mammillaria mystax*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 1; Karsten and Schenck, Vegetationsbilder 1: pl. 44; Schelle, Handb. Kakteenk. 272. f. 195, as *M. mutabilis*; Schelle, Handb. Kakteenk. 273. f. 196, as *M. mutabilis longispina*.

Plate 1x, figure 5, shows a plant collected by Dr. Rose at Tehuacán which flowered and fruited in Washington in 1907. Figure 89 is from a photograph of a potted plant obtained by Dr. Rose at Tehuacán in 1905.

36. *Neomammillaria petterssonii* (Hildmann).

Mammillaria petterssonii Hildmann, Deutsche Garten-Zeitung 1886: 185. 1886.
Mammillaria heeseana McDowell, Monatsschr. Kakteenk. 6: 125. 1896.

Plants rather large for this genus, cylindric, 2 dm. high or more, very spiny; tubercles arranged in 13 or 21 spirals, terete, setose in their axils; radial spines 10 to 12, white, with black tips; central spines, the longer ones 4.5 cm. long; flowers unknown; fruit small, naked, oblong.

Type locality: Mexico.

Distribution: Guanajuato, Mexico.

We have followed Schumann in uniting *Mammillaria petterssonii* and *M. heeseana* but have selected the older name.

Dr. Rose collected this plant in Guanajuato in 1889 (No. 4846) and Dr. Safford obtained it there a few years later.

Mammillaria heeseana brevispina and *M. heeseana longispina* are two varieties listed by Schelle.

Illustrations: Ann. Rep. Smiths. Inst. 1908: pl. 7; Schelle, Handb. Kakteenk. 265. f. 188, as *Mammillaria heeseana*; Blanc, Cacti 70. f. 1350, as *M. krameri* (this is the same figure as that used by Schelle as *M. heeseana*); Cact. Journ. 1: pl. for March; Blanc, Cacti 73. No. 1460; Deutsche Garten-Zeitung 1886: 186. f. 45, as *M. petterssonii*.*

Figure 90 is a reproduction of the first illustration cited above.

37. *Neomammillaria eichlamii* (Quehl).

Mammillaria eichlamii Quehl, Monatsschr. Kakteenk. 18: 6. 1898.

Solitary or growing in large clumps of 25 or more, but loosely held together; plant-body cylindric, 6 to 15 cm. long; tubercles yellowish green, very milky, only slightly angled; axils filled with dense yellow (sometimes whitish) wool and longer white bristles; radial spines 7 or 8, ascending, whitish with brown tips; central spines usually 1, rarely 2, stouter, darker colored than the radials; spine-areoles when young filled with short yellow wool, in age glabrate; flower-buds covered with long wool; outer perianth-segments narrow, acuminate, with a dark red stripe down the center, otherwise cream-colored, slightly ciliate; inner perianth-segments narrowly lanceolate, acuminate, entire, cream-colored to light lemon-yellow; style longer than the stamens, pale; stigma-lobes linear, 4 to 6, yellow, obtuse.

Type locality: Guatemala.

Distribution: Guatemala and Honduras.

This plant differs from the other Guatemalan species in the yellow wool in the axils of the tubercles and in the areoles.

Our first knowledge of this species came from a photograph and living and herbarium material collected by Dr. William R. Maxon in Guatemala in 1905. In 1908 Quehl described it as new from specimens sent by F. Eichlam; the plant since then has been common in cultivation. It flowered first in Washington, December '909.

* This name appears as *M. petersonii* in Blanc and Schumann.

The plant is named for Federico Eichlam (1862-1911), an enthusiastic cactus collector who made very valuable discoveries in Guatemala. He published a cactus list in 1911 (Kakteen-Verzeichnis Abgeschlossen Ende 1910).

Illustrations: Monatsschr. Kakteenk. 19: 7; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 14, as *Mammillaria eichlamii*.

Figure 91 is from a photograph of a plant collected in Guatemala by F. Eichlam in 1908.

38. *Neomammillaria karwinskiana* (Martius).

Mammillaria karwinskiana Martius, Nov. Act. Nat. Cur. 16: 335. 1832.

(?) *Mammillaria fischeri* Pfeiffer, Allg. Gartenz. 4: 257. 1836.

Mammillaria centripina Pfeiffer, Allg. Gartenz. 4: 258. 1836.

Mammillaria karwinskiana flavescens Zuccarini in Pfeiffer, Enum. Cact. 19. 1837.

(?) *Mammillaria virens* Scheidweiler, Allg. Gartenz. 9: 43. 1841.

Mammillaria karwinskiana virens Salm-Dyck, Cact. Hort. Dyck. 1844. 10. 1845.

Mammillaria karwinskiana centripina Salm-Dyck, Cact. Hort. Dyck. 1844. 10. 1845.

Cactus centripinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus fischeri Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus karwinskianus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus virens Kuntze, Rev. Gen. Pl. 1: 261. 1891.



FIG. 91.—*Neomammillaria eichlamii*.




FIG. 92.—*Neomammillaria karwinskiana*.

Globose to cylindric, somewhat flattened above; tubercles terete, yielding milk when pricked; axils very woolly and with long conspicuous white or brown-tipped bristles, much longer than the tubercles; spines 4, 5, or 6, all radial, sometimes one nearer the center than the others, nearly equal, short, brown or blackish at the tips or throughout; flowers nearly 2 cm. long, the scales and outer perianth-segments narrow, reddish except at the margins, ciliate; inner perianth-segments broader, cream-colored, not ciliate, mucronate-tipped; stamens cream-colored, much shorter than the inner perianth-segments; style a little longer than the stamens; stigma-lobes 5, cream-colored; fruit 15 mm. long, red; seeds brown.

Type locality: Mexico.

Distribution: Oaxaca, Mexico.

This species is near *Neomammillaria mystax* but the spines are usually radial, short, and nearly equal. Specimens sent to Washington in 1918 had some of the lowermost spines much elongated and curved backward, sometimes 2.5 cm. long.

The plant flowers readily in cultivation. Professor C. Conzatti has repeatedly sent it to us from Oaxaca.

Illustrations: Nov. Act. Nat. Cur. **16**: pl. 22; Ann. Rep. Smiths. Inst. **1908**: pl. 14, f. 4, as *Mammillaria karwinskiana*.

Plate XI, figure 2, shows a plant collected by Dr. Rose in Oaxaca in 1906, which flowered in Washington, April 16, 1907; plate IX, figure 2, shows a plant collected by B. P. Reko also in Oaxaca, which fruited in the New York Botanical Garden in 1918. Figure 92 is from a photograph of a plant collected by Dr. Rose in Oaxaca in 1906.

Related to this species is the following:

MAMMILLARIA KNIPPELIANA Quehl, Monatsschr. Kakteenk. **17**: 59. 1907.

Stem solitary, about 7 cm. high by 6 cm. in diameter, slightly depressed at apex; tubercles when young pyramidal, 4-sided, 8 mm. long, their axils setose; areoles circular, at first white-woolly, soon glabrate; spines usually 6, up to 6 cm. long, whitish with blood-red or brown tips, sometimes accompanied with smaller spines; flowers and native country unknown.




FIG. 93.—*Neomammillaria Standleyi*.




FIG. 94.—*Neomammillaria parkinsonii*.

39. *Neomammillaria praelii* (Mühlenpfordt).

Mammillaria praelii Mühlenpfordt, Allg. Gartenz. **14**: 372. 1846.

Mammillaria viridis praelii Salm-Dyck, Cact. Hort. Dyck. 1849. 16. 1850.

Mammillaria viridis Salm-Dyck, Cact. Hort. Dyck. 1849. 116. 1850.

Mammillaria inclinis Lemaire, Illustr. Hort. **5**: Misc. 9. 1858.

Cactus praelii Kuntze, Rev. Gen. Pl. **1**: 261. 1891.


Cactus viridis Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

Globose, light green, sunken at the apex; axils of the tubercles lanate and setose; tubercles somewhat 4-angled; spine-areoles villous; spines 4, radial, forming a cross, the uppermost and lowermost elongated; flowers and fruit unknown.

Type locality: Guatemala.

Distribution: Guatemala.

We do not know this species but we are following previous authors in our classification of it. When flowers and fruit become known this may be subject to modification. Until recently it and *Neomammillaria woburnensis* were the only species of this genus known from Guatemala; neither was known in cultivation. Through the efforts of Dr. William R. Maxon, Mr. F. Eichlam, Professor Kellermann, and others, much material has been collected, new species discovered, and *N. woburnensis* rediscovered, but not *N. praelii*.



M. E. Eaton del. 1 to 4

E. I. Schutt del. 2

D. G. Passmore del. 5

A. Hoen & Co. Baltimore

1. Flowering plant of *Neomammillaria magnimamma*.
2. Flowering plant of *Neomammillaria karwinskiana*.

3. Flowering plant of *Neomammillaria gigantea*.
4. Flowering plant of *Neobesseyia missouriensis*.

Schumann described the plant in some detail, but apparently confused it with another species, possibly *Mammillaria karwinskiana*, inasmuch as he reported it from Oaxaca as well as from Guatemala. He referred here as a synonym *M. viridis* Salm-Dyck (Cact. Hort. Dyck. 1849. 16. 1850), which may be the Mexican element.

40. *Neomammillaria standleyi* sp. nov.

Plants usually solitary, nearly globular, often 10 cm. in diameter, pale green, densely covered with spines; axils of tubercles containing white bristles, the flowering and fruiting ones filled with dense white wool; radial spines about 16, slightly spreading, white except the dark tips; central spines 4, longer and stouter than the radials, porrect, reddish brown; flowers rather small, about 12 mm. long, purplish; inner perianth-segments oblong, entire; filaments pale; stigma-lobes green; fruit scarlet, 12 to 16 mm. long; seeds brownish.

Collected by Rose, Standley, and Russell on rocks in the Sierra de Alamos, Sonora, Mexico, March 14, 1910 (No. 12849).

It is common in dry stony places above Alamos, where both living and herbarium specimens were obtained, and is an attractive plant flowering freely in cultivation.

The plant is named for Paul C. Standley of the U. S. National Museum.

Figure 93 is from a photograph of the type specimen which flowered in Washington.




FIG. 95.—*Neomammillaria evermanniana*.

41. *Neomammillaria evermanniana* sp. nov.

Globose to elongate-turbinate, 5 to 7 cm. in diameter, lactiferous; tubercles closely set, tere, nearly hidden under the numerous slender spines; axils of tubercles at first very woolly and setose; spines white except at tip and there brown; radial spines 12 to 15; central spines 3, erect or nearly so; fruit red, about 1 cm. long; seeds brown.

Collected by Ivan M. Johnston on Cerralbo Island, Gulf of California, 1921 (No. 4058). Mr. Johnston writes of it as follows:

"I found it growing wedged in narrow dirt-filled cracks on the canyon side of the island. It is quite common on this island, usually growing singly, but one cespitose mass with 19 unequal heads was observed."

The species is named for Dr. Barton W. Evermann, Director of the Museum of the California Academy of Sciences, who organized the scientific expedition to the Gulf of California in 1921, which obtained this as well as many other new and rare plants.

Related to this species, but perhaps distinct from it, is Johnston's No. 3121 from Nolasco Island, Gulf of California. It has fewer spines (about 10 radials and 1 or 2 centrals).

Figure 95 is from a photograph of plants from the type collection.

42. *Neomammillaria parkinsonii* (Ehrenberg).

Mammillaria parkinsonii Ehrenberg, *Linnaea* 14: 375. 1840.
Cactus parkinsonii Kuntze, *Rev. Gen. Pl.* 1: 261. 1891.

Cespitose, somewhat depressed to cylindric, 15 cm. high, 7.5 cm. in diameter, globose, glaucous, green; axils of tubercles lanate and setose; tubercles milky, short, conic; radial spines numerous (20 or more), setaceous, short, white; central spines 2 or sometimes 4 or 5, brownish at tip; flowers surrounded by a mass of wool, small, yellowish; inner perianth-segments apiculate; stigma-lobes elongated; fruit clavate, scarlet, 1 cm. long; seeds brown.

Type locality: At San Onofre in the Mineral del Doctor, Mexico.

Distribution: Central Mexico.

We have a photograph, identified as this plant, sent us by L. Quehl in 1921, and also specimens which are like this photograph, collected by Dr. Rose near Higuierillas, Queretaro, Mexico, in 1905 (No. 9798).

The plant was named for John Parkinson, at one time British Consul-General in Mexico, who died in Paris, April 3, 1847.

Mammillaria parkinsonii rubra (Förster, *Handb. Cact.* 196. 1846) is only a name.

Mammillaria parkinsonii waltonii we do not know, although it is frequently referred to in cactus literature. Haage and Schmidt offer it for sale in their catalogue (1920) under the name of *M. waltonii* Quehl.

Illustrations: *Cact. Journ.* 1: pl. for March, as *Mammillaria waltonii*; *Gartenwelt* 14: 232; Möllers *Deutsche Gärt. Zeit.* 25: 475. f. 8, No. 15; Rother, *Praktischer Leitfaden Kakteen* 39, as *M. parkinsonii*.

Figure 94 is from a photograph sent by L. Quehl.

43. *Neomammillaria geminispina* (Haworth).

Mammillaria geminispina Haworth in Gillies, *Phil. Mag.* 63: 42. 1824.
Mammillaria bicolor Lehmann, *Samen. Hamb. Gart.* 7. 1830.
Mammillaria nivea Wendland in Pfeiffer, *Enum. Cact.* 27. 1837.
Mammillaria daedalea Scheidweiler, *Hort. Belge* 4: 16. 1837.
Mammillaria toaldoae Lehmann, *Linnaea* 12: 13. 1838.
Mammillaria eburnea Miquel, *Linnaea* 12: 14. 1838.
Mammillaria nivea daedalea Lemaire, *Cact. Gen. Nov. Sp.* 101. 1839.
Mammillaria nobilis Pfeiffer, *Allg. Gartenz.* 8: 282. 1840.
Mammillaria bicolor longispina Salm-Dyck, *Cact. fort. Dyck.* 1844. 6. 1845.
Mammillaria bicolor cristata Salm-Dyck, *Cact. Hort. Dyck.* 1844. 6. 1845.
Mammillaria bicolor nobilis Förster, *Handb. Cact.* 198. 1846.
Cactus geminispinus Kuntze, *Rev. Gen. Pl.* 1: 260. 1891.
Cactus niveus Kuntze, *Rev. Gen. Pl.* 1: 261. 1891.
Cactus nobilis Kuntze, *Rev. Gen. Pl.* 1: 261. 1891. Not Lamarck, 1783.
Mammillaria bicolor nivea Schumann, *Gesamtb. Kakteen* 569. 1898.

Cespitose, or single in cultivation, cylindric, somewhat glaucous; axils woolly; tubercles terete, conic; radial spines numerous (16 to 20), very short, setaceous, white; central spines 2 to 4, stouter and longer than the radials, about 25 mm. long, black-tipped; flowers dark red; inner perianth-segments oblong, obtuse, serrate.

Type locality: Mexico.

Distribution: North-central Mexico.

Mammillaria daedalea, which is referred here by Schumann, is based on an abnormal specimen which has elongated, contorted stems and looks very unlike the typical plant. Scheidweiler illustrated his species.

Mammillaria nivea cristata Salm-Dyck (Walpers, Repert. Bot. 2: 270. 1843) is only a name. *M. nivea wendlei* Pfeiffer (Labouret, Monogr. Cact. 57. 1853) was given as a synonym of *M. bicolor*.

To this relationship we would refer the plant which has long been known in collections under the name of *Mammillaria potosina** and *M. potosina* var. *longispina*. It resembles *M. celsiana* in the spines, but the tubercles are milky and the stem is more elongated. We have seen the following illustration: Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 9, as *M. potosina*.

De Candolle (Prodr. 3: 459. 1828) referred here *Cactus columnaris* Mociño and Sessé.




FIG. 96.—*Neomammillaria collinsii*.




FIG. 97.—*Neomammillaria geminispina*.

Illustrations: Wiener Ill. Gart. Zeit. 11: pl. 3, in part, as *Mammillaria nobilis*; Hort. Belge 4 pl. 1, as *M. daedalea*; Möllers Deutsche Gärt. Zeit. 25: 75. f. 8, No. 4, as *M. bicolor nobilis*; Cact. Journ. 1: pl. for March, as *M. nivea cristata*; Cact. Journ. 1: pl. for March, as *M. nivea longispina*; Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 3; De Laet, Cat. Gén. f. 50, No. 8; Wiener Ill. Gart. Zeit. 29: f. 22, No. 8; Knippel, Kakteen pl. 19, as *M. bicolor*.

Plate v, figure 3, shows a flowering plant sent by Carl Ackerman which flowered in the New York Botanical Garden, October 9, 1920; plate VIII, figure 5, shows a plant which flowered in the New York Botanical Garden, November 11, 1911. Figure 97 is from a photograph by Ernest Branton showing a plant grown in southern California.

44. *Neomammillaria pyrrhocephala* (Scheidweiler).

Mammillaria pyrrhocephala Scheidweiler, Allg. Gartenz. 9: 42. 1841.

Mammillaria mallettiana Cels, Portef. Hort. 2: 222.

Mammillaria senkei† Förster, Handb. Cact. 227. 1846.

Mammillaria pyrrhocephala donkelaeri Salm-Dyck, Cact. Hort. Dyck. 1849. 17, 121. 1850.

Cactus pyrrhocephalus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cylindric; axils lanate and setose; tubercles angled, green or subglaucous; areoles bearing yellowish wool; spines all black when young, when old becoming gray below; radial spines 6, spreading, the upper ones a little longer; central spines single, erect; flowers red.

Type locality: Real del Monte, Mexico.

Distribution: Hidalgo and, perhaps, Oaxaca.

* This name is sometimes credited to Rebut (Möllers Deutsche Gärt. Zeit. 25: 475. 1910) but if he published it we are unaware of it.

† This was originally written *M. senkii*, although the plant was named for F. Senke of Leipzig.

We have followed Schumann and others who refer this species also to Oaxaca but the plants from that state may represent more than one species. In fact, the plant figured in *Blühende Kakteen* we have described as new (see No. 50), while the one illustrated by Mr. H. H. Thompson is like others sent by Dr. Reko and Professor Conzatti, which we have referred here.

Illustration: Thompson, U. S. Dept. Agr. Bur. Pl. Ind. Bull. **262**: pl. 2, f. 2, as *Mammillaria pyrrhocephala*.

Figure 100 is from a photograph of the plant sent to Washington by Dr. Reko from Oaxaca in 1919.

45. *Neomammillaria woburnensis* (Scheer).

*Mammillaria woburnensis** Scheer, Lond. Journ. Bot. **4**: 136. 1845.

Cactus woburnensis Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

Mammillaria chapinensis Eichlam and Quehl, Monatsschr. Kakteenk. **19**: 1. 909.




FIG. 98.—*Neomammillaria woburnensis*.




FIG. 99.—*Neomammillaria chionocephala*.

Growing in clumps, giving off new plants from all parts of the body, globose to cylindric, dull green, milky; tubercles angled, setose and woolly in their axils; radial spines 5 to 9, yellowish or white; central spines 1 to 8, often long, reddish or yellow; flowers yellow, small, about 1 cm. long; fruit red, clavate, 18 to 25 mm. long; seeds minute, brown.

Type locality: Guatemala.

Distribution: Guatemala.

For a long time little was known about this plant, but a few years ago it was discovered in abundance by Wm. R. Maxon (1905) and by F. Eichlam (1908). It was given a new name, *Mammillaria chapinensis*, under which it is to be found in most collections.

The plant was described by Frederick Scheer from a barren specimen in the Royal Botanical Garden at Kew, sent from Guatemala. It was named for Woburn Abbey, where there was once a very large collection of cacti under the care of James Forbes.

Illustrations: Monatsschr. Kakteenk. **24**: 87; Succulenta **4**: 40; Möllers Deutsche Gärt. Zeit. **25**: 475. f. 8, No. 12, as *Mammillaria chapinensis*.

Figure 98 is from a photograph of a plant sent to Washington by F. Eichlam in 1908.

* This name was originally printed by Scheer as *Mamillaria woburnensis*.

46. *Neomammillaria collinsii* sp. nov.

Plants becoming large clumps, the individuals globose, 4 cm. in diameter; tubercles terete, milky, green, but becoming bronzed or even a deep purple; axils of tubercles both lanate and setose; radial spines usually 7, pale yellowish below, with dark brown or blackish tips, subequal, 5 to 7 mm. long; central spine 1, similar to or a little longer and usually darker than the radials; flowers 12 to 15 mm. long; outer perianth-segments reddish with a yellowish margin, ciliate; inner perianth-segments lighter, entire, acuminate; fruit clavate, 1.5 to 2 cm. long, deep red; seeds brownish.

Collected by G. N. Collins at San Gerónimo, near Tehuantepec, Mexico, December 1906, and flowered in Washington, July and August 1909, type, and near the same locality by A. Groeschner, February 1923.

Figures 96 and 103 are from photographs showing the type plant in flower and fruit.

47. *Neomammillaria chinocephala* (J. A. Purpus).

Mammillaria chinocephala J. A. Purpus, Monatsschr. Kakteenk. 16: 41. 1906.

Plant-body globose, sometimes 8 cm. in diameter, almost hidden by the white spines; tubercles very milky; axils of tubercles densely filled with white wool and numerous hair-like bristles; tubercles low; radial spines 35 to 40, somewhat pectinate, spreading; central spines 2 to 7, more or less divergent, much stouter than the radials, rigid, white with brownish tips; flowers 1 cm. long, rose-red; fruit clavate, red; seeds small, brown.

Type locality: Sierra de Parras, Coahuila, Mexico.

Distribution: Highlands of central Mexico.

This species is common in collections, both living and dried, and it is surprising that it remained so long undescribed. It was distributed by Pringle in 1890 as *Mammillaria acanthophlegma*. It resembles very much a large plant of *Mammillaria elegans*, but the tubercles are milky and bear setae in their axils.

Illustrations: Monatsschr. Kakteenk. 16: 3; 20: 46, as *Mammillaria chinocephala*. Figure 99 is from a photograph of a plant collected by Dr. Purpus at Minas de San Rafael, Mexico, in 1910.

48. *Neomammillaria tenampensis* sp. nov.

Globose, light green, 5 to 6 cm. in diameter; tubercles 6 to 7 mm. long, 4-sided, milky, pointed; axils of upper tubercles naked, but those producing flowers filled with yellow wool and numerous yellow bristles, while in the older axils the wool disappears and the bristles become white; spines 4 to 6, brownish with dark tips, ascending, surrounded at base by 8 to 10 small white bristles; wool in young spine-areoles yellowish; outermost perianth-segments small, brownish, the outer ones lanceolate, acuminate, similar to the inner ones, all ciliate; inner perianth-segments reddish purple, 8 to 10 mm. long, lanceolate, apiculate, denticulate; stamens much shorter than the perianth-segments; filaments pale below, purplish above; style reddish; stigma-lobes 4 or 5.

Collected by C. A. Purpus in the Barranca de Tenampa, Mexico, in 1909 and flowered in Washington in November 1910.

Figure 102 is from a photograph of the type specimen.

49. *Neomammillaria polygona* (Salm-Dyck).

Mammillaria polygona Salm-Dyck, Cact. Hort. Dyck, 1849. 120. 1850.

Cactus polygonus Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Lamarck, 1783.

Subclavate, 10 cm. high, simple; axils of tubercles lanate and setose; tubercles 4-angled; radial spines about 8, 2 or 3 upper ones minute, the lateral ones and the lowermost one longer; central spines 2, stout, brownish at tip, often long and recurved; flowers pale rose-colored; stigma-lobes 5 or 6, linear.

Type locality: Not cited.

Distribution: Mexico, according to Labouret.

Schumann lists this species among those unknown to him. Rümpler refers it to *Mammillaria subpolyedra*, but it must be related more nearly to *M. polyedra*, with which it was compared by Salm-Dyck. We know it only from descriptions.

Mammillaria polyedra spinosior Salm-Dyck (Cact. Hort. Dyck. 1849. 17. 1850) is usually referred here, but was never described.

Related to this species is the following:

MAMMILLARIA ECHINOPS Scheidweiler, Hort. Belge 5: 5. 1838.

Simple, globose or a little broader than high, 8 cm. in diameter, lactiferous; tubercles ovoid, light green, somewhat 4-angled, lanate and setose in their axils; radial spines 12 or 13, the upper three much shorter, setose, the others about equal; central spines 4, stout when young, white, with rosy brown tips, these black in age; flowers not known; fruit red, clavate, 8 mm. long.

Type locality: Mexico.

We have not been able to associate this description or illustration with any species which we know. The author believed that it was related to *Mammillaria polyedra*. The setae in the axils of the tubercles suggest this relationship, but we believe that it is very distinct from that species.

The original description seems to have been unknown to the compilers of the Index Kewensis and to Schumann, for they refer the name to Förster's Handbuch, where it is used as a synonym of another species. Förster, followed by the Index Kewensis, refers it as a synonym of *Mammillaria oothele*, which is a very different plant if we can judge from the description.

Illustration: Hort. Belge 5: pl. 5.

50. *Neomammillaria confusa* sp. nov.

At first solitary, becoming cespitose, globose to short-cylindric, deep green; axils densely white-woolly and setose; tubercles short, a little flattened, 4-angled, pointed; spines 4 to 6, all radial, ascending, at first yellowish with brown tips, in age white below, 2 to 3 mm. long; flowers yellow, small, about 8 mm. long, opening for 2 or 3 successive days; outer perianth-segments ovate, ciliate, with a black tip; inner perianth-segments spreading, acute; filaments and style yellowish white; stigma-lobes 6, greenish yellow.

In 1912 Dr. Rose obtained a plant from W. Mundt near Berlin which flowered in the New York Botanical Garden in April 1914 and in 1918 and which we have designated as the type. It is not known in the wild state, but is doubtless from Mexico.

This is the plant which Schumann described and figured as *Mammillaria pyrrhocephala*, but it does not accord with the original description.

Illustration: Blühende Kakteen 1: pl. 20, as *Mammillaria pyrrhocephala*.

Plate v, figure 2, shows the type plant.

51. *Neomammillaria villifera* (Otto).

Mammillaria villifera Otto in Pfeiffer, Enum. Cact. 18. 1837.

Cactus villifer Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria carnea villifera Gürke, Blühende Kakteen 1: under pl. 60. 1905.

Subglobose, proliferous; axils lanate and setose; tubercles angled; areoles at first lanate, in age naked; spines 4, rigid, straight, the lowest one longer (8 mm. long), at first purplish, in age black; flowers pale rose-colored; inner perianth-segments 14, acute; stigma-lobes 4 or 5.

Type locality: Mexico.

Distribution: Mexico, but range not known.

The species is often referred to *Mammillaria carnea*, but the axils are setose.

52. *Neomammillaria polyedra* (Martius).

Mammillaria polyedra Martius, Nov. Act. Nat. Cur. 16: 326. 1832.

Mammillaria polytricha Salm-Dyck, Allg. Gartenz. 10: 289. 1842.

Mammillaria polytricha hexacantha Salm-Dyck, Allg. Gartenz. 10: 289. 1842.

Mammillaria polytricha tetracantha Salm-Dyck, Allg. Gartenz. 10: 290. 1842.

Mammillaria polyedra laevior Salm-Dyck in Labouret, Monogr. Cact. 105. 1853.

Mammillaria polyedra scleracantha Labouret, Monogr. Cact. 105. 1853.

Cactus polyedrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus polytrichus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Solitary, cylindric or somewhat thicker above; axils of tubercles setose; tubercles 12 mm. long, flattened dorsally, angled, pointed; spines 4, ascending, short, grayish with purplish tips; flowers inconspicuous, reddish; inner perianth-segments short-acuminate; anthers white; style white, longer than the stamens; stigma-lobes 8, greenish; fruit unknown.

Type locality: Near Oaxaca, Mexico.

Distribution: Southern Mexico.

This species was collected by Baron Karwinsky near Oaxaca City, about 1832. It has been reported over a large area of central Mexico, but is doubtless much more restricted in range. One small specimen from near the type locality was sent to Washington in 1909.




FIG. 100.—*Neomammillaria pyrrocephala*.




FIG. 101.—*Neomammillaria polyedra*.

Mammillaria anisacantha Hortus first appeared as a synonym of *M. polyedra anisacantha* Salm-Dyck (Cact. Hort. Dyck. 1844. 11. 1845) and then as a synonym of *M. polyedra laevior* Salm-Dyck (Cact. Hort. Dyck. 1849. 17. 1850); neither of the varieties was here described, but the latter was briefly characterized by Labouret. *Mammillaria scleracantha* is cited from Monville's Catalogue of 1846 but we have not seen this publication; it does occur as a synonym of *M. polyedra scleracantha* in Labouret's Monograph, p. 105.

Illustrations: Martius, Nov. Act. Nat. Cur. 16: pl. 18; Blühende Kakteen 2: pl. 112; Schelle, Handb. Kakteenk. 271. f. 194, as *Mammillaria polyedra*.

Plate XII, figure 5, shows the plant sent from the Berlin Botanical Garden in 1914 which flowered in the New York Botanical Garden on April 1, 1918. Figure mi shows the type plant, being a reproduction of the first illustration cited above.

53. *Neomammillaria konzattii* sp. nov.

Short-cylindric, 8 cm. high, sometimes branched at apex, dark green, very milky; axils of young tubercles bearing abundant white wool and conspicuous white bristles; tubercles short, 4 to 5 mm. long, somewhat angled; young spine-areoles woolly; spines 4 or 5, all radial, somewhat spreading, brownish, the tips usually darker than the bases; flowers opening in bright sunlight, white, campanulate, sometimes tinged with red, about 2 cm. long, the segments somewhat spreading, narrowly oblong, the outer ones serrulate, apiculate; style pale green; stigma-lobes, white.

Collected by C. Konzatti on Cerro San Felipe, Oaxaca, in 1907 and flowered in 1913 (type); collected again in 1921 (No. 4140) and flowered in April 1922.

Figure 104 is from a photograph of the plant collected by C. Konzatti in 1921.

54. *Neomammillaria napina* (Purpus).*Mammillaria napina* Purpus, Monatsschr. Kakteenk. 22: 161. 1912.

Roots thick, but when in a cluster of 3 or 4 somewhat spindle-shaped; plants globose, 4 to 6 cm. in diameter; tubercles low, terete in section, not at all milky; spines all radial, 10 to 12, pectinate, white or yellowish, spreading and interlacing; flowers unknown.

Type locality: Mountains west of Tehuacán, Mexico.

Distribution: Southern Mexico.

The plant was collected by C. A. Purpus in 1911. In 1901 Dr. Rose collected near Tehuacán three small plants which we now believe are to be referred here; these differ from the type plant chiefly in having usually one porrect central spine 5 to 8 mm. long. Some of the spine-clusters have no central spines and then they look very much like those of *Neomammillaria napina*. Dr. Rose's plants were globose when collected but now are cylindric, and after 20 years are less than 6 cm. high; they have never flowered.

Illustration: Monatsschr. Kakteenk. 23: 123, as *Mammillaria napina*.




FIG. 102.—*Neomammillaria tenampensis*.




FIG. 103.—*Neomammillaria collinsii*.

55. *Neomammillaria lanata* sp. nov.

Small, short-cylindric; tubercles short, 2 to 4 mm. long; spine-areoles short-elliptic; spines 12 to 14, all radial, widely spreading, white except the brown bases; flowering areoles very woolly, the young flowers surrounded by a mass of long white hairs; flowers very small, 6 to 7 mm. long, red; inner perianth-segments about 15, oblong, obtuse or acutish, spreading above; stigma-lobes 3, short, obtuse.

Collected by C. A. Purpus near Rio de Santa Luisa, Mexico, in 1907 and since grown in Washington.

Figure 105 is from a photograph of the type specimen.

56. *Neomammillaria kewensis* (Salm-Dyck).*Mammillaria kewensis* Salm-Dyck, Cact. Hort. Dyck. 1849. 112. 1850.

Globose to cylindric, 3 to 4 cm. in diameter; tubercles short, terete, when young short-woolly in the axils and at the areoles; spines 5 or 6, all radial, 4 or 5 mm. long, brown with dark tips; axils of tubercles bearing crisp hairs; flowers about 15 mm. long, reddish purple; perianth-segments lanceolate, acute; stigma-lobes 5, reddish.

Type locality: Not cited.

Distribution: Doubtless Mexico.

We have had a living plant from Haage and Schmidt and one from Quehl which we have used in our description.

Salm-Dyck (Cact. Hort. Dyck. 1849. 15. 1850) mentions *Mammillaria kewensis* var. *albispina* and also *M. spectabilis* Hortus as synonyms.

This plant was named for the Royal Botanic Gardens, Kew.

Illustration: Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 3, as *Mammillaria kewensis*.

Figure 106 is reproduced from a photograph sent us by L. Quehl in 1921.

57. *Neomammillaria subpolyedra* (Salm-Dyck).

Mammillaria subpolyedra Salm-Dyck, Hort. Dyck. 343. 1834.

Cactus subpolyedrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Solitary, subcylindric, 10 cm. high, 6 cm. in diameter; tubercles pointed, strongly angled; axils and spine-areoles white-woolly; spines 4, at first blackish purple, becoming paler but the tips remaining purplish, the lowest one the largest; flowers 2.5 cm. broad; perianth-segments obtuse, erose, with a darker midrib; fruit red, 2.5 cm. long, pyriform, 12 mm. in diameter at apex.




FIG. 104.—*Neomammillaria conzattii*.




FIG. 105.—*Neomammillaria lanata*.

Type locality: Not cited.

Distribution: According to Rümpler, Zimapan and Ixmiquilpan, Mexico.

Some of the illustrations here cited do not correspond very well with the original description. This species is listed by Schumann with those unknown to him, and it is known to us only from descriptions and illustrations.

Mammillaria polygona Zuccarini (Pfeiffer, Enum. Cact. 17. 1837) is referred here but it was never described. Salm-Dyck afterwards used the name for a very different plant.

Mammillaria jalappensis and *M. anisacantha* are referred by Pfeiffer (Enum. Cact. 17. 1837) as synonyms of *M. subpolyedra*.

Illustrations: (?) Förster, Handb. Cact. ed. 2. 357. f. 37; (?) Dict. Gard. Nicholson 4: 565. f. 39; Suppl. 518. f. 557; Watson, Cact. Cult. 176. f. 71; ed. 3. f. 48, as *Mammillaria subpolyedra*.

Figure 107 is reproduced from the illustration used in Nicholson's Dictionary.

58. *Neomammillaria galeottii* (Scheidweiler).

Mammillaria galeottii Scheidweiler, Hort. Belge 4: 93. 1837.

Mammillaria obconella galeottii Scheidweiler, Hort. Belge 4: 93. 1837.

Mammillaria dolichocentra galeottii Salm-Dyck in Förster, Handb. Cact. 213. 1846.

Mammillaria dolichocentra phaeacantha Labouret, Monogr. Cact. 50. 1853.

Solitary or cespitose, globose; tubercles pointed; spines 4, elongated, the upper ones erect and connivent over apex of plant, on older tubercles weak and spreading, 2.5 cm. long, pale rose to crimson.

Type locality: Mexico.

Distribution: Mexico.

We have not seen this plant, but we have examined the illustration which accompanies the original description. L. Quehl has had it in cultivation, and sent us a photograph.

This must be a very distinct species and not at all closely related to *Mammillaria dolichocentra*, to which Schumann referred it as a variety, crediting himself as the authority; the name, however, had been used by Förster in 1846. The illustrations in Förster's Handbuch der Cacteenkunde and in Nicholson's Dictionary cited below probably are not to be referred here and they certainly should not be referred to *Mammillaria dolichocentra*.

Mammillaria obscura galeottii Salm-Dyck (Förster, Handb. Cact. 213. 1846) is mentioned as a synonym of this species, but so far as we can learn it was never described.

Illustrations: Hort. Belge 4: pl. 6; Rother, Praktischer Leitfaden Kakteen 37, as *Mammillaria galeottii*; Förster, Handb. Cact. ed. 2. 323. f. 32; Dict. Gard. Nicholson 2: 321. f. 508, as *Mammillaria dolichocentra*.




FIG. 106.—*Neomammillaria kewensis*.




FIG. 107.—*Neomammillaria subpolyedra*.

59. *Neomammillaria tetracantha* (Salm-Dyck).

Mammillaria tetracantha Salm-Dyck in Pfeiffer, Enum. Cact. 18. 1837.

Mammillaria obconella Scheidweiler, Hort. Belge 4: 93. 1837.

Mammillaria dolichocentra Lemaire, Cact. Aliq. Nov. 3. 1838.

Mammillaria dolichocentra staminea Labouret, Monogr. Cact. 50. 1853.

Cactus obconella Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Cactus dolichocentrus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus tetracanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Mammillaria rigidispina Hildmann, Monatsschr. Kakteenk. 3: 112. 1893.

Mammillaria dolichocentra brevispina Runge, Monatsschr. Kakteenk. 3: 112. 1893.

Nearly globular, 6 to 8 cm. in diameter; axils of tubercles with scanty persistent wool; tubercles 8 to 10 mm. long, obscurely 4-angled; areoles small, at first lanate, somewhat 4-angled; spines, all radial, slender, the 3 lower equal, the upper one incurved, longer, 25 mm. long, when young all yellowish white, in age grayish yellow or brown; flowers numerous from towards top of plant, small, pinkish to rose-colored; inner perianth-segments narrowly lanceolate, acuminate.

Type locality: Mexico, but no definite locality cited.

Distribution: Mexico, but range unknown.

Schumann refers here *Mammillaria longispina* Reichenbach (Suppl. Terscheck Cact. Verz.; see also Walpers, Repert. Bot. 2: 301. 1843) and *M. obconella* Scheidweiler (Hort. Belge 4: 93. f. 6. 1837), but we are uncertain as to their relationship. To the former Walpers refers as a synonym *M. galeottii* Otto.

Mammillaria dolichacantha Lemaire (Förster, Handb. Cact. 213. 1846) and *M. dolichocentra picta* (Salm-Dyck, Cact. Hort. Dyck. 1844. 9. 1845) were never described.

Illustrations: Curtis's Bot. Mag. 70: pl. 4060, as *Mammillaria tetracantha*; Cassell's Dict. Gard. 2: 48; Karsten, Deutsche Fl. 887. f. 501, No. 2; ed. 2. 2: 456. f. 605, No. 2; Schelle, Handb. Kakteenk. 260. f. 182; Watson, Cact. Cult. 155. f. 58; ed. 3. f. 36; Lemaire, Icon. Cact. pl. 5; Schumann, Gesamtb. Kakteen 558. f. 91; Förster, Handb. Cact. ed. 2. 322. f. 31; Gartenwelt 9: 265; Lemaire, Cactées 37. f. 3, as *Mammillaria dolichocentra*; Rev. Hort. 1861: 270. f. 72, as *Mammillaria*; Thomas, Zimmerkultur Kakteen 54; Monatsschr. Kakteenk. 3: 11 3, as *Mammillaria rigidispina*.

Figure 108 is a reproduction of the first illustration cited above.

60. *Neomammillaria elegans* (De Candolle).

- Mammillaria geminispina* * De Candolle, Mém. Mus. Hist. Nat. Paris 17: 30. 1828. Not Haworth, 1824.
Mammillaria elegans De Candolle, Mém. Mus. Hist. Nat. Paris 17: 111. 1828.
Mammillaria elegans minor De Candolle, Mém. Mus. Hist. Nat. Paris 17: 111. 1828.
 ? *Mammillaria elegans globosa* De Candolle, Mém. Mus. Hist. Nat. Paris 17: 111. 1828.
Mammillaria acanthophlegma Lehmann, Del. Sem. Hamb. 1832.
Mammillaria supertexta Martius in Pfeiffer, Enum. Cact. 25. 1837.
Mammillaria dyckiana Zuccarini in Pfeiffer, Enum. Cact. 26. 1837.
Mammillaria elegans micrantha Lemaire, Cact. Gen. Nov. Sp. 100. 1839.
Mammillaria geminispina tetracantha Lemaire, Cact. Gen. Nov. Sp. 100. 1839.
Mammillaria klugii Ehrenberg, Bot. Zeit. 2: 834. 1844.
Mammillaria meisneri Ehrenberg, Bot. Zeit. 2: 834. 1844.
Mammillaria kunthii Ehrenberg, Bot. Zeit. 2: 835. 1844.
Mammillaria splendens Ehrenberg, Allg. Gartenz. 57: 242. 1849.
Mammillaria acanthophlegma decandollii Salm-Dyck, Cact. Hort. Dyck. 1849. 9. 1850.
Mammillaria elegans klugii Salm-Dyck, Cact. Hort. Dyck. 5849. 9. 1850.
Mammillaria acanthophlegma meisneri Salm-Dyck, Cact. Hort. Dyck. 1849. 9. 1850.
Mammillaria supertexta tetracantha Salm-Dyck in Labouret, Monogr. Cact. 61. 1853.
Mammillaria acanthophlegma elegans Monville in Labouret, Monogr. Cact. 63. 1853.
Mammillaria acanthophlegma monacantha Monville in Labouret, Monogr. Cact. 63. 1853.
Mammillaria acanthophlegma leucocephala Monville in Labouret, Monogr. Cact. 63. 1853.
Mammillaria acanthophlegma abducta Monville in Labouret, Monogr. Cact. 64. 1853.
Cactus acanthophlegma Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus dyckianus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus elegans Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Link, 1822.
Cactus kunthii Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus klugii Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus meisneri Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus supertextus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Simple, obovate to globose, 5 cm. in diameter, somewhat umbilicate at apex; tubercles ovate, naked in their axils, not lactiferous; spine-areoles tomentose when young; radial spines stiff, bristle-like, 25 to 30, white, spreading; central spines 1 (sometimes 2 or 3), rigid.

Type locality: Mexico.

Distribution: Central Mexico.

This species was based on Thomas Coulter's No. 48 from Mexico but no definite locality was cited. The type was not preserved nor is there any illustration extant of the original. De Candolle may have had more than one species before him when he drew up his description, for he described two varieties, one of which has bristles in the axils of the tubercles, which are never found in *Neomammillaria elegans* as we have treated it here.

Plants named *Mammillaria elegans* are to be found in most collections of cacti, but the name is often applied to several closely allied species. A plant from northern Mexico, *Mammillaria chionocephala*, resembles it very much but has milky tubercles. Other species which

* Here De Candolle referred *Cactus columnaris* Mociño and Sessé (De Candolle, Prodr. 3: 459. 188), which Schumann has inadvertently taken up as *Mammillaria columnaris* Mociño and Sessé (Gesamtb. Kakteen 565. 1898).

have passed as *M. elegans* have recently been described as *Mammillaria pseudoperbella* and *M. perbella*.

Mammillaria supertexta caespitosa Monville (Salm-Dyck, Cact. Hort. Dyck. 1844. 6. 1845) is only a name; *M. supertexta compacta* Scheidweiler (Labouret, Monogr. Cact. 61. 1853) was given as a synonym of *M. supertexta tetracantha* but may not belong here.

The name *Mammillaria leucocephala* Hortus is given by Pfeiffer as a synonym of *M. acanthophlegma*. *M. recta* Miquel (Labouret, Monogr. Cact. 63. 1853) occurs only as a synonym for the same species.

Illustrations: Blühende Kakteen 3: pl. 139; Cact. Journ. 1: pl. for February; Schelle, Handb. Kakteenk. 261. f. 183; Schumann, Gesamtb. Kakteen 564. f. 92, as *Mammillaria elegans*; Mém. Mus. Hist. Nat. Paris 17: pl. 3, as *Mammillaria geminispina*; Cact. Journ. 1: pl. for February in part, as *Mammillaria supertexta*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 24, as *Mammillaria dyckiana*.




FIG. 108.—*Neomammillaria tetracantha*.



FIG. 109.—*Neomammillaria elegans*.

Figure 109 is from a photograph of the plant grown in the Huntington Collection near Los Angeles, California, as this species.

Of this relationship are the following:

MAMMILLARIA CONSPICUA J. A. Purpus, Monatsschr. Kakteenk. 22: 163. 1912.

Simple, cylindric to globose, not milky; spine-areoles small, short-elliptic, when young a little woolly, in age glabrate; radial spines 10 to 25, rigid; central spines 2, a little curved; fruit red; seeds 1 mm. long.

Type locality: Near Zapotitlán, Puebla, Mexico.

Illustration: Monatsschr. Kakteenk. 24: 37.

MAMMILLARIA MICROTHELE Mühlenpfordt, Allg. Gartenz. 16: 11. 1848.

Cactus bispinus Coulter, Contr. U. S. Nat. Herb. 3: 101. 1894.

Cespitose, many-headed; joints globose, small; tubercles when dry 6 mm. long, naked or woolly in their axils; radial spines 22 to 24, white-setiform, spreading, 2 to 4 mm. long; central spines 2, much stouter than the radials, 2 mm. long or less; flowers flesh-colored without, white within, small, only 3 to 4 mm. long when dried; fruit clavate, 10 mm. long; seeds rather large, probably black.

Type locality: Not known but supposed to be Mexico.

Distribution: Mexico.

Our description is drawn from the original, supplemented by specimens in the Engelmann Herbarium obtained from Salm-Dyck's garden in January 1857, which consist of two packets, one containing a few spine-clusters and the other several withered flowers and nearly ripe fruits; these latter are labeled "Baumann 857." Engelmann and Coulter compare this species with *Mammillaria micromeris* but we believe that it is related to *M. elegans* and its allies.

It seems to have been described from specimens of Haage of unknown origin but supposed to be from Mexico; Coulter's reference, on the statement of Budd, that it occurs within the southern border of Pecos County, Texas, is to be doubted.

Coulter renamed *Mammillaria microthele* because of an older *Cactus microthele*. Martius used the name *M. microthele* in 1829 (Hort. Reg. Monac. 127) but without description. The names *M. brongniartii* Hortus, *M. microthele brongniartii*, and *M. compacta* Hortus (not Engelmann, 1848) have been used (Salm-Dyck, Cact. Hort. Dyck. 1849. 9. 1850) but without descriptions.




FIG. 110.—*Neomammillaria pseudoperbella*.




FIG. 111.—*Neomammillaria dealbata*.

61. *Neomammillaria pseudoperbella* (Quehl).

Mammillaria pseudoperbella Quehl, Monatsschr. Kakteenk. 19: 188. 1909.

Mammillaria pseudoperbella rufispina Quehl, Monatsschr. Kakteenk. 26: 94. 1916.

Solitary, or few together, globose to short-cylindric, very spiny, depressed at apex; tubercles short-cylindric; radial spines 20 to 30, setaceous, white, short; central spines 2, one erect, the other turned backwards; flowers small, purple; perianth-segments narrow-oblong, with an ovate acute tip; style longer than the filaments, pinkish; stigma-lobes, obtuse.

Type locality: Mexico.

Distribution: Central Mexico.

The flowers of this plant were not known when first described nor was its exact origin known. An illustration of it was given. We have also received a dead plant from Bödeker. This illustration and specimen seem to point to a species which has been frequently sent to us from Oaxaca by Conzatti, Reko, and Soils. These plants from Oaxaca normally have 2 short, stout, divergent, central spines. In one specimen sent by Professor Conzatti in 1922 the central spines are often 2 and 4, with one of the centrals more elongated and those near the top of the plant connivent.

Illustration: Monatsschr. Kakteenk. 19: 189, as *Mammillaria pseudoperbella*.

Plate XII, figure 1, shows a plant sent by C. Conzatti from Oaxaca, in 1921. Figure 110 is from a photograph of the type specimen.

62. *Neomammillaria dealbata* (Dietrich).

Mammillaria dealbata Dietrich, Allg. Gartenz. 14: 309. 1846.
Cactus dealbatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Globose to short-cylindric, glaucous, more or less depressed at apex but almost hidden by the many closely appressed spine-clusters; axils of tubercles and young spine-areoles densely lanate but in age glabrate; radial spines about 20, white, short, appressed; central spines 2, much stouter and longer than the radials, sometimes 1 cm. long, the upper ones often erect, white below, brown or black at tip; flowers small, carmine; fruit clavate, red; seeds brown.

Type locality: Mexico.

Distribution: Central Mexico, especially on the pedregal about the City of Mexico.

We have referred to this species a plant which is very common in the Valley of Mexico and which is known in collections as *Mammillaria peacockii*. The name, first used by Rümpler (Förster, Handb. Cact. ed. 2. 286. 1885), was given as a synonym of *Mammillaria dealbata*. It was offered for sale by Grässner as *M. elegans dealbata* (Monatsschr. Kakteenk. February 1920).

Illustration: Grässner, Haupt-Verz. Kakteen 1912: 23, as *Mammillaria peacockii*.

Plate XII, figure 3, shows a plant from Mexico, sent to the New York Botanical Garden in 1911. Figure 111 is from a photograph of a plant sent by Dr. Reiche from the Valley of Mexico in 1922.



FIG. 112.—*Neomammillaria haageana*.




FIG. 113.—*Neomammillaria mundtii*.

63. *Neomammillaria haageana* (Pfeiffer).

Mammillaria haageana Pfeiffer, Allg. Gartenz. 4: 257. 1836.
Mammillaria diacantha Haage in Steudel, Nom. ed. 2. 2: 96. 1841. Not Lemaire, 1838.
Mammillaria haageana validior Monville in Labouret, Monogr. Cact. 54. 1853.
Cactus haageanus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Somewhat caespitose, the individual plants globose or somewhat elongated in age; axils slightly woolly; radial spines about 20, radiating, white; central spines 2, a little longer than the radials, black; flowers small, carmine-rose.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

Pfeiffer (Enum. Cact. 26. 1837) refers here *Mammillaria diacantha nigra* which Haage had listed in his Catalogue of 1836. Here Pfeiffer also refers *M. perote* (Allg. Gartenz. 4: 257. 1836) of gardens.

Illustrations: Dict. Gard. Nicholson 2: 321. f. 509; Cact. Journ. 1: 165; Knippel, Kakteen f. 21; Förster, Handb. Cact. ed. 2. 284. f. 29; Watson, Cact. Cult. 163. f. 62; ed. 3. f. 9; Schelle, Handb. Kakteenk. 262. f. 184; Rümpler, Sukkulanten 201. f. 114, as *Mammillaria haageana*.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering plant of *Neomammillaria pseudoperbella*.
2. Flowering plant of *Neomammillaria spinosissima*.
3. Flowering plant of *Neomammillaria dealbata*.
4. Flowering plant of *Neomammillaria amoena*.
5. Flowering plant of *Neomammillaria polyedra*.
5. Flowering plant of *Neomammillaria celsiana*.

Figure 112 is reproduced from the first illustration cited above. Nicholson recorded the receipt of the plant figured by him from Haage.

64. *Neomammillaria perbella* (Hildmann).

Mammillaria perbella Hildmann in Schumann, Gesamtb. Kakteen 567. 1898.

Solitary or somewhat cespitose, depressed-globose, glaucous-green; tubercles short-conic, their axils lanate; radial spines 14 to 18, 1 to 1.5 mm. long, setaceous, white; central spines 2, very short (4 to 6 mm. long); flowers 9 to 10 mm. long, reddish; stigma-lobes red.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

We know this species from description only; Schumann places it near *Mammillaria donatii*.




FIG. 114.—*Neomammillaria donatii*.




FIG. 115.—*Neomammillaria collina*.

65. *Neomammillaria collina* (J. A. Purpus).

Mammillaria collina J. A. Purpus, Monatsschr. Kakteenk. 22: 162. 1912.

Solitary, globose, 12 to 13 cm. in diameter, somewhat depressed at apex; tubercles cylindric, 1 cm. long or less, woolly in their axils; radial spines 16 to 18, white, 4 mm. long; central spines 1 or 2, longer than the radials; flowers rose-colored, 1.5 to 2 cm. long; fruit 2 cm. long, red.

Type locality: Esperanza, Puebla, Mexico.

Distribution: Puebla, Mexico.

We refer here specimens collected near the type locality in 1912 by Dr. C. A. Purpus.

Illustrations: Monatsschr. Kakteenk. 23: 99; Grässner, Haupt-Verz. Kakteen 1914: 28, as *Mammillaria collina*.

Figure 115 shows a plant sent by Dr. Purpus to Washington.

66. *Neomammillaria donatii* (Berge).

Mammillaria donatii Berge in Schumann, Gesamtb. Kakteen Nachtr. 135. 1903.

Usually simple, stout and globose, but sometimes cespitose, glaucous-green; tubercles small, conic, naked in their axils; radial spines 16 to 18, 8 mm. long, glassy; central spines 2, yellowish black, 10 mm. long; flowers reddish, 15 mm. long; style and stigma-lobes white.

Type locality: Mexico.

Distribution: Mexico.

We do not know the exact type locality or distribution of this plant. It is now in the trade and we recently obtained a specimen from Haage and Schmidt.

Figure 114 is from a photograph of the plant received from Haage and Schmidt in 1920, referred to above.

67. *Neomammillaria mundtii* (Schumann).

Mammillaria mundtii Schumann, Monatsschr. Kakteenk. 13: 141. 1903.

Solitary, so far as known, globose, 6 to 7 cm. in diameter; tubercles not milky, nearly terete, dark green, rather short and stubby, their axils naked; spine-areoles circular, somewhat lanate when young; radial spines 8 to 19, swollen at base, spreading or somewhat curved backward, 6 to 8 mm. long, brownish when young, the tips usually darker; central spines 2, a little stouter and longer than the radials, porrect; flower from toward the center of the plant, 2 cm. long.

Type locality: Not cited.

Distribution: Mexico, but known only from cultivated plants.

We know this plant from a specimen sent to Washington in 1921 by W. Mundt, in whose honor the species had been named.

Illustration: Monatsschr. Kakteenk. 13: 142, as *Mammillaria mundtii*.

Figure 113 is a reproduction of a photograph sent us by L. Quehl in 1921.




FIG. 116.—*Neomammillaria celsiana*.

68. *Neomammillaria celsiana* (Lemaire).

Mammillaria celsiana Lemaire, Cact. Gen. Nov. Sp. 41. 1839.

Mammillaria muehlenpfordtii Förster, Allg. Gartenz. 15: 49. 1847.

Mammillaria schaeferi Fennel, Allg. Gartenz. 15: 66. 1847.

Mammillaria schaeferi longispina Haage, Hamb. Gartenz. 17: 160. 1861.

Cactus muehlenpfordtii Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus celsianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus schaeferi Kuntze, Rev. Gen. Pl. 1: 261. 1891.

(?) *Mammillaria perringii* Hildmann, Gartenwelt 10: 250. 1906.

Plant-body subglobose, becoming cylindric, 10 to 12.5 cm. high, 7.5 cm. in diameter, deep green; axils of tubercles woolly; tubercles conic, compact; spine-areoles small, round, woolly when young; radial spines 24 to 26, about equal, white, setaceous; central spines 4 to 6, rarely 7, somewhat longer than the radials, terete, rigid, pale yellow, more or less recurved and unequal, 8 to 16 mm. long; flowers red; fruit described as green.

Type locality: Not cited.


Distribution: Southern Mexico.

In 1920 Professor Konzatti sent us two specimens from the District of Cuicatlán, Oaxaca, which we refer here; these are the only plants of this species we have seen.

According to Salm-Dyck, *Mammillaria celsiana* differs from *M. rutila* in its columnar stem and in its spines.

Schumann refers *Mammillaria perringii* to *M. celsiana*, while Hildmann claims that it is possible that the two may be distinct, but we do not have the material at hand to decide definitely.

Mammillaria lanifera Haworth (Phil. Mag. 63: 41. 1824; *Cactus lanifer* Kuntze, Rev. Gen. Pl. 1: 260. 1891) is referred here by Schumann; it is probably different but, if not, the name has priority over *M. celsiana*. To *M. lanifera* De Candolle (Prodr. 3: 459. 1828) refers *Cactus canescens* Mociño and Sessé. *M. geminispina monacantha* Lemaire (Cact. Gen. Nov. Sp. 100. 1839) was supposed to be the same as *M. lanifera*. *Mammillaria polycephala* Mühlenpfordt (Allg. Gartenz. 13: 347. 1845; *Cactus polycephalus* Kuntze, Rev. Gen. Pl. 1: 261. 1891) was referred by Schumann to *M. elegans*, but it was described with 4 central spines. It seems to be related to *M. crucigera*, which we have tentatively referred to *M. celsiana*, which has yellow central spines, while both *M. polycephala* and *M. elegans* have white centrals.



FIGS. 117 and 118.—*Neomammillaria aureiceps*.

Mammillaria supertexta dichotoma (Salm-Dyck, Cact. Hort. Dyck. 1849. 9. 1850) is based on *M. polycephala*.

Mammillaria crucigera Martius (Nov. Act. Nat. Cur. 16: 340. pl. 25, f. 2. 1832; *Cactus cruciger* Kuntze, Rev. Gen. Pl. 1: 260. 1891) is related to this species, judging from the description, but the illustration suggests that it is a distinct species. It was collected by Karwinsky in Mexico, but he does not give a definite locality. It was unknown to Schumann.

Illustrations: Gartenwelt 10: 250; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 29, as *Mammillaria celsiana*; Gartenwelt 10: 250, as *Mammillaria perringii*; ?Mém. Mus. Hist. Nat. Paris 17: pl. 4, as *Mammillaria lanifera*; ?Martius, Nov. Act. Nat. Cur. 16: pl. 25, f. 2, as *Mammillaria crucigera*.

Plate XII, figure 6, shows a plant in the New York Botanical Garden which flowered October 16, 1911. Figure 116 is from a photograph of two plants sent by Professor Conzatti in 1920.

69. Neomammillaria aureiceps (Lemaire).*Mammillaria aureiceps* Lemaire, Cact. Aliq. Nov. 8. 1838.*Mammillaria rhodantha aureiceps* Salm-Dyck, Cact. Hort. Dyck. 1844. 7. 1845.*Cactus aureiceps* Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Globose to short-oblong, 8 to 10 cm. in diameter; tubercles short, terete in section, woolly and setose in their axils; radial spines about 20, bristle-like, white, 5 to 8 mm. long, spreading; central spines several, sometimes as many as 9, yellow, stouter and longer than the radials, 10 to 14 mm. long, somewhat spreading and a little curved inward; flowers small, dark red.

Type locality: Mexico.

Distribution: Valley of Mexico.

Our description is based on specimens recently sent us by Dr. Karl Reiche as *Mammillaria rhodantha*, under which name it usually passes. *M. rhodantha*, however, has different spines and is more strictly a mountain species.

Plate IX, figure 3, shows a plant sent from the Edinburgh Botanical Garden in 1902 as *Mammillaria rhodantha* which flowered in the New York Botanical Garden, October 15, 1912. Figures 117 and 118 give two views of this plant sent us by Dr. Reiche from the Valley of Mexico.




FIG. 119.—*Neomammillaria yucatanensis*.




FIG. 120.—*Neomammillaria ruestii*.

70. Neomammillaria yucatanensis sp. nov.

Plants in clumps of 4, erect, cylindric, not milky, 10 to 15 cm. long, 3 to 6 cm. in diameter, very spiny; tubercles conic, woolly in their axils but not setose; radial spines about 20, white, spreading, acicular; central spines 4 or rarely, much stouter than the radials, 6 to 8 mm. long, slightly spreading above, yellowish brown; flowers very small, rose; fruit oblong, bright red."

Collected by George F. Gaumer at Progreso, Yucatan, Mexico, in 1918 (No. 23939) and again in 1921 (No. 24367, type).

We have not seen this species in flower or fruit but Dr. Gaumer has described them as above. He says that the plant is rare on the land side of the coastal marshes.

Figure 119 is from a photograph of the plant sent in 1921 by Dr. Gaumer.

71. *Neomammillaria ruestii* (Quehl).*Mammillaria ruestii* Quehl, Monatsschr. Kakteenk. 15: 173. 1905.*Mammillaria celsiana guatemalensis* Eichlam, Monatsschr. Kakteenk. 19: 59. 1909.

Cylindric, 6 to 7 cm. high, 4 to cm. in diameter, light green, almost hidden by the spines; axils of tubercles more or less woolly, at least when young; flowering areoles at first quite woolly; radial spines 20 or more, white, glossy, to 6 mm. long, spreading; central spines usually 4, sometimes, much stouter than the radials, yellow, swollen at base, ascending, 7 to 8 mm. long; flowers small, sometimes almost hidden by the spines, 8 mm. long; inner perianth-segments about 25, lanceolate, acute, pale purple, the margins almost colorless; filaments colorless below, purplish above; style pale; stigma-lobes 4, linear, elongated, reflexed; fruit clavate, red; seeds brown.

Type locality: Honduras.

Distribution: Honduras and Guatemala.

We have had the Guatemala plant under observation for 14 years and it has both flowered and fruited.

Figure 120 is from a photograph of a plant sent by Dr. A. W. Kellermann from Guatemala in 1908.



FIG. 121.—*Neomammillaria pringlei*.

72. *Neomammillaria pringlei* (Coulter).*Cactus pringlei* Coulter, Contr. U. S. Nat. Herb. 3: 109. 1894.*Mammillaria pringlei* K. Brandege, Zoe 5: 7. 1900.

Solitary, with long fibrous roots, usually globose, but sometimes depressed or short-cylindric, 6 to 16 cm. high, 6 to 7 cm. in diameter; tubercles dull green, terete, conic, 6 to 10 mm. long; axils of tubercles woolly and setose; spines all yellow; radial spines 18 to 20, setaceous, spreading, 5 to 8 mm. long; central spines 5 to 7, much stouter and longer than the radials, more or less recurved, 2 to 2.5 cm. long, those from the upper areoles curved over the apex of the plant; flowers deep red, 8 to 10 mm. long; fruit borne in a circle near the middle of the plant, oblong, 12 to 15 mm. long; seeds small, brown.

Type locality: Cited as San Luis Potosí, but doubtless Tultenango Canyon, state of Mexico, according to Pringle, who collected the type.

Distribution: Known only from the type locality.

Dr. Rose collected living specimens from the type locality some years ago but these never flowered. In April 1921 we sent Dr. Reiche to the type locality and he obtained thirteen beautiful specimens, one of which was in fruit.

Coulter (Contr. U. S. Nat. Herb. 3: 109. 1894) states that *Cactus pringlei* was near *Cactus rhodanthus sulphureospinus*, which was based on *M. sulphurea* Förster.

Figure 121 is from a photograph of the plants collected at Tultenango Canyon in 1921.

73. *Neomammillaria cerralboa* sp. nov.

Cylindric, solitary, 1 to 1.5 dm. high, 5 to 6 cm. in diameter; tubercles not milky, yellowish, terete, obtuse, closely set; spines all yellow, very much alike, about 11, one usually more central, the longer ones nearly 2 cm. long; flowers small, 1 cm. long or less, forming a circle around the plant about 3 cm. below the top.

Collected by Ivan M. Johnston on Cerralbo Island, Gulf of California, June 6, 1921 (No. 4038). The next day on the same island he collected three more plants (No. 4053) which seem to be referable here, except that two of them have hooked spines; Dr. Rose also collected on this same island (No. 16877) in 1911 specimens with hooked spines which are like Mr. Johnston's plant. Whether this plant has normally these two forms or whether the hooked-spined one is a hybrid we are unable to determine.

Figure 121a is a photograph of the type plant, collected by Johnston (No. 4038).




FIG. 121a.—*Neomammillaria cerralboa*.

74. *Neomammillaria phaeacantha* (Lemaire).

Mammillaria phaeacantha Lemaire, Cact. Gen. Nov. Sp. 47. 1839.

Mammillaria nigricans Fennel, Allg. Gartenz. 15: 66. 1847.

Cactus nigricans Kuntze, Rev. Gen. Pl. 1: 261. 1891. Not Haworth, 1803.

Cactus phaeacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.




FIG. 122.—*Neomammillaria phaeacantha*.




FIG. 123.—*Neomammillaria graessneriana*.

Globose or somewhat depressed, green; axils of tubercles woolly; tubercles conic, hardly, if at all, angled; spine-areoles small, yellowish tomentose (probably so only when young); radial spines

16 to 20, white, setaceous; central spines 4, black, subulate, spreading or reflexed, the lowest one longest; flowers from upper part of plant, dark red; perianth-segments oblong, acuminate.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

This species has not been recognized by recent writers, and while we have seen no specimens we believe it deserves specific rank.

Schumann refers *Mammillaria nigricans* definitely to *M. rhodantha* but, it appears to us, without justification; the Index Kewensis has referred it, we believe properly, to *M. phaeacantha*.

Mammillaria phaeacantha rigidior (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845) is only a name.

Illustration: Pfeiffer, Abbild. Besch. Cact. 2: pl. 23, as *Mammillaria nigricans*.

Figure 122 is reproduced from the illustration cited above.

75. *Neomammillaria graessneriana* (Bödeker).

Mammillaria graessneriana Bödeker, Monatsschr. Kakteenk. 30: 84. 1920.



Solitary, or becoming cespitose, globose, 6 to 8 cm. in diameter, dark bluish green, somewhat depressed at apex; tubercles 4-angled, 8 mm. long, not milky, obtuse or truncate at apex, not setose in their axils; spine-areoles circular, white-woolly when young, nearly naked in age; radial spines 18 to 20, acicular, 6 to 8 mm. long, white; central spines 2 to 4, stouter than the radials, spreading, 8 mm. long, reddish brown; flowers small, somewhat distant from the apex of the plant.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

Illustration: Monatsschr. Kakteenk. 30: 85, as *Mammillaria graessneriana*.

Figure 123 is reproduced from the illustration cited above.



FIGS. 124 and 125.—*Neomammillaria spinosissima*.

76. *Neomammillaria spinosissima* (Lemaire).

Mammillaria spinosissima Lemaire, Cact. Aliq. Nov. 4. 1838.

Mammillaria polycentra Berg, Allg. Gartenz. 8: 130. 1840.

Mammillaria auricoma Dietrich, Allg. Gartenz. 14: 308. 1846.

Mammillaria polyacantha Ehrenberg, Allg. Gartenz. 16: 265. 1848.

Mammillaria polyacantha Ehrenberg, Allg. Gartenz. 16: 266. 1848.

Mammillaria hepatica Ehrenberg, Allg. Gartenz. 16: 267. 1848.

Mammillaria pomacea Ehrenberg, Allg. Gartenz. 16: 267. 1848.

Mammillaria pulcherrima Ehrenberg, Allg. Gartenz. 17: 249. 1849.

Mammillaria pretiosa Ehrenberg, Allg. Gartenz. 17: 250. 1849.

- Mammillaria caesia* Ehrenberg, Allg. Gartenz. 17: 251. 1849.
Mammillaria mirabilis Ehrenberg, Allg. Gartenz. 17: 251. 1849.
Mammillaria pruinosa Ehrenberg, Allg. Gartenz. 17: 261. 1849.
Mammillaria seegeri Ehrenberg, Allg. Gartenz. 17: 261. 1849.
Mammillaria baseloffii Ehrenberg, Allg. Gartenz. 17: 261. 1849.
Mammillaria herrmannii Ehrenberg, Allg. Gartenz. 17: 303. 1849.
Mammillaria aurea Ehrenberg, Allg. Gartenz. 17: 303. 1849.
Mammillaria linkeana Ehrenberg, Allg. Gartenz. 17: 308. 1849.
Mammillaria vulpina Ehrenberg, Allg. Gartenz. 17: 308. 1849.
Mammillaria eximia Ehrenberg, Allg. Gartenz. 17: 309. 1849.
Mammillaria isabellina Ehrenberg, Allg. Gartenz. 17: 309. 1849.
Mammillaria spinosissima brunnea Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria spinosissima flavida Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria spinosissima rubens Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria herrmannii flavicans Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria seegeri gracilispina Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria seegeri pruinosa Salm-Dyck, Cact. Hort. Dyck. 1849. 8. 1850.
Mammillaria uhdeana Salm-Dyck, Cact. Hort. Dyck. 1849. 83. 1850.
Mammillaria spinosissima hepatica Labouret, Monogr. Cact. 35. 1853.
Mammillaria castaneoides Lemaire in Labouret, Monogr. Cact. 37. 1853.
Mammillaria seegeri mirabilis Labouret, Monogr. Cact. 37. 1853.
Mammillaria sanguinea Haage jr. in Regel, Act. Hort. Petrop. 8: 276. 1883.
Mammillaria poselgeriana Haage in Förster, Handb. Cact. ed. 2. 269. 1885.
Mammillaria pretiosa cristata Hildmann in Förster, Handb. Cact. ed. 2. 273. 1885.
Cactus auricomus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus aureus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus eximius Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus isabellinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus linkeanus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus mirabilis Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus polycentrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pomaceus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pretiosus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pulcherrimus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus spinosissimus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus vulpinus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Mammillaria spinosissima sanguinea Haage in Brandegee, Cycl. Amer. Hort. Bailey 2: 976. 1900.
Mammillaria spinosissima aurea Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima auricoma Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima eximia Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima baseloffii Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima herrmannii Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima isabellina Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima linkeana Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima mirabilis Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima pruinosa Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima pulcherrima Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima seegeri Gürke, Blühende Kakteen 2: under pl. 71. 1905.
Mammillaria spinosissima vulpina Gürke, Blühende Kakteen 2: under pl. 71. 1905.

Cylindric, 7 to 30 cm. long, 2.5 to 10 cm. in diameter, almost hidden under a dense covering of spines; axils of tubercles setose; tubercles very short, 2 to 3 mm. long; spines brownish to red, usually weak, hardly pungent; radial spines about 20, 1 cm. long or less; central spines 7 or 8, 2 cm. long or more; flowers from the upper part of the plant, purplish, 12 mm. long; inner perianth-segments acute; filaments much shorter than the perianth-segments, purple.

Type locality: Not cited.

Distribution: Mountains of central Mexico.

The above description is drawn from collections obtained in the high mountains between the City of Mexico and Cuernavaca. There seems to be little doubt but that they are the *M. sanguinea* Haage which Schumann refers to *M. spinosissima*.

We are disposed to refer here *Echinocactus spinosissimus* (Forbes, Journ. Hort. Tour Germ. 152. 1837). Forbes did not have much knowledge of the cacti but was the gardener of the Duke of Bedford, who sent him to the Continent of Europe in 1835, where he obtained many cacti and on his return to England published a list of them, sometimes with brief descriptions. The names had been given to him by Pfeiffer and others who were studying this family. As he published his list very promptly after his return to England many names appear there first or in the same year as in Pfeiffer's Enumeratio. *Mammillaria*

spinosissima may have been in cultivation at the time of Forbes's visit to Germany, for it was published in 1838.

Illustrations: Möllers Deutsche Gärt. Zeit. 25: 475. f. 8. No. 26, as *Mammillaria poselgeriana*; Gartenflora 32: pl. 111; Dict. Gard. Nicholson 2: 322. f. 510; Förster, Handb. Cact. ed. 2. 271. f. 28; Watson, Cact. Cult. 172. f. 68; ed. 3. f. 46, as *Mammillaria sanguinea*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 11, as *Mammillaria eximia*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 18, as *Mammillaria spinosissima auricoma*; Balt. Cact. Journ. 2: 150, as *M. spinosissima brunnea*; Möllers Deutsche Gärt. Zeit. 25: 487. f. 21; Cact. Journ. 2: 93; Blanc, Cacti 74. No. 1580; Schelle, Handb. Kakteenk. 253. f. 174; Blühende Kakteen 2: pl. 71, as *Mammillaria spinosissima*.

Plate XII, figure 2, shows a plant collected by Dr. Rose at El Parque, Mexico, in 1906. Figure 124 is from a photograph of a plant sent to the New York Botanical Garden by Frank Weinberg in 1906 as *Cactus spinosissimus*; figure 125 is from a photograph of a plant sent by William Brockway from the mountains above the City of Mexico.




FIG. 126.—*Neomammillaria densispina*.




FIG. 127.—*Neomammillaria nunezii*.

77. *Neomammillaria densispina* (Coulter).

Cactus densispinus Coulter, Contr. U. S. Nat. Herb. 3: 96. 1894.

Mammillaria pseudofuscata Quehl, Monatsschr. Kakteenk. 24: 114. 1914.

Globose, 6 to 10 cm. in diameter, entirely hidden by the dense covering of spines; tubercles short and thick, green, not milky; radial spines 25 or more, slightly spreading, about x cm. long, whitish or pale yellow; central spines 5 or 6, longer than the radials, 10 to 12 mm. long, the upper half or third dark brown; flowers purple without, yellowish within, 1.5 cm. long; seeds obovate, reddish brown, 1 mm. in diameter.

Type locality: San Luis Potosí, Mexico.

Distribution: San Luis Potosí, Mexico.

We have had this plant in cultivation since 1912, specimens having been sent to Washington by Mrs. Irene Vera from San Luis Potosí. Our plant is probably a part of the type collection of Quehl's *Mammillaria pseudofuscata*, as Mrs. Vera wrote us that she had sent specimens to Germany which had been identified as *M. fuscata*. Our plant has been compared with Eschancier's specimen from the same locality which is the type of Coulter's

Cactus densispinus and we are convinced that they are the same; Coulter's type is now in the Field Museum of Natural History.

Illustration: Monatsschr. Kakteenk. 24: 115, as *Mammillaria pseudofuscata*.

Figure 126 shows the plant sent by Mrs. Vera from San Luis Potosí.

78. *Neomammillaria nunezii* sp. nov.

Globose to cylindric, 1.5 cm. long, 6 to 8 cm. in diameter; tubercles closely set, short, terete in section, setose in their axils; radial spines white, stiff, about 30, widely spreading; central spines 2 to 4, stout, 10 to 15 mm. long, brown to nearly blackish at tips; fruit 2.5 cm. long, clavate, white or tinged with pink; seeds small, brown.




FIG. 128.—*Neomammillaria nunezii*.

FIG. 129.—*Neomammillaria rhodantha*.

Collected by Professor C. Núñez at Buenavista de Cuellar, Guerrero, Mexico, in 1921 (Nos. 1, 2 and 3), and communicated to us by Octavio Solís. This species is rather variable in habit and spines and is very unlike anything that we have heretofore studied.

Figures 127 and 128 are from photographs of the top and side of two plants of this collection.

79. *Neomammillaria amoena* (Hoppfer).

Mammillaria amoena Hoppfer in Salm-Dyck, Cact. Hort. Dyck. 1849. 99. 1850.

Stems robust, columnar; tubercles green, ovoid, obtuse, subglaucous; radial spines 16, slender, radiating, white; central spines 2, rigid, yellowish brown, 8 to 10 mm. long, the upper one longer and recurved; flowers appearing from axils above middle of plant, 2 cm. long; tube cone-shaped, green; outer perianth-segments somewhat brownish; inner perianth-segments with a pale-brown central stripe; margins nearly white, obtuse, entire; stamens short; filaments pale; anthers red; style pale green; stigma-lobes green, linear.

Type locality: Not cited.

Distribution: Central Mexico.

Förster's Handbuch (254. 1846) is often given as the place of publication, but while the name is found in the place cited it is without description.

Plate XII, figure 4, shows a plant which flowered in the New York Botanical Garden in 1912, sent from Cuernavaca, Mexico, by Wm. Brockway the preceding year. Figure 130 is from a photograph of a plant from the same collection which flowered in Washington.




FIG. 130.—*Neomammillaria amoena*.



FIG. 131.—*Neomammillaria plumosa*.

80. *Neomammillaria rhodantha* (Link and Otto).

- Mammillaria rhodantha* Link and Otto, Icon. Pl. Rar. 51. 1829.
Mammillaria pulchra Haworth in Edwards's Bot. Reg. 16: pl. 1329. 1830.
Mammillaria fulvispina Haworth, Phil. Mag. 7: 108. 1830.
 ? *Mammillaria inuncta* Hoffmannsegg, Preiss-Verz. ed. 7. 23. 1833.
Mammillaria erinacea Wendland, Cact. Herrenh. 1835.
Mammillaria chrysacantha Otto in Pfeiffer, Enum. Cact. 28. 1837.
Mammillaria fuscata Pfeiffer, Enum. Cact. 28. 1837.
Mammillaria tentaculata Otto in Pfeiffer, Enum. Cact. 29. 1837.
Mammillaria rhodantha rubens Pfeiffer, Enum. Cact. 31. 1837.
Mammillaria rhodantha andreae * Otto in Pfeiffer, Enum. Cact. 31. 1837.
Mammillaria rhodantha prolifera Pfeiffer, Enum. Cact. 31. 1837.
Mammillaria rhodantha neglecta Pfeiffer, Enum. Cact. 31. 1837.
Mammillaria rhodantha wendlandii Pfeiffer, Enum. Cact. 31. 1837.
Mammillaria ruficeps Lemaire, Cact. Gen. Nov. Sp. 37. 1839.
Mammillaria odieriana Lemaire, Cact. Gen. Nov. Sp. 46. 1839.
Mammillaria pyrrhocbracantha Lemaire, Cact. Gen. Nov. Sp. 51. 1839.
Mammillaria rhodantha major Monville in Lemaire, Cact. Gen. Nov. Sp. 98. 1839.
Mammillaria pfeifferi Booth in Scheidweiler, Bull. Acad. Sci. Brux. 6: 93. 1839.
Mammillaria pfeifferi altissima Scheidweiler, Bull. Acad. Sci. Brux. 6: 93. 1839.
Mammillaria pfeifferi dichotoma Scheidweiler, Bull. Acad. Sci. Brux. 6: 93. 1839.
Mammillaria pfeifferi flaviceps Scheidweiler, Bull. Acad. Sci. Brux. 6: 93. 1839.
Mammillaria pfeifferi fulvispina Scheidweiler, Bull. Acad. Sci. Brux. 6: 3. 1839.
Mammillaria pfeifferi variabilis Scheidweiler, Bull. Acad. Sci. Brux. 6: 93. 1839.
 ? *Mammillaria crassispina* Pfeiffer, Allg. Gartenz. 8: 406. 1840.
 ? *Mammillaria stenocephala* Scheidweiler, Allg. Gartenz. 9: 43. 1841.
 ? *Mammillaria imbricata* Wegener, Allg. Gartenz. 12: 66. 1844.
Mammillaria crassispina gracilior Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845.
Mammillaria rhodantha centrispina Link in Förster, Handb. Cact. 198. 1846.
Mammillaria sulphurea Sencke in Förster, Handb. Cact. 200. 1846.
Mammillaria robusta Otto in Förster, Handb. Cact. 207. 1846.
Mammillaria tentaculata ruficeps Förster, Handb. Cact. 207. 1846.
Mammillaria stueberi Otto in Förster, Handb. Cact. 517. 1846.
Mammillaria fulvispina rubescens Salm-Dyck, Cact. Hort. Dyck. 1849. 10. 1850.

* *Mammillaria andreae* was used by Schumann (Gesamtb. Kakteen 598. 1898).

- Mammillaria rhodantha sulphurea* Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850.
Mammillaria rhodantha ruficeps Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850.
Mammillaria chrysacantha fuscata Salm-Dyck, Cact. Hort. Dyck. 1849. 12. 1850.
Mammillaria rhodantha rubescens Salm-Dyck, Cact. Hort. Dyck. 1849. 97. 1850.
Mammillaria odieriana rigidior Salm-Dyck, Cact. Hort. Dyck. 1849. 98. 1850.
Mammillaria lanifera Salm-Dyck, Cact. Hort. Dyck. 1849. 98. 1850. Not Haworth, 1824.
? *Mammillaria russea* Dietrich, Allg. Gartenz. 19: 347. 1851.
Mammillaria odieriana rubra Sencke in Förster, Handb. Cact. ed. 2. 295. 1885.
Mammillaria odieriana cristata Hortus in Förster, Handb. Cact. ed. 2. 295. 1885.
Mammillaria tentaculata picta Förster, Handb. Cact. ed. 2. 309. 1885.
Mammillaria crassispina rufa Rümpler in Förster, Handb. Cact. ed. 2. 311. 1885.
Cactus chrysacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus crassispinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus fuscatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.
Cactus odieranus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus pyrrhocbroacanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus rhodanthus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus ruficeps Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus stenocephalus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus stueberi Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus tentaculatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Cactus capillaris Coulter, Contr. U. S. Nat. Herb. 3: 107. 1894.
Cactus rhodanthus sulphureospinus Coulter, Contr. U. S. Nat. Herb. 3: 107. 1894.
Mammillaria rhodantha pfeifferi Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha rubra Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha ruberrima Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha pyramidalis Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha callaena Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha crassispina Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha droegeana Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha chrysacantha Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha stenocephala Schumann, Gesamtb. Kakteen 550. 1898.
Mammillaria rhodantha fuscata Schumann, Gesamtb. Kakteen 551. 1898.
Mammillaria rhodantha odieriana Schelle, Handb. Kakteenk. 257. 1907.
Mammillaria rhodantha fulvispina Schelle, Handb. Kakteenk. 257. 1907.
Mammillaria rhodantha tentaculata Hortus in Schelle, Handb. Kakteenk. 257. 1907.

Cylindric, 1 to 3 dm. long, erect, dull green; tubercles terete, somewhat narrowed toward the apex, 3 to 5 mm. long, not yielding milk when pricked; axils of tubercles sometimes bearing bristles, often naked; radial spines 15 to 20, white, 5 to 7 mm. long; central spines 4 to 6, reddish brown, straight, ascending, much stouter than the radials, 10 to 12 mm. long; flowers numerous, rose-colored, 12 mm. broad; inner perianth-segments linear, somewhat spreading, pointed; filaments red; stigma-lobes 4 or 5, rose-colored; fruit 2.5 cm. long, cylindric, lilac to red; seeds brownish.

Type locality: Mexico.

Distribution: Probably central Mexico.

We have had this plant in cultivation but it has never flowered with us; however, it is very distinct from anything else we know.

Mammillaria flaviceps is referred by Labouret to *M. crassispina*, now usually referred to *M. rhodantha*.

Mammillaria floccigera (and its variety *longispina*) Förster (Handb. Cact. 254. 1846), as well as *M. aurata* and *M. hybrida* (Pfeiffer, Enum. Cact. 31. 1837), are given by Schumann as synonyms of *M. rhodantha*, but none of them was described at the places cited.

Mammillaria erinacea Wendland is unknown to us; it is referred to *Mammillaria rhodantha* by the Index Kewensis, but whether it was described or not we do not know.

Mammillaria fulvispina was said by Haworth to come from Brazil, while the Index Kewensis refers it to Brazil and Mexico. If of this relationship, it is from Mexico. *Mammillaria radula* Scheidweiler (Förster, Handb. Cact. 208. 1846), referred by Schumann as a synonym of this species, was given by Förster as a synonym of *Mammillaria phaeacantha*.

Mammillaria pyramidalis Link and Otto (Verh. Ver. Beförd. Gartenb. 6: 429. 1830), given as a synonym of this species by Schumann, is only a name.

Mammillaria atrata Mackie (Curtis's Bot. Mag. 65: pl. 3642. 1839) is also referred here by Schumann. The plant is supposed to have come from Chile and is probably referable to *Neoporteria*, which see (Cactaceae 3: 97. 1922).

Mammillaria pyrrhocentra Otto, its var. *gracilior* (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845), and *M. fulvispina pyrrhocentra* Salm-Dyck (Cact. Hort. Dyck. 1849. 10. 1850) were referred as synonyms of *M. rhodantha* by Schumann, but were not described at the places cited.

Mammillaria aurea Pfeiffer (Förster, Handb. Cact. 200. 1846; *M. rhodantha aurea* Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850) is referred here. We have found no description of it. *M. odieriana aurea* Salm-Dyck (Cact. Hort. Dyck. 1844. 7. 1845), also undescribed, may be the same.

Mammillaria rhodantha cristata (Förster, Handb. Cact. ed. 2. 292. 1885) is only an abnormal form.

Mammillaria recurvispina Hildmann (Schelle, Handb. Kakteenk. 257. 1907) is given without synonymy or description. *M. rhodantha schochiana* (*M. schochiana* Hortus) is also given at the same place, but so far as we can learn has not been published.

Mammillaria tentaculata conothele Monville is given by Labouret (Monogr. Cact. 55. 1853) as a synonym of *M. stueberi*, while he refers *M. tentaculata fulvispina* (Monogr. Cact. 44. 1853) to *M. fulvispina*.

Mammillaria tentaculata rubra (Förster, Handb. Cact. 207. 1846) was given as a synonym of *M. tentaculata ruficeps*.

Mammillaria olivacea was cited by Pfeiffer (Enum. Cact. 180. 1837) as a synonym of *M. tentaculata*.

Mammillaria neglecta was given as a synonym of *M. rhodantha neglecta* by Salm-Dyck (Cact. Hort. Dyck. 1849. 11. 1850).

Mammillaria rhodantha var. *inuncta* Hoffmannsegg was listed by Labouret (Monogr. Cact. 45. 1853) as one of the synonyms of *M. rhodantha*. *M. rhodantha rubra* was given by Rümpler (Förster, Handb. Cact. ed. 2. 292. 1885) as a synonym of *M. rhodantha ruficeps*, but afterwards was formally published by Schumann.

Mammillaria rhodantha celsii Lemaire (Labouret, Monogr. Cact. 48. 1853) was given as a synonym of *M. lanifera*. It probably belongs to *M. rhodantha*. *Cactus capillaris* was made by Coulter because of the older *Mammillaria lanifera* of Haworth. Palmer's plant from Saltillo (1880), preserved in the Missouri Botanical Garden, is very different and suggests that the labels have been mixed.

Illustrations: Knippel, Kakteen pl. 24; De Laet, Cat. Gén. f. 50, No. 9; Wiener Illustr. Gart. Zeit. 29: f. 22, No. 9; Haage and Schmidt, Haupt-Verz. Cact. 1912: 37; Link and Otto, Icon. Pl. Rar. pl. 26; Gard. Chron. 111. 42: 290. f. 116; Schelle, Handb. Kakteenk. 256. f. 179; Gartenwelt 1: 200; Abh. Bayer. Akad. Wiss. München 2: pl. 1, 1. f. 3; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 22, as *Mammillaria rhodantha*; Grässner, Haupt-Verz. Kakteen 1912: 24; Schelle, Handb. Kakteenk. 258. f. 181, as *Mammillaria rhodantha pfeifferi*; Grässner, Haupt-Verz. Kakteen 1912: 24, as *M. rhodantha fuscata*; Schelle, Handb. Kakteenk. 258. f. 180, as *M. rhodantha fulvispina*; Blanc, Cacti 73. No. 1434; Cact. Journ. 1: 43, as *M. odieriana*; Cact. Journ. 1: 43; pl. for February, as *M. pfeifferi*; Edwards's Bot. Reg. 16: pl. 1329, as *M. pulchra*; Nov. Act. Nat. Cur. 19: pl. 16, f. 8, as *Mammillaria tentaculata*.

Figure 129 is from a photograph of a plant obtained by Dr. Rose through W. Mundt in 1913, which is now growing at Washington.

81. *Neomammillaria plumosa* (Weber).

Mammillaria plumosa Weber, Dict. Hort. Bois 804. 1898.

Small, growing in dense clusters sometimes 15 cm. broad, entirely covered by the mass of white spines; tubercles small, somewhat woolly in their axils, 2 to 3 mm. long; spines about 40, all radial, weak, plumose, 3 to 7 mm. long; flowers white, small, 3 to 4 mm. long; perianth-segments with a red line running down the center; seeds black.

Type locality: Northern Mexico.

Distribution: Northern Mexico.

This plant for a long time passed in the trade under the name of *Mammillaria lasiocantha*, but it is, of course, very different. It is a very striking species and differs from all the others in its feather-like spines. We have had it under observation since 1907 and it has only once flowered (1921).

According to Walton, it is called the feather ball on account of the feather-like spines.

Illustrations: Möllers Deutsche Gart. Zeit. 25: 475. f. 8, No. 16; Schelle, Handb. Kakteenk. 252. f. 173; Ann. Rep. Smiths. Inst. 1908: pl. 3, f. 6; Haage, Cact. Kultur ed. 2. 189; Journ. Hort. Home Farm. 111. 60: 7, as *Mammillaria plumosa*; Cact. Journ. 1: pl. for February, in part; Darel, Illustr. Handb. Kakteenk. 94. f. 76; Blanc, Hints on Cacti 70. f. 1355; Blanc, Illustr. Price List Cacti 13, as *M. lasiacantha*.

Figure 131 is from a photograph, furnished by Dr. Safford, showing the spines.

82. *Neomammillaria prolifera* (Miller).

Cactus proliferus Miller, Gard. Dict. ed. 8. No. 6. 1768.

Cactus glomeratus Lamarck, Encycl. 1: 537. 1783.

Cactus mammillaris prolifer Aiton, Hort. Kew. 2: 150. 1789.

Mammillaria prolifera Haworth, Syn. Pl. Succ. 177. 1812.

Cactus pusillus De Candolle, Cact. Hort. Monsp. 184. 1813. Not Haworth, 1803.

Cactus stellatus Willdenow, Enum. Pl. Suppl. 30. 1813.

Mammillaria stellaris Haworth, Suppl. Pl. Succ. 72. 1819.

Mammillaria pusilla Sweet, Hort. Brit. 171. 1826.

Mammillaria stellata Sweet, Hort. Brit. 171. 1826.

Mammillaria glomerata De Candolle, Prodr. 3: 459. 1828.

Mammillaria pusilla major Pfeiffer, Enum. Cact. 36. 1837.

Cactus haworthianus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Mammillaria pusilla haitiensis Schumann, Blühende Kakteen 1: under pl. 46. 1904.

Low, growing in colonies often 6 dm. in diameter, the individual plants globose or cylindric, 3 to 6 cm. in diameter, of soft texture; tubercles conic, about 8 mm. long, spreading; axils of tubercles with long, hair-like bristles; radial spines many, hair-like; central spines 5 to 12, much stouter than the radials, with bright yellow tips, puberulent; flowers borne in old axils but toward top of plant, small, yellowish white; inner perianth-segments erect, pale yellow, with brownish mid-rib, acute; filaments pale rose-colored; anthers at first deflexed inward; style shorter than filaments; stigma-lobes, yellow; fruit crowned by persistent withering perianth, clavate, somewhat curved, 1.5 to 2 cm. long, scarlet; seeds black, pitted, a little depressed; aril white, triangular.

Type locality: West Indies.

Distribution: Cuba and Hispaniola. Loddiges reports it from South America, doubtless in error.

At the United States Naval Station, Guantánamo Bay, Cuba, the plant grows in low, dry thickets and is quite inconspicuous but abundant.

Dr. Shafer referred to this species (Bull. N. Y. Bot. Gard. 13: 139) as *Mammariella*, without description or citation.

Burmann's plate (201, f. 1) of this plant shows most of the tubercles without spines or hairs but these have doubtless been omitted by the artist, for Plumier says (Cat. p. 19): "*Melocactus minimus, lanuginosus et tuberosus*."

Haworth (Phil. Mag. 7: 114. 1830) would exclude *Mammillaria pusilla* (Mém. Mus. Hist. Nat. Paris 17: pl. 2, f. ,) as figured by De Candolle. His illustration is evidently faulty, but his description seems to answer our plant.

The name *Mammillaria pusilla minor* occurred in the Index of the Cacti in the Botanical Garden of Berlin for 1829 (Verh. Ver. Beförd. 6: 429. 1830), but it is without description. It is mentioned again by Salm-Dyck (Hort. Dyck. 156. 1834), who credits the name to Otto, but he does not describe it.

* Otto Kuntze (Rev. Gen. Pl. 1: 259. 1891) publishes this binomial as *Cactus prolifer*. Pfeiffer (Enum. Cact. 9. uses this later binomial for another species, crediting it to Willdenow, but we do not find it used elsewhere.

Mammillaria granulata Meinshausen (Wöchenschr. Gärt. Pflanz. 1: 264. 1858; *Cactus granulatus* Kuntze, Rev. Gen. Pl. 1: 260. 1891) was described without the flowers and fruit being known and it has never been identified. Meinshausen says that it has the habit of *M. pusilla*, but he considered it different otherwise.

Cactus stellaris was given by Haworth (Suppl. Pl. Succ. 72. 1819) instead of *C. stellatus* Willdenow.

Mammillaria pusilla cristata (Schelle, Handb. Kakteenk. 249. 1907) is probably only a form.

Illustrations: Loudon, Encycl. Pl. 410. f. 6842, as *Cactus stellaris*; Loddiges, Bot. Cab. 1: pl. 79, as *Cactus stellatus*; Plukenet, Opera Bot. 1: pl. 29, f. 2, as *Ficoides* etc.; Dict. Hort. Nicholson Suppl. 514. f. 547; Abh. Bayer. Akad. Wiss. München 2: pl. 1, VIII, f. 7; Rümpler, Sukkulente 197. f. 110; Monatsschr. Kakteenk. 8: 73; Mém. Mus. Hist. Nat. Paris 17: pl. 2, f. 1; Ann. Rep. Smiths. Inst. 1908: pl. 2, f. 4; Blanc, Cacti 74, No. 1500; Schumann, Gesamtb. Kakteen f. 87; Blühende Kakteen 1: pl. 46; Ann. Inst. Roy. Hort. Fromont 2: pl. 1, f. B; Watson, Cact. Cult. ed. 2. 255. f. 96; ed. 3. f. 45; Remark, Kakteenfreund 15; Cact. Journ. 2: 6, as *Mammillaria pusilla*.

Figure 132 is from a photograph by Ernest Branton of a clump of plants growing in the Huntington collection near Los Angeles, California.




FIG. 132.—*Neomammillaria prolifera*.



FIG. 133.—*Neomammillaria multiceps*.

83. *Neomammillaria multiceps* (Salm-Dyck).

Mammillaria multiceps Salm-Dyck, Cact. Hort. Dyck. 1849.81. 1850.

Mammillaria multiceps elongata Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria multiceps grisea Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria multiceps humilis Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria multiceps perpusilla Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria pusilla texana Engelm., Cact. Mex. Bound. 5. 1859.

Mammillaria texana Poselger in Young, Fl. Texas. 279. 1873.

Cactus multiceps Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus stellatus texanus Coulter, Contr. U. S. Nat. Herb. 3: 108. 1894.

Cactus texanus Small, Fl. Southeast. U. S. 812. 1903.

Cespitose, often forming large clumps; separate plants globose to short-oblong, often only 1 to 2 cm. in diameter; tubercles small, terete, hairy in their axils; radial spines hair-like, white; central spines several, pubescent, yellowish at base, dark brown above; flowers about 12 mm. long, whitish to yellowish salmon, often becoming reddish on outside; fruit oblong, 8 to 12 mm. long, scarlet; seeds black, 1 mm. long, punctate.

Type locality: Not cited.

Distribution: Texas and northeastern Mexico.

It is sometimes classified as a variety of *Mammillaria prolifera*, from which it differs in having the central spines always brown-tipped instead of golden yellow; it is somewhat smaller, with slightly smaller seeds.

Mr. Robert Runyon says that this plant forms clumps usually about 10 cm. broad, but sometimes broader. It is never very plentiful but has a rather wide distribution, and seems to prefer mesquite thickets where the soil is very rich, but occasionally is found on rocky hillsides.

Mammillaria pusilla mexicana, offered for sale by Grässner (Monatsschr. Kakteenk. February 1920), probably belongs here.

Mammillaria caespititia Hortus was referred by Salm-Dyck as a synonym of *M. multiceps*. *M. pusilla caespititia* (Schelle, Handb. Kakteenk. 249. 1907) is the same.

Mammillaria parvissima Karwinsky (Wöchenschr. Gärt. Pflanz. 1: 27. 1858) is sometimes credited to Meinshausen, but seems never to have been described. *M. perpusilla* Meinshausen, given only as a synonym, belongs here and the name occurs on the page mentioned above.




Fig. 134.—*Neomammillaria multiceps*.

Illustrations: Cact. Mex. Bound. pl. 5; Cact. Journ. 2: 93; Förster, Handb. Cact. ed. 2. 262. f. 25; Schelle, Handb. Kakteenk. 249. f. 68, as *Mammillaria pusilla texana*.

Plate XIV, figure 5, shows a very small plant in flower, collected by Robert Runyon near Brownsville, Texas, in 1921; figure 6 shows a plant received from the Missouri Botanical Garden in 1904 which flowered in the New York Botanical Garden in March 1912. Figure 134 is from a photograph of a plant collected near Victoria, Mexico, by Dr. Edward Palmer, which was grown for many years in Washington; figure 133 shows a small plant photographed by Robert Runyon on July 10, 1921.

84. *Neomammillaria camptotricha* (Dams).

Mammillaria camptotricha Dams, Gartenwelt 10: 14. 1905.

Plants globose, caespitose, deep green, 5 cm. in diameter; tubercles somewhat elongated, often curved, 2 cm. long, terete, not at all milky, bearing bristles in the axils; spines 2 to 4, described as up to as many as 8, yellowish, bristle-like, spreading and twisted or bent, often 3 cm. long; spine-areoles small, circular, a little woolly at first; axils of tubercles bristly; flowers small, about cm. long; outer perianth-segments greenish; inner perianth-segments white, 10 mm. long, acute.

Type locality: Mexico.

Distribution: Deserts of eastern Queretaro, Mexico.

This plant was collected by Rose and Painter between Higuierillas and San Pablo, August 23, 1905 (No. 11536), and flowered in Washington on October 3, 1905. In 1913 L. Quehl of Halle sent us some flowers of this species.

Illustrations: Blühende Kakteen 3: pl. 151; Möllers Deutsche Cart. Zeit. 25: 475. f. 8, No. 6, as *Mammillaria camptotricha*.

Figure 135 is from a photograph of the plant collected by Dr. Rose in 1905.

85. *Neomammillaria eriacantha* (Link and Otto).

Mammillaria eriacantha Link and Otto in Pfeiffer, Enum. Cact. 32. 1837.

Cactus eriacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Solitary or cespitose, 10 to 15 cm. high, cylindric, 5 cm. in diameter; tubercles spiraled, in 22 rows; radial spines about 20, delicate, spreading, pubescent; central spines 2, widely spreading, stouter than the radials, also pubescent, yellowish; flowers borne in a ring above the middle of the plant, yellow, 14 mm. broad; inner perianth-segments about 14, linear, acute; stigma-lobes; fruit at first greenish white, afterwards tinged with red, short-clavate.




Fig. 135.—*Neomammillaria camptotricha*.




Fig. 136.—*Neomammillaria schiedeana*.

Type locality: Mexico.

Distribution: Central Mexico.

A plant, collected by McDowell, was seen in the collection of the Instituto Medico Nacional in the City of Mexico, but no specimen was obtained.

Mammillaria columbiana Salm-Dyck (Cact. Hort. Dyck. 1849. 99. 1850) is probably to be referred here. It is doubtless of Mexican rather than of Colombian origin.

Mammillaria eriantha (Pfeiffer, Enum. Cact. 32. 1837), referred here by Pfeiffer, was never described.

Mammillaria cylindracea De Candolle (Mém. Mus. Hist. Nat. Paris 17: 111. 1828) is referred here by Schumann and also by Pfeiffer and Otto, but the description of it would suggest a different species. Kuntze changes the name to *Cactus cylindraceus* (Rev. Gen. Pl. 1: 260. 1891). Here is also referred *Mammillaria cylindrica flavispina* (Labouret, Monogr. Cact. 88. 1853).

Illustrations: Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 25; Schelle, Handb. Kakteenk. 256. f. 178, as *Mammillaria eriacantha*.

Figure 138 is a reproduction of the first illustration above cited.

86. *Neomammillaria schiedeana* (Ehrenberg).

- Mammillaria schiedeana* Ehrenberg in Schlechtendal, Allg. Gartenz. 6: 249. 1838.
 ? *Mammillaria sericata* Lemaire, Cact. Gen. Nov. Sp. 44. 1839.
Cactus schiedianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
Mammillaria dumetorum J. A. Purpus, Monatsschr. Kakteenk. 22: 149. 1912.
 ? *Mammillaria cephalophora* Quehl, Monatsschr. Kakteenk. 24: 158. 1914. Not Salm-Dyck, 1850.

Densely cespitose, somewhat soft in texture; axils of tubercles bearing long bristle-like white hairs; tubercles green, terete; radial spines about 30, white, spreading, bristle-like, puberulent; central spines 6 to 10, spreading and appressed against the radials, a little stouter, often tinged with yellow; flowers 15 mm. long; inner perianth-segments white; filaments white; style cream-colored; stigma-lobes 4, short, obtuse.

Type locality: Near Puente de Dios, Mexico.

Distribution: Central Mexico.

The Index Kewensis refers *Mammillaria sericata* Lemaire to *M. magnimamma*.




FIG. 137.—*Neomammillaria lenta*.




FIG. 138.—*Neomammillaria eriacantha*.

Illustrations: Schumann, Gesamt. Kakteen f. 113; Blühende Kakteen 1: pl. 13; Monatsschr. Kakteenk. 8: 12; 13: 92, f. A, as *Mammillaria schiedeana*; Monatsschr. Kakteenk. 23: 89, as *Mammillaria dumetorum*; (?) Monatsschr. Kakteenk. 24: 158, as *Mammillaria cephalophora*.

Figure 136 is from a photograph of a plant collected by Dr. C. A. Purpus at San Rafael, Mexico, in 1910.

87. *Neomammillaria lasiacantha* (Engelmann).

- Mammillaria lasiacantha* Engelmann, Proc. Amer. Acad. 3: 261. 1856.
Mammillaria lasiacantha minor Engelmann, Cact. Mex. Bound. 5. 1859.
Cactus lasiacanthus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Globose, 2 to 2.5 cm. in diameter; tubercles small, their axils naked; spines 40 to 60, in more than one series, white, puberulent, 2 to 4 mm. long; flowers 12 mm. long, whitish or pink; fruit 1 to 2 cm. long; seeds blackish, pitted.

Type locality: On the Pecos in western Texas.

Distribution: Western Texas and northern Chihuahua. Reported also from Arizona, but doubtless incorrectly.

We have seen no specimens of *N. lasiacantha*, except the type, but the following species, first described as a variety of *lasiacantha*, is very common in eastern Texas and northern Mexico. Possibly the two should be united, the typical form simply representing a juvenile phase.

Illustrations: Cact. Mex. Bound. pl. 3; Schumann, Gesamtb. Kakteen 522. f. 86; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 56, A; Blanc, Cacti 70. No. 1335; West Amer. Sci. 13: 39, as *Mammillaria lasiacantha*.

88. Neomammillaria denudata (Engelmann).

Mammillaria lasiacantha denudata Engelmann, Cact. Mex. Bound. 5. 1859.

Cactus lasiacanthus denudatus Coulter, Contr. U. S. Nat. Herb. 3: 100. 1894.

Mammillaria lasiandra denudata Quehl, Monatsschr. Kakteenk. 19: 79. 1909.

Globose, 2.5 to 3.5 cm. in diameter; tubercles 5 to 6 mm. long; spines 50 to 80, glabrous or nearly so, 3 to 5 mm. long, the innermost usually much shorter; flowers and fruit from near the center but not from the axils of young tubercles; flowers 10 to 12 mm. long; perianth-segments few, about 12, oblong, obtuse, the margins white, the center light purple; stamens white; style and stigma-lobes green; fruit clavate, red, 1.5 to 2 cm. long; seeds black with basal hilum.




FIG. 139.—*Neomammillaria denudata*.



FIG. 140.—*Neomammillaria lenta*.

Type locality: Western Texas.

Distribution: Western Texas and northern Coahuila, Mexico.

The flowers open about mid-day and close at night; in one case which we recorded the flowers opened for six consecutive days.

Mammillaria rungii (Schumann, Gesamtb. Kakteen 522. 1898), an unpublished garden name, was supposed by Schumann to be referable to *M. lasiacantha denudata*.

Illustrations: Cact. Mex. Bound. pl. 4; Möllers Deutsche Gärt. Zeit. 25: 75. f. 8, No. 21, as *Mammillaria lasiacantha denudata*.

Figure 139 is from a photograph of a plant collected by Elmer Steams in 1909, which afterwards flowered in Washington.

89. Neomammillaria lenta (K. Brandegee).

Mammillaria lenta K. Brandegee, Zoe 5:194. 1904.

Described as cespitose; individuals globose to short-cylindric, almost hidden by the white delicate spines; tubercles very slender, light green; spine-areoles naked; spines about 40, very fragile; axils woolly and occasionally bearing a single bristle; flowers whitish, 7 mm. long; perianth-segments pointed; fruit red, clavate; seeds 1 mm. in diameter, dull black.

Type locality: Near Viesca, in Coahuila, Mexico.

Distribution: Coahuila, Mexico.

Illustration: Monatsschr. Kakteenk. **16**: 40, as *Mammillaria lenta*.

Figure 137 is from a photograph obtained from L. Quehl in 1921; figure 140 is from a photograph of a fruiting plant sent from Parras, Mexico, by C. A. Purpus in 1905.

90. *Neomammillaria candida* (Scheidweiler).

Mammillaria candida Scheidweiler, Bull. Acad. Sci. Brux. **5**: 496. 1838.

Mammillaria sphaerotricha Lemaire, Cact. Gen. Nov. Sp. 33. 1839.

Mammillaria humboldtii Ehrenberg, Linnaea **14**: 378. 1840.


Mammillaria sphaerotricha rosea Salm-Dyck, Cact. Hort. Dyck. 1849. 85. 1850.

Cactus humboldtii Kuntze, Rev. Gen. Pl. **1**: 260. 1891. Not Humboldt, Bompland, and Kunth, 1823.

Cactus sphaerotrachus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

Mammillaria candida rosea Salm-Dyck in Schumann, Gesamtb. Kakteen 525. 1898.

Cespitose; individual plant globose, 5 to 7 cm. in diameter, almost hidden by the white spines; radial spines numerous, radiating; central spines 8 to 12, porrect, often brownish at tip, a little stouter than the radials; axils setose; flowers 2 cm. long, rose-colored; perianth-segments serrulate towards the apex; fruit red; seeds black.



FIGS. 141 and 142.—*Neomammillaria candida*.

Type locality: Near San Luis Potosí.

Distribution: Central Mexico.

Illustrations: Monatsschr. Kakteenk. **29**: 141, as *Mammillaria candida rosea*; Hort. Belge **5**: pl. 117; Möllers Deutsche Gärt. Zeit. **25**: 475. f. 8, No. 27; Blühende Kakteen **3**: pl. 169, as *Mammillaria candida*.

Figure 141 is from a photograph of a plant obtained by Dr. Palmer near San Luis Potosí in 1905; figure 142 is from a photograph of a plant collected by C. A. Purpus from near the same locality in 1910.

91. *Neomammillaria vetula* (Martius).

Mammillaria vetula Martius, Nov. Act. Nat. Cur. **16**: 338. 1832.

Cactus vetulus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

Plant somewhat club-shaped, small, 4 to 5 cm. high; tubercles terete, light green, somewhat shining; axils of tubercles naked or sometimes with a small tuft of wool; radial spines about 25, spreading, white, bristle-like; central spines 1 to 6, stouter than the radials, brownish; flowers 12 to 15 mm. long, borne at upper part of plant; outer perianth-segments red with yellowish margins; inner perianth-segments cream-colored; filaments greenish; style green; stigma-lobes 5, white.

Type locality: San José del Oro, Hidalgo, Mexico.

Distribution: Hidalgo, Mexico.

The above description was drawn in part from a plant which flowered in Washington on November 8, 1912, and which had been sent to us by L. Buscacioni from Catania, Italy. This plant gave off numerous young ones from the axils of the tubercles, but it has died.

Mammillaria vetula major Salm-Dyck (Walpers, Repert. Bot. 2: 270. 1843) is said to be the same as *M. grandiflora* Hortus. If so, this must be different from *M. grandiflora* Otto, which we have referred to *Neolloydia conoidea*.

Illustration: Nov. Act. Nat. Cur. 16: pl. 24, as *Mammillaria vetula*.

Figure 143 is reproduced from the illustration above cited.




FIG. 143.—*Neomammillaria vetula*.




FIG. 144.—*Neomammillaria discolor*.

92. *Neomammillaria fertilis* (Hildmann).

Mammillaria fertilis Hildmann in Schumann, Gesamtb. Kakteen 503. 1898.

Cespitose, the individual plant globose to short-cylindric, dark green; tubercles arranged in 8 or 13 rows, a little woolly in their axils; radial spines 7 to 10, acicular, 6 mm. long; central spines 1 or 2, straight, stouter than the radials, 10 mm. long; flowers deep crimson, 2 cm. long; inner perianth-segments linear-lanceolate, acute.

Type locality: Mexico, but definite station not given.

Distribution: Mexico, but range unknown.

We have not seen living specimens of this plant but L. Quehl of Halle had it growing in 1913 and sent us flowers which we have used in this description.

93. *Neomammillaria decipiens* (Scheidweiler).

Mammillaria decipiens Scheidweiler, Bull. Acad. Sci. Brux. 5: 496. 1838.

*Mammillaria anacistrina** Lemaire, Cact. Gen. Nov. Sp. 39. 1839.

Mammillaria guillemianiana Lemaire, Cact. Gen. Nov. Sp. 48. 1839.

Mammillaria glochidiata inuncinata Lemaire, Cact. Gen. Nov. Sp. 102. 1839.

Cactus decipiens Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus guillemianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cactus ancistrus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Usually cespitose, deep green; tubercles soft, cylindric, about 1 cm. long, their axils bearing 2 or 3 bristles each; radial spines 7 to 9, spreading, slender, white, sometimes yellowish with brown

* Spelled *M. ancistrina* by Walpers (Repert. Bot. 2: 296. 1843.)

tips, puberulent when young; central spine x, much longer than the radials, erect or ascending, 15 to 18 mm. long, dark brown; flower-buds pinkish, acute; flower 15 mm. long, broadly funnel-shaped; inner perianth-segments nearly white or faintly tinged with pink, acute; filaments white to pinkish; stigma-lobes, white or pinkish, slender, filiform.

Type locality: Not cited.

Distribution: San Luis Potosí.

The above description is drawn from plants growing in the top of *Calibanus caespitosus*, a curious, globose, liliaceous plant of the desert of central Mexico, sent by Dr. E. Palmer from San Luis Potosí in 1905.

Schumann says that the axils of the tubercles are naked, while K. Brandegees describes them as bearing bristles as in our plant and so called for in the original description.

In some plants one or two of the upper radial spines are brown like the central spine; the flowers are delicately fragrant, remaining open during cloudy days. In cultivation this is one of the earliest species to flower; in 1918 it began to bloom early in January.

Mammillaria inuncinata (Lemaire, Cact. Gen. Nov. Sp. 39. 1839) was never described but belongs here.

Mammillaria ancistroides inuncinata Lemaire and *M. deficum* (Förster, Handb. Cact. 185. 1846), as synonyms, were referred here. *M. deficiens* Hortus (Salm-Dyck, Cact. Hort. Dyck. 1849. 7. 1850) is another name, used only as a synonym of this species.

Illustrations: Schumann, Gesamtb. Kakteen 528. f. 88; Knippel, Kakteen pl. 20; Schelle, Handb. Kakteenk. 249. f. 169; Blanc, Cacti 68. No. 100, as *Mammillaria decipiens*.

Plate XIV, figure 3, is from a plant sent to the New York Botanical Garden by Weinberg in 1903, which flowered November 14, 1911.

94. *Neomammillaria discolor* (Haworth).

Mammillaria discolor Haworth, Syn. Pl. Succ. 177. 1812.

Cactus depressus De Candolle, Cact. Hort. Monsp. 84. 1813. Not Haworth, 1812.

Cactus pseudomammillaris Salm-Dyck, Liste Pl. Gr. 1: 1. 1815.

Cactus spinii Colla, Mem. Accad. Sci. Torino 33: 133. 1826.

Mammillaria pseudomammillaris Pfeiffer, Allg. Gartenz. 3: 57. 1835.

Mammillaria discolor prolifera Pfeiffer, Enum. Cact. 28. 1837.

Mammillaria albida Haage in Pfeiffer, Enum. Cact. 28. 1837.

Mammillaria aciculata Otto in Pfeiffer, Enum. Cact. 29. 1837.

Mammillaria discolor monstrosa Monville in Lemaire, Cact. Gen. Nov. Sp. 99. 1839.

Mammillaria discolor albida Salm-Dyck, Cact. Hort. Dyck. 1844. 7. 1845.

? *Mammillaria curvispina* Otto in Dietrich, Allg. Gartenz. 14: 204. 1846.

? *Mammillaria discolor pulchella* Otto in Förster, Handb. Cact. 206. 1846.

Mammillaria curvispina parviflora A. Dietrich, Allg. Gartenz. 14: 204. 1846.

Mammillaria nitens Otto in Linke, Allg. Gartenz. 16: 331. 1848.

Mammillaria pulchella Otto in Linke, Allg. Gartenz. 16: 331. 1848.

Mammillaria discolor aciculata Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850.

Mammillaria discolor curvispina Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850.

Mammillaria discolor nitens Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850.

Mammillaria polythele aciculata Salm-Dyck, Cact. Hort. Dyck. 1849. 15. 1850.

Mammillaria pulchella nigricans Monville in Labouret, Monogr. Cact. 40. 1853.

Cactus aciculatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus discolor Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus pulchellus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Globose or somewhat depressed, often solitary, about 7 cm. in diameter; tubercles ovoid-conic, arranged in 13 to 15 spirals, their axils naked; radial spines 16 to 20, white, setaceous, widely spreading; central spines about 6, stouter than the radials, straight, at first black with white bases; flowers 15 mm. broad when fully open; inner perianth-segments linear, white, with a violet-rose stripe; fruit red, 2.5 cm. long.

Type locality: Not cited.

Distribution: Puebla, according to Schumann.

We have been unable to identify definitely this species. As there seems to be no type preserved we must rely upon the short original description and the early illustrations. The illustration of Loddiges (Bot. Cab. 17: pl. 1871) shows a plant with yellowish-brown spines and must belong elsewhere.

Mammillaria depressa was credited by mistake to De Candolle by Pfeiffer in listing the synonyms of *M. discolor* (Enum. Cact. 28. 1837).

Mammillaria confinis Haage, according to Pfeiffer (Enum. Cact. 28. 1837), appeared in "Haage, Catal. Cact. 1836" and he lists it as a synonym of *M. albida*.

Mammillaria canescens Hortus (Pfeiffer, Enum. Cact. 28. 1837) was given as a synonym of *M. discolor*. This is different from *M. canescens* Jacobi (Allg. Gartenz. 24: 89. 1856) which Schumann lists among his unknown plants. (See also Lemaire, Cact. Gen. Nov. Sp. 99. 1839.)

Mammillaria coniflora Hortus and *M. discolor coniflora* Salm-Dyck (Cact. Hort. Dyck. 1849. 11. 1850) are only names which belong here.

Mammillaria discolor fulvescens Salm-Dyck (Cact. Hort. Dyck. 1844. 7. 1845) was not formally published at the place here cited.

Mammillaria discolor breviflora (Förster, Handb. Cact. 206. 1846), although not described at the place here cited, is usually referred here.

Cactus pseudomammillaris appeared simply as a name in 1815 (Desfontaines, Tab. Bot. ed. 2. 191), and again in Pfeiffer's Enumeratio (28. 1837) as a synonym of *Mammillaria discolor prolifera*. Pfeiffer credits the name to Salm-Dyck and gives the reference to Allgemeine Gartenzeitung (3: 57. 1835), but the name appeared there under *Mammillaria* along with *spinii* and *canescens*. *M. spinii*, credited to Colla, is given by Salm-Dyck (Cact. Hort. Dyck. 1849. 11. 1850) as a synonym of *M. discolor*.

Schumann lists *Mammillaria rhodacantha* Salm-Dyck (Cact. Hort. Dyck 1849. 96. 1850) among his unknown species. *M. rhodacantha pallidior* (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845) is only a name, while *M. discolor rhodacantha* (Walpers, Repert. Bot. 2: 271. 1843), although never described, seems to be the same as *M. rhodacantha*.

Illustrations: Mém. Mus. Hist. Nat. Paris 17: pl. 2, f. 2: Ann. Inst. Roy. Hort. Fromont 2: pl. 1, f. A; Loddiges, Bot. Cab. 17: pl. 1671 (?), as *Mammillaria discolor*; Mem. Accad. Sci. Torino 33: pl. 11, as *Cactus spini*.

Figure 144 is reproduced from the first illustration cited above.



FIG. 145.—*Neomammillaria fragilis*.




FIG. 146.—*Neomammillaria elongata*.

95. *Neomammillaria fragilis* (Salm-Dyck).

Mammillaria fragilis Salm-Dyck, Cact. Hort. Dyck. 1849. 103. 1850.

Stems usually oblong or club-shaped, sprouting freely towards the top; branches globose and breaking off at the slightest touch; tubercles bright green, terete, their axils nearly naked; radial spines 12 to 14, white, naked, spreading; central spines usually wanting, especially on branches, if present 1 or 2, elongated, erect, brownish especially at tip; young spine-areoles with white wool; flowers from upper part of plant but not from center, small, lasting for several days; cream-colored with outer segments somewhat pinkish; petals broad with a mucronate tip; filaments and style pale.

Type locality: Not cited.

Distribution: Doubtless Mexico, but not known from wild plants.

Mrs. K. Brandegee, some years ago (Zoe 5: 5. 1900), called attention to the fact that this fragile little plant did not answer Pfeiffer's description of *Mammillaria gracilis* and that Salm-Dyck had suggested the very appropriate name of *M. fragilis*, which we have adopted here. The plant is known in the trade as *Mammillaria gracilis pulchella*, under which designation we received plants from Haage and Schmidt in 1921.

Illustrations: Schumann, Gesamtb. Kakteen 552. f. 90 (?); Blühende Kakteen 2: pl. 68; Monatsschr. Kakteenk. 6: 2; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 19; Gartenwelt 12: 333, as *Mammillaria gracilis*.

Figure 145 is from a photograph sent us by L. Quehl.

96. *Neomammillaria elongata* (De Candolle).

Mammillaria elongata De Candolle, Mém. Mus. Hist. Nat. Paris 17: 109. 1828.

Mammillaria subcrocea De Candolle, Mém. Mus. Hist. Nat. Paris 17: 110. 1828.

Mammillaria intertexta De Candolle, Mém. Mus. Hist. Nat. Paris 17: 110. 1828.

Mammillaria tenuis De Candolle, Mém. Mus. Hist. Nat. Paris 17: 110. 1828.

Mammillaria tennis media De Candolle, Mém. Mus. Hist. Nat. Paris 17: 110. 1828.

? *Mammillaria densa* Link and Otto, Icon. Pl. Rar. 69. 1830.

Mammillaria echinata densa Pfeiffer, Enum. Cact. 6. 1837.

? *Mammillaria stella-aurata* Martius in Zuccarini, Abh. Bayer. Akad. Wiss. München 2: 201. 1837.

? *Mammillaria minima* Reichenbach in Terschek, Suppl. Cact. Verz. 1.

Echinocactus densus Steudel, Nom. ed. 2. 1: 536. 1840.

Mammillaria stella-aurata Salm-Dyck in Walpers, Repert. Bot. 2: 272. 1843.

Mammillaria subcrocea intertexta Salm-Dyck, Cact. Hort. Dyck. 1844. 13. 1845.

Mammillaria elongata intertexta Salm-Dyck, Cact. Hort. Dyck. 1849. 12. 1850.

Mammillaria elongata subcrocea Salm-Dyck, Cact. Hort. Dyck. 1849. 12. 1850.

Mammillaria subcrocea rufescens Salm-Dyck, Cact. Hort. Dyck. 1849. 100. 1850.

Mammillaria stella-aurata gracilispina Salm-Dyck, Cact. Hort. Dyck. 1849. 101. 1850.

? *Mammillaria anguinea* Otto in Salm-Dyck, Cact. Hort. Dyck. 1849. 201. 1850.

? *Mammillaria subechinata* Salm-Dyck, Cact. Hort. Dyck. 1849. 101. 1850.

? *Mammillaria rufocrocea* Salm-Dyck, Cact. Hort. Dyck. 1849. 102. 1850.

Cactus anguineus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus densus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus elongatus Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Willdenow, 1813.

Cactus intertextus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus minimus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus stella-auratus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus subcroceus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus subechinatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus tenuis Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Mammillaria elongata tenuis Schumann, Gesamtb. Kakteen 520. 1898.

Mammillaria elongata stella-aurata Schumann, Gesamtb. Kakteen 520. 1898.

Mammillaria elongata anguinea Schumann, Gesamtb. Kakteen 521. 1898.

Mammillaria elongata rufocrocea Schumann, Gesamtb. Kakteen 521. 1898.

Densely caespitose, forming small clumps, erect, ascending or prostrate, 3 to 10 cm. long, 1 to 1.5 cm. in diameter, almost covered by a mass of interlocking spines; tubercles arranged in a few rows, usually in spirals, short, their axils naked; spines usually all radial but sometimes with 1 porrect central spine, yellow or with brown tips, more or less recurved, 8 to 12 mm. long; spine-areoles pubescent when young; flowers at the upper part of the plant, white or nearly so, 6 to 7 mm. long; perianth-segments about 12, rather broad, obtuse or sometimes apiculate.

Type locality: Mexico.

Distribution: Eastern Mexico.

Mammillaria supertexta rufa is referred to *M. elongata intertexta* by Labouret (Monogr. Cact. 68. 1853).

Mammillaria caespitosa was first listed by De Candolle (Prodr. 3: 460. 1828). It next appears in 1830 as a synonym in a list of the cacti of the Botanical Garden of Berlin. In 1837 Pfeiffer (Enum. Cact. 6) gives it as a synonym of *M. echinata densa*.

The three varieties *Mammillaria tenuis arrecta*, *M. tenuis coerulescens*, and *M. tenuis deprubescens* were garden names in the Botanical Garden at Berlin, listed by Förster (Handb. Cact. 240. 1846).

Walpers (Repert. Bot. 2: 272. 1843) records *M. intertexta rufocrocea*, but without any description.

Labouret (Monogr. Cact. 67. 1853) records the variety *M. stella-aurata minima* Salm-Dyck.

The two varieties of *Mammillaria subcrocea*, *anguinea*, and *rutila* (Walpers, Repert. Bot. 2: 272. 1843) are without descriptions.

Mammillaria elongata rufescens Salm-Dyck (Cact. Hort. Dyck. 1844. 12. 1845) was not described at the place here cited, while the variety *straminea* was a garden name (Förster, Handb. Cact. 240. 1846).

Illustrations: Schumann, Gesamtb. Kakteen 519. f. 85; Blühende Kakteen 3: pl. 174, as *Mammillaria elongata*; Schelle, Handb. Kakteenk. 247. f. 165, as *M. elongata minima*; Blanc, Cacti 72. No. 1398, as *M. minima*; Link and Otto, Icon. Pl. Rar. pl. 35, as *M. densa*; Abh. Bayer. Akad. Wiss. München 2: pl. 1. VIII. f. 5, as *M. stella-aurata*; Curtis's Bot. Mag. 65: pl. 3646; Edwards's Bot. Reg. 18: pl. 1523; De Candolle, Mém. Cact. pl. 1; Loudon, Encycl. Pl. ed. 2 and 3. 1201. f. 17359, as *M. tenuis*.

Figure 146 is from a photograph of the common form in cultivation.




FIG. 147.—*Neomammillaria oliviae*.

97. *Neomammillaria oliviae* (Orcutt).

Mammillaria oliviae Orcutt, West Amer. Sci. 12: 163. 1902.

Globose to short-cylindric, up to 10 cm. high, simple or becoming cespitose, sometimes as many as 8 together; tubercles ovoid, their axils naked; radial spines 25 to 36, snowy white or sometimes reddish brown, slender, rigid, 6 mm. long, the upper ones shorter; central spines 1 to 3, the lower one erect, rigid, white or tipped with chocolate brown; flowers about 3 cm. broad; perianth-segments lanceolate, acute, magenta, the upper part of the margins and tip with a narrow band of white; filaments deep magenta; style light pink; stigma-lobes olive-green; fruit scarlet, clavate, up to 2.5 cm. long; seeds small, black.

Type locality: West of Vail, a flag station on the Southern Pacific Railroad, near Tucson, Arizona.

Distribution: Mountains and deserts of Arizona.

Our description of the flowers is drawn from the notes and photograph of F. E. Lloyd's specimen sent us from Oro Blanco Mountains, Arizona. This is the only record we have had of this plant blooming, but fruiting plants were collected by C. R. Orcutt in 1922 (No. 802). It was first collected in considerable quantity by Mr. Orcutt, but his supply soon died out and most of the skeletons were sent to the U. S. National Herbarium, where

they are now preserved. In April 1921 Mr. Vernon Bailey rediscovered the species in Arizona and sent in a number of living specimens, but none has yet flowered. Mr. Orcutt reports that he has collected specimens which have hooked spines.

Mr. Orcutt dedicated this species to his wife, Mrs. Olivia Orcutt.

Figure 147 is from a photograph of two plants sent by Mr. Vernon Bailey from Continental, Arizona, in 1920.

98. *Neomammillaria echinaria* (De Candolle).

Mammillaria echinaria De Candolle, Mém. Mus. Hist. Nat. Paris 17: 110. 1828.

Mammillaria echinata De Candolle, Mém. Cact. 3. 1834.

Mammillaria gracilis Pfeiffer, Allg. Gartenz. 6: 275. 1838.

Cactus echinaria Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus gracilis Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Miller, 1770.

Mammillaria elongata echinata Schumann, Gesamtb. Kakteen 521. 1898.

Plants cespitose, often forming large clumps, ascending or spreading, about 1 dm. long, 1 to 1.5 cm. in diameter; tubercles short, terete, their axils naked; spines pale yellow to glassy white; radial spines about 1, spreading; central spine one, straight, acicular, about 1 cm. long; flowers and fruit not known.




FIG. 148.—*Neomammillaria echinaria*.




FIG. 149.—*Neomammillaria rekoii*.

Type locality: Mexico.

Distribution: Hidalgo, Mexico.

The above description is based on a plant collected by Dr. Rose in 1905 near Ixmiquilpan, and this we have had growing ever since.

The two varieties of *Mammillaria echinata*, *gracilior* Ehrenberg and *pallida*, published by Förster (Handb. Cact. 239. 1846), are probably only forms of the species.

The varieties of *Mammillaria gracilis* may or may not belong here. They are as follows: var. *laetevirens* Salm-Dyck (as a synonym of var. *pulchella*), var. *pulchella* Hoppfer and *virens*, all given by Förster in 1846 (Handb. Cact. 242). *Mammillaria elongata centrispina* (Förster, Handb. Cact. 240. 1846), which is only a name, may belong here.

Illustrations: Gartenflora 34: pl. 1208, f. d, e, as *Mammillaria echinata*.

Figure 148 is from a photograph of the plant collected by Dr. Rose (No. 8990), mentioned above.

99. *Neomammillaria pottsii* (Scheer).

Mammillaria pottsii Scheer in Salm-Dyck, Cact. Hort. Dyck. 1849. 104. 1850.

Mammillaria leona Poselger, Allg. Gartenz. 21: 94. 1853.

Echinocactus pottsi Poselger, Allg. Gartenz. 21: 107. 1853.

Cactus pottsii Kuntze, Rev. Gen. Pl. 1: 261. 1891.

More or less cespitose, the individual plants cylindric, 12 cm. long or more; tubercles almost hidden by the spines; radial spines about 30, white, weak, short; central spines 6 to 12, much stouter and longer, more or less ascending, grayish with brown tips; axils of tubercles woolly; flowers borne in a circle about 2 cm. below top of plant, about 1 cm. long; inner perianth-segments light purple, somewhat spreading at tip, acute; stamens pale, much shorter than the style, purplish above; stigma-lobes narrow; fruit red, clavate; seeds blackish brown, the surface deeply pitted.

Type locality: Not cited.

Distribution: In the highlands of the Rio Grande, Texas; Nuevo Leon and Coahuila to Chihuahua and Zacatecas, Mexico.

This species is widely grown in collections but the flowers are inconspicuous.

In the Engelmann Collection, now in the Missouri Botanical Garden, is a specimen labeled "*Mammillaria pottsii vera*—original coll. Dyck. Jan. 1857." This proves to be identical with the plant well known in our collections as *M. leona*. With specimens of this plant in hand Salm-Dyck's description, which heretofore we had not understood, is clearly




Fig. 150.—*Neomammillaria pottsii*.




Fig. 151.—*N. mazatlanensis*.

interpreted, except that he states that the tubercle is slightly sulcate above. From the fact that Engelmann says that his specimen is "*M. pottsii vera*" we suspect that he may have had a plant like *M. tuberculosa* mixed with it. This seems to have been Poselger's idea, for he refers the plant to *Echinocactus*, doubtless on account of this supposed groove. The plant which Poselger describes under *Echinocactus pottsonianus*, collected at Guerrero, south of the Rio Grande, is very different from Salm-Dyck's plant; his fragment, also deposited in the Missouri Botanical Garden, consists of a fruit, a few brownish seeds, and a spine-cluster, one attached to the top of a grooved tubercle, and is to be referred to *Escobaria tuberculosa*, or a related species. The specimen is too fragmentary to identify definitely. Poselger's misunderstanding of Salm-Dyck's plant left the way open for his species, *Mammillaria leona*, described shortly afterwards.

The description of the flower and fruit as given by Coulter is doubtless taken from Poselger but does not apply to the true *M. pottsii*. Our only Texas record is based on J. H. Ferriss's plant from the Big Bend of the Rio Grande, November 15, 1922.

Coryphantha pottsii occurs in C. R. Orcutt's Circular to Cactus Fanciers 1922 (unsigned and undated) to which he assigns *M. leona*.

Illustrations: Ann. Rep. Smiths. Inst. 1908: pl. 2, f. 3; Blanc, Cacti 70. No. 1359, as *Mammillaria leona*.

Figure 150 is from a photograph of a cluster of plants obtained in Zacatecas by F. E. Lloyd in 1908.

100. *Neomammillaria mazatlanensis* (Schumann).

Mammillaria mazatlanensis Schumann, Monatsschr. Kakteenk. 11: 154. 1901.

Mammillaria littoralis K. Brandegee, Kew Bull. Misc. Inf. 1908: App. 91. 1908.

Plants cespitose, often forming broad clumps with many oblong heads, 4 to 10 cm. long, about 2 cm. in diameter; tubercles terete, 3 to 4 mm. long, their axils naked; radial spines 12 to 15, setaceous, spreading, white; central spines 4 to 6,* stouter than the radials, reddish, ascending, 8 to 10 mm. long; flowers from the axils of the old tubercles but towards the top of the plant, 3 cm. long or more, red; perianth-segments oblong, spreading; stigma-lobes 8, very long and slender.

Type locality: Mazatlán.

Distribution: On the hills near the sea, about Mazatlán, Mexico.

Dr. Rose collected this plant in 1897 and again in 1910. From this last collection we still have growing plants, but these have never flowered.

Mammillaria littoralis K. Brandegee, first mentioned in 1907 (Monatsschr. Kakteenk. 17: 80), seems never to have been described by Mrs. Brandegee but was described in the Kew Bulletin as mentioned above, where it was stated to be from "California(?)." It was doubtless sent by Mrs. Brandegee from California but collected at Mazatlán.

Illustration: Monatsschr. Kakteenk. 15: 155, as *Mammillaria mazatlanensis*.

Figure 151 is from a photograph sent by L. Quehl, showing a flowering plant.

101. *Neomammillaria sphaelata* (Martius).

Mammillaria sphaelata Martius, Nov. Act. Nat. Cur. 16: 339. 1832.

Echinocactus sphaelatus Poselger, Allg. Gartenz. 21: 107. 1853.

Cactus sphaelatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Usually densely cespitose, often grayish, forming clumps 3 to 4 dm. in diameter, the individual plants cylindric, more or less elongated, often 1 to 2 dm. high; radial spines 54 to 20, usually white with black tips; central spines 3 or 4, usually black or reddish throughout, sometimes becoming white in age; axils of tubercles often bearing tufts of short hairs and occasionally a few bristles; flowers about 15 mm. long, purplish; fruit red, clavate; seeds black, the surface deeply pitted.

Type locality: Mexico, possibly in Oaxaca or Puebla; it was collected by Karwinsky.

Distribution: Puebla and Oaxaca; Schumann reports it, but doubtless erroneously, from Hidalgo (Zimapán) and Sonora (Guaymas).

Illustrations: Nov. Act. Nat. Cur. 116: pl. 25, f. 1; Monatsschr. Kakteenk. 28: 74; Grässner, Haupt-Verz. Kakteen 1914: 36, as *Mammillaria sphaelata*.

102. *Neomammillaria albicans* sp. nov.

Plants at first globose but becoming cylindric and then 10 to 20 cm. long, up to 6 cm. in diameter, often in clumps of 5 to 15; spines almost hiding the plant body and often pure white; radial spines numerous, short, stiff, widely spreading; central spines several, straight, stiff, often



FIG. 152.—*Neomammillaria albicans*.

* Sometimes one of the central spines is hooked, as is shown in plants from near the type locality collected by Señor J. G. Ortega in 1922.

brownish or blackish at tip; spine-areoles when young densely white-woolly; fruit clavate, red, 10 to 18 mm. long; seeds black with basal hilum.

Collected on Santa Cruz Island, Gulf of California, by J. N. Rose, April 16, 1911 (No. 16842, type), and by Ivan M. Johnston in 1921 (No. 3912) also on the adjacent island of San Diego by Mr. Johnston (No. 3923).

This is a very beautiful plant which grows in small clusters and is covered with nearly pure white spines. A number of plants were brought back to the New York Botanical Garden in 1911 by Dr. Rose but they have all since died. We now have living plants sent in by Mr. Johnston from two localities.

Figure 152 is from a photograph of a plant sent by Mr. Johnston to Washington from the type locality.




FIG 153.—*Neomammillaria slevinii*.

103. *Neomammillaria slevinii* sp. nov.

Plants simple, cylindric, 1 dm. high or more, 5 to 6 cm. in diameter, entirely hidden under the many closely set spines; spines at top of plant pinkish below, with brown to blackish tips, on lower part of plant bleaching white; radial spines numerous, acicular, widely spreading; central spines about 6, a little longer and stouter than the radials, slightly spreading; flowers about 2 cm. broad; outer perianth-segments with a pinkish mid-rib; inner perianth-segments white; filaments pinkish; style nearly white; stigma-lobes nearly white; fruit red, about 1 cm. long; seeds black, nearly globular, with a projection at base and a large basal hilum.

Collected by J. N. Rose, March 31, 1911 (No. 16550, type), on San Josef Island, and by Ivan M. Johnston in 1921 (No. 3943) on San Francisco Island just off the southern end of San Josef Island.

This species is related to *Neomammillaria albicans*, but it has darker spines and the spine-areoles are not densely lanate.

The plant is named for J. R. Slevin, who was in charge of the scientific expedition of the California Academy of Sciences to the Gulf of California in 1921, at which time the plant was collected.

Figure 153 is from a photograph of one of the plants collected by Mr. Johnston and sent to Washington.

104. *Neomammillaria palmeri* (Coulter).

Cactus palmeri Coulter, Contr. U. S. Nat. Herb. 3: 108. 1894.

Mammillaria dioica insularis K. Brandegee, Erythea 5: 115. 1897.

Densely cespitose; individuals small; axils densely woolly and bristly; radial spines 25 to 30, slender, white, 5 mm. long, radiating; central spines 3 to 5, stouter and longer than the radials, brownish with black tips, straight, 7 to 8 mm. long; flowers cream-colored, sometimes tinged with pink; fruit clavate, scarlet; seeds black.

Type locality: "San Benito Island." *

Distribution: San Benito Islands and possibly Guadalupe Island off the west coast of Lower California.

Plate XIV, figure 7, shows the plant, collected on the San Benito Islands, which flowered in the New York Botanical Garden, April 1, 1912.

105. *Neomammillaria uncinata* (Zuccarini).

Mammillaria uncinata Zuccarini in Pfeiffer, Enum. Cact. 34. 1837.

Mammillaria bibamata Pfeiffer, Allg. Gartenz. 6: 274. 1838.

Mammillaria depressa Scheidweiler, Bull. Acad. Sci. Brux. 5: 494. 1838.

Mammillaria uncinata biuncinata Lemaire, Cact. Gen. Nov. Sp. 96. 1839.

Mammillaria uncinata spinosior Lemaire, Cact. Gen. Nov. Sp. 96. 1839.

Mammillaria uncinata rhodacantha Hortus in Förster, Handb. Cact. ed. 2. 347. 1885.

Cactus bibamatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus depressus Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not De Candolle, 5813.

Cactus uncinatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Globose or somewhat depressed, usually half-buried in the soil, 8 to 30 cm. in diameter; tubercles lactiferous, short, obtuse; axils of old tubercles naked, of young ones lanate, forming a mass of wool at top; young spine-areoles also lanate; radial spines 4 to 6, usually white, subulate, 4 to 5 mm. long; central spines usually solitary, sometimes 2 or 3, much stouter than the radials, 8 to 12 mm. long, brown, hooked at apex; flowers small, reddish white, about 2 cm. long; inner perianth-segments linear-oblong; stigma-lobes pinkish; fruit clavate, 10 to 18 mm. long, red; seeds small, brown.

Type locality: Mexico.

Distribution: Common in central Mexico, especially in Hidalgo and San Luis Potosí. Schumann reports it from Chihuahua, as collected by Wislizenus, but we suspect that there is an error. Pfeiffer does not give a definite locality for this species but Zuccarini, who re-described the plant soon afterwards, says that Karwinsky obtained it in the mountains near Pachuca, Mexico.

This species and the following two are the only milk-bearing *Neomammillaria* which have hooked spines.

Mammillaria adunca Scheidweiler (Förster, Handb. Cact. 222. 1846), referred here as a synonym, was never described.

Illustrations: Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 19; Schumann, Gesamtb. Kakteen f. 94; Abh. Akad. Bayer. Wiss. München 2: pl. 4, f. 3; Schelle, Handb. Kakteenk. 269. f. 191, as *Mammillaria uncinata*.

106. *Neomammillaria hamata* (Lehmann).

Cactus cylindricus Ortega, Nov. Rar. Pl. 528. 1800. Not Lamarck, 1783.

Mammillaria hamata Lehmann in Pfeiffer, Enum. Cact. 34. 1837.

Cactus hamatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Stem 6 dm. long, cylindric, somewhat branched at base, described as milky; tubercles conic or a little compressed; radial spines 15 to 20, white, spreading; central spines several, brownish, stouter than the radials, one of them hooked; flowers small, probably scarlet, from near top of plant

*Although San Benito Island is given as the type locality, San Benito is really a group of three small islands. Dr. Rose found this species on two of these islands in 1911 (No. 16042).

but from axils of old tubercles; inner perianth-segments lanceolate, acute; filaments half length of perianth-segments, white; stigma-lobes 4, yellowish; fruit slender, clavate, probably red; seeds minute, brown.

Type locality: Mexico.

Distribution: Mexico, but range not known.

Schumann referred both *Cactus cylindricus* and *Mammillaria hamata* to *M. coronaria*, but the last name must be excluded from this genus. The specific name, *cylindricus*, which has been used four times in the genus *Cactus*, can not be transferred to *Neomammillaria* on account of the earlier use of this specific name by Lamarck.

Mammillaria hamata was first mentioned in the Seed Catalogue of the Hamburg Garden in 1832.




FIG. 154.—*Neomammillaria hamata*.




FIG. 155.—*Neomammillaria wildii*.

The following are usually referred as synonyms of *Mammillaria coronaria*, but probably belong here: *Mammillaria hamata brevispina* and *M. hamata principis* Salm-Dyck (Labouret, Monogr. Cact. 34. 1853) and *M. hamata longispina* Salm-Dyck (Cact. Hort. Dyck. 1844. 8. 1845). *Mammillaria principis* Monville (Labouret, Monogr. Cact. 34. 1853) was given as a synonym of the last variety here cited.

Illustration: Ortega, Nov. Rar. Pl. pl. 16, as *Cactus cylindricus*.

Figure 154 is reproduced from the illustration above cited.

107. *Neomammillaria rekoi* sp. nov.

Globular to short-cylindric, becoming 12 cm. long, 5 to 6 cm. in diameter, sometimes milky; tubercles green, terete, 8 to 10 mm. long, not very closely set, each bearing in its axil a tuft of short white wool and 1 to 8 long white bristles; radial spines spreading, about 20, white, delicately acicular, 4 to 6 mm. long; central spines 4, brown, much stouter than the radials, 10 to 15 mm. long, the

lower one sometimes strongly hooked; flowers from axils of old tubercles, near top of plant; 1.5 cm. long, deep purple; inner perianth-segments narrowly oblong, apiculate; filaments and style purplish; stigma-lobes greenish; fruit clavate, red, 12 mm. long; seeds minute, brown.

This species has been sent to us repeatedly from Oaxaca, Mexico, by Dr. B. P. Reko and it has been named in his honor; we have selected as the type his specimen of 1921, which flowered in Washington.

This is a remarkable species, being the only one we know, except the following, which has the characters of watery tubercles, a hooked spine, and brown seeds, but some plants give out a very diluted milk and have no hooked spines.

Dr. Reko sent us a single plant in April 1922, which was about 12 cm. long and short-clavate; the central spines were mostly 4, but sometimes 5, and none of them hooked. In this specimen we obtained a diluted milky juice from the upper tubercles while the lower ones are entirely devoid of milk. It flowered in April 1923 and seemed to be referable here.




FIG. 155.—*Neomammillaria rekoii*.

Figure 149 shows a plant sent by Dr. B. P. Reko from Oaxaca, Mexico, in 1919; figure 155*a* shows the plant collected by Dr. Reko in 1922, referred to above.

108. *Neomammillaria solisii* sp. nov.

Simple, globular or nearly so, 5 to 7 cm. in diameter, green or becoming purplish; tubercles 8 mm. long, terete in section, a little narrow towards the tip and thus separated above from the adjoining tubercles, their axils without wool even when quite young, and usually with 1 to many bristles; radial spines about 10 to 20, spreading, 6 to 7 mm. long, white, bristle-like; central spines 3 or 4, a little stouter than the radials, becoming brown, one of them strongly hooked (sometimes 2 cm. long).

Collected by Octavio Solís in Cerro de Buenavista de Cuellar, Guerrero, Mexico, in 1920 (No. 5) and in 1921, type, and at the same station by Professor C. Núñez in April and November 1921 (Nos. 4 and 6).

Figure 156 is from a photograph of a plant sent by Octavio Solís from Guerrero, Mexico, in 1920; figure 157 is from a photograph of a plant sent by Professor C. Núñez in 1922.

109. *Neomammillaria pygmaea* sp. nov.

Plant very small, globose to cylindric, 2 to 3 cm. in diameter; tubercles small, obtuse; radial spines about 15, white, stiff, hardly puberulent even under a lens; central spines 4, ascending, golden yellow, the lower one hooked, 5 to 6 mm. long; flowers about 1 cm. long, the outer segments tinged with red, apiculate; inner perianth-segments about 10, cream-colored; filaments greenish, much shorter than the perianth-segments; style greenish.

Collected by J. N. Rose near Cadereyta, Querétaro, Mexico, in 1905 (No. 9863). It has repeatedly flowered but was only 3 cm. high in 1921 when it died.

The species is known only from the single collection recorded above. It grows on stony hills in a very arid part of Queretaro. It is very inconspicuous and is easily over looked in the field.

110. *Neomammillaria wildii* (Dietrich).

Mammillaria wildii Dietrich, Allg. Gartenz. 4: 137. 1836.


Mammillaria wildiana Otto in Pfeiffer, Enum. Cact. 37. 1837.

Mammillaria wildiana compacta Hortus in Förster, Handb. Cact. ed. 2. 258. 1885.

Mammillaria wildiana cristata Hortus in Förster, Handb. Cact. ed. 2. 258. 1885.

Cactus wildianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cylindric to globose, cespitose at base; axils of tubercles bearing rose-colored hairs and bristles; tubercles slender, elongated, 8 to 10 mm. long, obtuse, green or somewhat rose-colored at base; young areoles tomentose; spines all pubescent; radial spines 8 to 10, 8 mm. long, setiform, white; central spines 4, yellow, one of them hooked; flowers white, 12 mm. in diameter; inner perianth-segments acuminate; stigma-lobes 4 or 5, straw-colored; fruit clavate, red.



FIGS. 156 and 157.—*Neomammillaria solisii*.

Type locality: Mexico.

Distribution: State of Hidalgo, Mexico, according to Schumann.

We have had this plant growing for a number of years, obtained from other collectors, but we do not know its natural habitat. It sprouts freely and new plants are easily started. Dr. Rose examined a specimen, labeled *Mammillaria wildii*, in the Botanical Garden at Halle in 191; we have a cluster of spines and a flower of that plant.

Mammillaria glochidiata aurea (Pfeiffer, Enum. Cact. 37. 1837), although never described, is referred usually as a synonym of this species. The two varieties of *Mammillaria wildii*, *cristata* and *compacta*, are listed but not described by Schelle (Handb. Kakteenk. 251. 1907), the latter being offered for sale by Grässner in his Kakteen for 1914 as form *cristata*.

The two varieties, *Mammillaria wildiana major* and *M. wildiana spinosior*, were given by Walpers (Repert. Bot. 2: 270. 1843) as synonyms of *M. wildiana*. The variety *monstrosa* Cels was given by Rümpler (Förster, Handb. Cact. ed. 2. 258. 1885) as a synonym of *M. wildiana cristata*.

Illustrations: Blühende Kakteen 2: pl. 64; Monatsschr. Kakteenk. 32: 103, as *Mammillaria wildii*; Grässner, Haupt-Verz. Kakteen 1912: 27, as *M. wildii cristata*.

Plate XIV, figure 8, shows a plant from the Missouri Botanical Garden which flowered in the New York Botanical Garden, April 25, 1913. Figure 155 is reproduced from the first illustration cited above.

111. *Neomammillaria seideliana* (Quehl).*Mammillaria seideliana* Quehl, Monatsschr. Kakteenk. 21: 154. 1911.

Solitary, globose, becoming cespitose, 3 to 4. cm. in diameter; tubercles purplish, their axils naked; radial spines 20 to 25, white, long and slender, ascending, puberulent; central spines yellow, 3 or 4, puberulent when young, one hooked; flowers arising from near top of plant, about 15 to 18 mm. long, creamy yellow; outer perianth-segments brownish; inner perianth-segments oblong, acute; style cream-colored, much longer than stamens; stigma-lobes 5 or 6, cream-colored, obtuse; fruit persisting in axils of tubercles, apparently for a number of years; seeds black, with thick neck at base; the hilum basal, large.

Type locality: Zacatecas, Mexico.

Distribution: Known only from the state of Zacatecas.

Collected by F. E. Lloyd in Zacatecas, Mexico, in 1908 (No. 54), who states that he found but a single specimen, though he made diligent search for others.

Although the flowers appear to come from near the top of the plant they are all from axils of old tubercles. In the single specimen examined the flowers appeared before the plant began to form new tubercles. In *Mammillaria barbata*, a closely related species, the flowers occur at both the old and new tubercles, but so far as known no other species possesses that character, although there is no good reason for not finding it in closely related species.



FIGS. 157a and 158.—*Neomammillaria seideliana*.

We have also had a plant sent us by Haage and Schmidt; it is a profuse bloomer.

Illustration: Monatsschr. Kakteenk. 21: 155, as *Mammillaria seideliana*.

Figure 157a is from a photograph of a plant sent us from Zacatecas, Mexico, by Professor Lloyd in 1908; figure 158 is from a photograph sent by L. Quehl.

112. *Neomammillaria barbata* (Engelmann).*Mammillaria barbata* Engelmann in Wislizenus, Mem. Tour North. Mex. 105. 1848.*Cactus barbatus* Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Often densely cespitose, globose, 3 to 4 cm. in diameter; radial spines 20 or more, acicular, spreading or ascending, white, sometimes with brown tips; central spines several, subulate, brown, puberulent, 1 or 2 hooked; flowers 15 mm. long; outer perianth-segments ovate to lanceolate, ciliate; inner perianth-segments erect or spreading at tip, light straw-colored or greenish, brown without, acute; filaments numerous, short, purplish; stigma-lobes 5 to 7, greenish.

Type locality: Cosihuirachi, Mexico.

Distribution: Western Chihuahua, Mexico.

This species was collected by Dr. Wislizenus in 1846 and rediscovered and collected at the type locality in 1908 by Dr. Rose, and upon this latter collection the above description is based. Schumann did not recognize the species, but thought that it might be near *Mammillaria grahamii*.

Illustrations: Cact. Mex. Bound. pl. 6, f. 9 to 12; Monatsschr. Kakteenk. 20: 181; Gartenflora 34 pl. 1208, f. a, b, c; 43 pl. 1400, as *Mammillaria barbata*.

Figure 159 is from a photograph of the specimen collected by Dr. Rose in 1908 at the type locality.

113. *Neomammillaria mercadensis* (Patoni).

Mammillaria mercadensis Patoni, Alianza Científica Universal 1: 54. 1910.

Mammillaria ocamponis Ochoterena, Bol. Dirección Estudios Biol. 2: 355. 1918.

Solitary or cespitose, small, globose; radial spines numerous, sometimes 25, widely spreading, white; central spines 4 or 5, elongated, much longer than the flowers, one of them strongly hooked at apex; flowers small, pale rose-colored; perianth-segments oblong, obtuse.

Type locality: Cerro de Mercado, Durango.

Distribution: Durango, Mexico.

We know this plant only from descriptions and illustrations.




FIG. 159.—*Neomammillaria barбата*.

Illustrations: Alianza Científica Universal 3: pl. facing 223, as *Mammillaria barбата*; Bol. Dirección Estudios Biol. 2: facing 356, as *Mammillaria ocamponis*.

Figure 160 is from a photograph of the type plant, which has the same origin as the illustrations cited above.

114. *Neomammillaria kunzeana* (Bödeker and Quehl).

Mammillaria kunzeana Bödeker and Quehl, Monatsschr. Kakteenk. 22: 177. 1912.

Mammillaria bocasana kunzeana Quehl, Monatsschr. Kakteenk. 26: 46. 1916.

Cespitose, globose or sometimes becoming cylindric, light green; tubercles cylindric, setose in their axils; radial spines about 25, white, setaceous; central spines 3 or 4, brown, puberulent, one of them hooked; flowers white or yellowish white, rose-colored on the outside, 2 cm. long; inner perianth-segments acuminate; stigma-lobes 4, whitish yellow.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

This species is dedicated to Dr. Richard Ernest Kunze (1838-1919), who was an enthusiastic student of cacti and for many years a resident of Phoenix, Arizona. He sent the plant to Germany in 1910.

Illustration: Monatsschr. Kakteenk. 22: 178, as *Mammillaria kunzeana*.

Plate XIV, figure 1, is of a plant obtained by Dr. Rose in 1912 from W. Mundt as *Mammillaria bocasana*, which flowered in the New York Botanical Garden, April 21, 1914.

115. *Neomammillaria hirsuta* (Bödeker).

Mammillaria hirsuta Bödeker, Monatsschr. Kakteenk. 29: 130. 1919.

Solitary or becoming cespitose, globose, about 6 cm. in diameter; tubercles 10 mm. long, in 8 or 13 spiraled rows, cylindric, their axils setose; spine-areoles naked; radial spines about 20, white, 10 to 15 mm. long; central spines 3 or 4, the lower one hooked; flowers small, 10 mm. long; fruit and seeds unknown.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

The plant was exhibited by de Laet at Contich, Belgium, in 1914, as sent to him by Mrs. Nichols, presumably from northern Mexico.

Illustration: Monatsschr. Kakteenk. 29: 131, as *Mammillaria hirsuta*.




FIG. 160.—*Neomammillaria mercadensis*.

FIG. 161.—*N. multihamata*.

116. *Neomammillaria multihamata* (Bödeker).

Mammillaria multihamata Bödeker, Monatsschr. Kakteenk. 25: 76. 1915.

Short-cylindric, about 5 cm. in diameter; tubercles cylindric, setose in their axils; spine-areoles white-lanate; radial spines 25, acicular, white, 8 mm. long; central spines 7 to 9, several of them hooked; flowers numerous from near top of plant, small, 1.5 cm. long; inner perianth-segments narrow, acute, spreading; seeds blackish brown.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

This plant is in the trade. A specimen was sent us in 1914 by L. Quehl, but it never flowered and soon died.

Illustration: Monatsschr. Kakteenk. 25: 77, as *Mammillaria multihamata*.

Figure 161 is reproduced from a photograph furnished by L. Quehl.

117. *Neomammillaria longicoma* sp. nov.

Cespitose, often forming broad clumps; individual specimens 3 to cm. in diameter; tubercles conic, 4 to 5 mm. long, dark green, obtuse, bearing long white hairs in their axils; radial spines 25 or more, weak and hair-like, more or less interlocking; central spines 4, 10 to 12 mm. long, brown above, a little paler below, 1 or 2 hooked; flowers from axils of upper tubercles; outer perianth-segments pinkish, darker along the center; inner perianth-segments lanceolate, acute, nearly white or sometimes tinged with rose; stamens and style much shorter than the inner perianth-segments; stigma-lobes 3, cream-colored.

The plant is common about San Luis Potosí, Mexico, where it was collected by Dr. E. Palmer in 1905 (type) and by Mrs. Irene Vera in 1912. We have had it in cultivation since

1905. It differs considerably from *Neomammillaria kunzeana*, from the same region, in its hair-like radial spines. It is perhaps nearest *M. bocasana*, but that species has single central spines.

Illustration: Ann. Rep. Smiths. Inst. 1908: pl. 4, f. 4, as *Mammillaria bocasana*.

Figure 162 is from a photograph of a plant (type) collected by Dr. E. Palmer near San Luis Potosí in 1905 and figure 165 shows a cluster of plants from the same colony.

118. *Neomammillaria bocasana* (Poselger).

Mammillaria bocasana Poselger, Allg. Gartenz. 21: 94. 1853.

Cactus bocasanus Coulter, Contr. U. S. Nat. Herb. 3: 104. 1894.

Cespitose, often forming large mounds; individual plants globose, 3 to 4 cm. in diameter, light green; tubercles slender, 6 to 8 mm. long, terete, their axils sometimes hairy or bristly; radial spines represented by numerous long white silky hairs; central spines solitary, 5 to 8 mm. long, brown, but paler at base, hooked, much shorter than the radial hairy ones; flower-buds rose-colored; flowers described as white; perianth-segments lanceolate-linear, acute, spreading; fruit "green, 4 mm. long; seeds cinnamon brown, oblique, broadly obovate, with narrowly basal hilum."




Fig 162.—*Neomammillaria longicoma*.




Fig 163.—*Neomammillaria bocasana*.

Type locality: Sierra de Bocas,* Mexico.

Distribution: Northern central Mexico, especially in San Luis Potosí.

This species has not been well understood and is usually misnamed in collections.

The two varieties of *Mammillaria bocasana*, *cristata* and *glochidiata*, are listed by Schelle (Handb. Kakteenk. 250. 1907), but not described. The former is offered for sale by Grässner in his Kakteen for 1914. We do not find that *M. bocasana splendens* Liebner and *M. bocasana sericata* Lemaire, mentioned by Quehl (Monatsschr. Kakteenk. 19 46. 1909), have ever been described.

Mammillaria schelbsei lanuginosior Hildmann (Schumann, Gesamtb. Kakteen 531. 1898) we have not seen but it may belong here.

Mammillaria bocasana splendens, credited to Schlechtendal, is offered for sale by Haage and Schmidt in their 1922 Catalogue.

Illustrations: Schelle, Handb. Kakteenk. 250. f. 170; Blanc, Cacti 67, No. 1148; West Amer. Sci. 13: 40 (these three illustrations are from the same source); Blühende Kakteen 1: pl. 35; Monatsschr. Kakteenk. 31: 103; Schumann, Gesamtb. Kakteen f. 89, as *Mammillaria bocasana*; De Laet, Cat. Gén. 28. f. 42; Schelle, Handb. Kakteenk. 251. f. 171;

*Coulter (Contr. U. S. Nat. Herb. 3: 104) states that Poselger says the plant is from Texas "auf der Seira de Bocas," but in the original place of publication he does not give the state. Bocas, however, is in San Luis Potosí.

Rev. Hort. Belg. 40: after 186; Tribune Hort. 4: pl. 139 (these four illustrations are all from the same source); Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 25; Monatsschr. Kakteenk. 29: 81, as *Mammillaria bocasana cristata*.

Plate xiv, figure 2, shows a plant, collected by S. S. Hordes in 1915, which flowered in the New York Botanical Garden, May 11, 1916. Figure 163 shows a plant received from San Luis Potosí through Mrs. Irene Vera in 1912.

119. *Neomammillaria multiformis* sp. nov.

Cespitose, forming dense clumps, sometimes 25 or more from a single root, either globose or much elongated and 3 to 6 times as long as thick; tubercles terete, 6 to 8 mm. long, their axils bearing long white bristles and white wool; radial spines 30 or more, acicular, 8 mm. long,




FIG. 164.—*Neomammillaria multiformis*.

yellow or at least becoming so, ascending; central spines 4, a little longer and stouter than radials, nearly erect, reddish in upper part, one of them strongly hooked; flowers deep purplish red, 8 to 10 mm. long, usually broader than long; inner perianth-segments oblong, acute; filaments red; fruit nearly globose, at least when dry; seeds black.

Collected by Dr. E. Palmer at Alvarez, near San Luis Potosí, Mexico, in May 1905 (No. 591, type, and No. 592).

Figure 164 is from a photograph made from Dr. Palmer's specimen just after it was received in Washington.

120. *Neomammillaria scheidweileriana* (Otto).

Mammillaria glochidiata sericata Lemaire, Cact. Gen. Nov. Sp. 40. 1839.

Mammillaria scheidweileriana Otto in Dietrich, Allg. Gartenz. 9: 179. 1841.

Mammillaria wildiana rosea Salm-Dyck, Cact. Hort. Dyck. 1849. 81. 1850.

Cactus scheidweilerianus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

*Mammillaria monancistria** Berg in Schumann, Gesamtb. Kakteen 533. 1898.

* The publication of *Mammillaria monancistria* is usually referred to Förster's Handbuch (254. 1846), but the name occurs there without description.

Cespitose, globose to cylindric, light green; tubercles setose in their axils, in 8 and 13 spirals, cylindric; spines all puberulent; radial spines 9 to 10, setaceous, white, x cm. long; central spine, 1 to 4, brown, 1 or 2 hooked; flowers rose-colored, 12 to 13 mm. long.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

The plant is known to us from description only.

121. *Neomammillaria saffordii* sp. nov.

Plants small, globose to short-cylindric, 3 to 4 cm. high, dull green, nearly hidden under the dense covering of spines; axils naked; spine-areoles when quite young slightly woolly, but early glabrate, circular; spines all puberulent under a lens when young; radial spines 12 to 14, somewhat ascending, but in age more or less curved outward, when just developing with bright red tips and white bases, later the lower part becoming yellowish; central spines single, stout, reddish, 1.5 cm. long, hooked at apex; flowers 2.5 cm. long, rose-colored; outer perianth-segments tipped by long bristles, the inner obtuse; stigma-lobes green.

This beautiful little species was collected by W. E. Safford, February 3, 1907, near Icamole, Nuevo León (No. 1250). Two plants, which were sent to Washington, flowered June 21, 1912; but they have not done well in cultivation. The plants sprout freely in cultivation and in this way we hope to distribute material to other collections. It is near *Mammillaria carretii* and was so figured by Dr. Safford, but it differs in several important respects from that species. It is named for Dr. Safford, the author of a very interesting paper, entitled *Cactaceae of Northeastern and Central Mexico* (Ann. Rep. Smiths. Inst. 1908), frequently referred to in these volumes.

Illustration: Ann. Rep. Smiths. Inst. 1908: pl. 4, f. 2, as *Mammillaria carretii*.

Figure 168 is from a photograph of the type plant.

122. *Neomammillaria schelhasii* (Pfeiffer).

Mammillaria schelhasii Pfeiffer, Allg. Gartenz. 6: 274. 1838.

Mammillaria glochidiata purpurea Scheidweiler, Bull. Acad. Sci. Brux. 5: 495. 1838.

Cactus schelhasii Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Cespitose, forming a large hemispheric mound; individual plants globose to short-cylindric, olive-green; tubercles cylindric, their axils a little woolly, but not setose; radial spines 14 to 16, setaceous, white; central spines 3, brown, one hooked at apex; flowers large, 2.2 to 2.5 cm. long, salmon or rose-colored (Nicholson says white with line of rose down each petal); fruit 5 mm. long.

Type locality: Mineral del Monte, Mexico.

Distribution: Hidalgo, Mexico.

Salm-Dyck (Cact. Hort. Dyck. 1849. 7, 81. 1850) describes the three following varieties: *sericata*, *rosea*, and *triuncinata*, some of which may belong elsewhere. Of these Schumann recognizes only the last. The first Lemaire has referred to a different species, *Mammillaria glochidiata sericata* Lemaire (Cact. Gen. Nov. Sp. 40. 1839).

Illustrations: Schelle, Handb. Kakteenk. 252. f. 172; Dict. Gard. Nicholson 4: 565. f. 37; Suppl. 518. f. 555; Förster, Handb. Cact. ed. 2. 254. f. 24 (32, in error); Rümpler, Sukkulanten 198. f. 111; Watson, Cact. Cult. 173. f. 69; ed. 3. f. 47; Knippel, Kakteen pl. 25; Blühende Kakteen 3: pl. 170; Monatsschr. Kakteenk. 30: 163, as *Mammillaria schelhasii*; Gartenflora 6: pl. 207, as *M. schelhasii sericata*.

123. *Neomammillaria glochidiata* (Martius).

Mammillaria glochidiata Martius, Nov. Act. Nat. Cur. 16: 337. 1832.

Mammillaria ancistroides Lehmann, Del. Sem. Hort. Hamb. 1832.

Cactus glochidiatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus ancistrodes Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Densely cespitose, forming clusters sometimes 15 cm. high; tubercles cylindric, green, shining, 8 to 15 mm. long, well separated from one another towards the tip, obtuse, terete; radial spines 12 to 15, widely spreading, puberulent, white, setiform, 10 to 12 mm. long; central spines 4, brownish, one of them hooked; flowers white; inner perianth-segments lanceolate, acuminate; style longer than the stamens; stigma-lobes 4 or 5, yellow; fruit clavate, scarlet, 16 mm. long; seeds black.

Type locality: Mexico.

Distribution: Southern Mexico.

Martius, who described this species, based it on a plant of Karwinsky, but did not cite a definite locality; Hemsley, however, records Karwinsky's plant as from near San Pedro Nolasco, Hidalgo, at 7,000 to 8,000 feet altitude.

As it is a high mountain species it would doubtless not remain long in cultivation. Pfeiffer refers here *Mammillaria criniformis* De Candolle (Mém. Cact. 8. pl. 4. 1834) and transfers his two varieties *rosea* and *albida* to *M. glochidiata* as variety *rosea* and *albida* (Enum. Cact. 37. 1837). *Mammillaria criniformis* must be very different, for it has only 8 to 10 radial spines and one central spine, and this yellow. The two varieties also may belong elsewhere; in fact, the variety *rosea* has been referred to *Mammillaria decipiens*,

Mammillaria ancistrata Schelhase (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845) given as a synonym of *M. ancistroides* Lemaire, is referred here by Schumann, perhaps wrongly.




FIG 165.—*Neomammillaria longicoma*.




FIG 166.—*Neomammillaria glochidiata*.

Mammillaria ancistrina Hortus (Salm-Dyck, Cact. Hort. Dyck. 1849. 10. 1850) was given as a synonym of *M. ancistroides*.

To *Mammillaria ancistroides major* (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845) was referred *M. ancistrata* as a synonym. Afterwards it was briefly described in Förster's Handbuch.

Mammillaria bergeana, a name from Hildmann's Catalogue, is referred as a synonym of *M. glochidiata* (Schumann, Gesamtb. Kakteen 532. 1898), and so also is *M. glochidiata alba* (Förster, Handb. Cact. 188. 1846).

Mammillaria ancistroides Lehmann (Delect. Sem. Hort. Hamb. 1832) is usually referred to this species but it must go elsewhere; it has setae in the axils of the tubercles, the radial spines are 6 to 8, and the hooked spine is brown at tip.

Schumann (Gesamtb. Kakteen 532. 1898) describes two varieties, *crinita* and *prolifera*. The former is based on *Mammillaria crinita* De Candolle (Mém. Mus. Hist. Nat. Paris 17: 112. 1828; *Cactus crinitus* Kuntze, Rev. Gen. Pl. 1: 260. 1891), and has the central spines straight (at least so shown in the illustration, but described as hooked), and must be excluded

from this species. *Mammillaria crinita pauciseta* De Candolle (Mém. Mus. Hist. Nat. Paris 17: 112. 1828) may be of this relationship but we do not know it.

Other varietal names have been given, such as *M. glochidiata alba* (Förster, Handb. Cact. 188. 1846).

Illustrations: Blühende Kakteen 2: pl. 82; Nov. Act. Nat. Cur. 16: pl. 23, f. 1; Abh. Bayer. Akad. Wiss. München 2: pl. 1, I. f. 4; Monatsschr. Kakteenk. 29: 141, as *Mammillaria glochidiata*. The following illustrations we have not placed: De Candolle, Mém. Cact. pl. 3; Krock, Handb. Cact. 38, as *M. crinita*; De Candolle, Mém. Cact. pl. 4, as *M. criniformis*.

Figure 166 is reproduced from the original illustration of the type as shown in the second illustration cited above.




FIG. 167.—*Neomammillaria trichacantha*. FIG. 168.—*Neomammillaria saffordii*.

124. *Neomammillaria trichacantha* (Schumann).

Mammillaria trichacantha Schumann, Gesamtb. Kakteen Nachtr. 133. 1903.

Solitary, globose to short-cylindric, small; tubercles small, clavate, 4 to 5 cm. high, slightly glaucous; radial spines x to 18, pubescent, acicular, white, 8 mm. long; central spines 2, brownish, 12 mm. long, one of them hooked; flowers red or yellow, 1.5 cm. long; inner perianth-segments lanceolate, widely spreading, acuminate; style pale green; stigma-lobes white.

Type locality: Not cited.

Distribution: Undoubtedly Mexico, but known only from cultivated plants.

The relationship of this species is somewhat uncertain. Schumann placed it next to *Mammillaria carretii* and described the flowers as red, while Quehl stated that the inner perianth-segments are pale yellow, and this is clearly shown by an unpublished study of Mrs. Gürke, made May 26, 1907, now in our possession. We have received such flowers from Quehl.

Quehl refers here *Mammillaria hamuligera* (sometimes written *M. lamuligera*) while Bödeker would keep it distinct. We have received flowers from Quehl which correspond with Mrs. Gürke's painting of *M. trichacantha*, but her plant may be different from Schumann's type, which had red flowers.

Illustrations: Schumann, Gesamtb. Kakteen Nachtr. 133. f. 33; Monatsschr. Kakteenk. 14: 45, as *Mammillaria trichacantha*.

Figure 167 is reproduced from the first illustration cited above.

125. *Neomammillaria painteri* (Rose).

Mammillaria painteri Rose in Quehl, Monatsschr. Kakteenk. 27: 22. 1917.

Mammillaria erythrosperma Bödeker, Monatsschr. Kakteenk. 28: 101. 1918.

Mammillaria erythrosperma similis De Laet in Bödeker, Monatsschr. Kakteenk. 28: 102. 1918.

Plant globose, small, 2 cm. in diameter, almost hidden by the spines; tubercles without bristles

in their axils; radial spines about 20, stiff, white, puberulent under a hand lens; central spines 4 or 5, ascending, dark brown, one hooked, puberulent; flowers 15 mm. long, greenish white, the outer segments brownish; inner perianth-segments broad, with an ovate acute tip; stamens white; stigma-lobes cream-colored.

Type locality: Near San Juan del Rio, Queretaro.

Distribution: Central Mexico.

Collected in Querétaro, Mexico, in 1905 by J. N. Rose. It has flowered repeatedly in cultivation (August 1909, June 1911, 1912, April 1915), and is nearest perhaps to *Neomammillaria kunzeana* and *N. multihamata*, but the axils of the tubercles are naked.

Illustrations: Monatsschr. Kakteenk. 27: 23, as *Mammillaria painteri*; Monatsschr. Kakteenk. 28: 103, as *M. erythrosperma* and var. *similis*.

Figure 169 is from a photograph of the type plant.



FIG. 169.—*Neomammillaria painteri*.




FIG. 170.—*Neomammillaria microcarpa*.

126. *Neomammillaria wrightii* (Engelmann).

Mammillaria wrightii Engelmann, Proc. Amer. Acad. 3: 262. 1856.

Cactus wrightii Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Depressed-globose, simple; tubercles terete, 10 to 12 mm. long, with naked axils; radial spines 8 to 15, white, spreading, acicular; central spines 1 to 3, stouter than the radials, brown to black, x or sometimes 2 or 3 hooked at apex; flowers large, 25 mm. long and as broad as long when expanded; outer segments about 13, triangular-obtuse, fimbriate; inner perianth-segments bright purple; fruit obovoid, large, 25 mm. long, purple; seeds 1.5 mm. long, black, with a narrow ventral hilum.

Type locality: Anton Chico on the Pecos east of Santa Fe, New Mexico.

Distribution: Mountains of northeastern New Mexico.

Mammillaria wrightii as described by Dr. Engelmann is complex, his original description being based on two collections, one from the upper Pecos, the type, and one from the Santa Rita Copper mines in southwestern New Mexico. This latter specimen is referable to a new species described below. There has always existed much confusion regarding *M. wrightii*, and several species have been distributed under that name. It is very rare in collections. In the National Herbarium we have only a part of the type (clusters of spines) and spines and fruit collected by J. W. Tourney at White Oaks, New Mexico, October 20, 1896. Engelmann cites a specimen in Mexico (near Lake Santa Maria) which doubtless is to be referred elsewhere.

This species was named for Charles Wright (1811-1855), who explored extensively in Texas and Cuba.

Illustrations: Cact. Mex. Bound. pl. 8, f. 1 to 8; Monatsschr. Kakteenk. 14: 9; Möllers Deutsche Gärt. Zeit. 25: 475. f. 8, No. 5; West Amer. Sci. 13: 40; Förster, Handb. Cact. ed. 2. 249. f. 23 (as f. 31, in error); Schelle, Handb. Kakteenk. 255. f. 177; Remark, Kakteenfreund 16, 17, as *M. wrightii*.

Figure 171 is a reproduction of the first illustration cited above.




FIG. 171.—*Neomammillaria wrightii*.




FIG. 172.—*Neomammillaria mainae*.

127. *Neomammillaria viridiflora* sp. nov.

Globular to short-oblong, 5 to 10 cm. long, the plant-body well hidden under the closely appressed radial spines; tubercles terete, small, naked in their axils; radial spines 20 to 30, widely spreading, white with brown tip, bristle-like, 10 to 12 mm. long; central spines much stouter than the radials, 1.5 to 2 cm. long, brown, one or more of them hooked; flowers greenish, narrowly campanulate, 1.5 cm. long; fruit globose to ovoid, 10 to 15 mm. long, purplish, very juicy; seeds minute, 1 mm. long.

Collected by C. R. Orcutt on Superior-Miami Highway, near Boundary Monument, between Pinal and Gila counties, Arizona, 4,700 feet elevation, July, 1922 (No. 608, type), and by Mrs. Ruth C. Ross near Tula Spring, south of Aravaipa, Arizona, June 1922 (No. 14).

Here perhaps are to be referred plants collected in New Mexico by O. B. Metcalfe (Nos. 797, 803, and 820) and probably that part of *Mammillaria wrightii* which came from Santa Rita. Mr. Orcutt has repeatedly written to us about this green-flowered species, which we are now able to separate very distinctly from both *M. wrightii* and *M. wilcoxii*.

Dr. Forrest Shreve has also reported a green-flowered species from Arizona which he states is common in oak-woods.

128. *Neomammillaria wilcoxii* (Tourney).

Mammillaria wilcoxii Tourney in Schumann, Gesamtb. Kakteen 545. 1898.

Solitary, almost globose, flabby in texture, 10 cm. in diameter, almost covered by a mass of interlocking spines; axils of tubercles naked; radial spines 14 to 20, widely spreading, often 15 mm. long, bristle-like, white with colored tips; central spines 1 to 3, brown, 2 cm. long, or more hooked; flowers pink to purple, large, 3 cm. long, 4 cm. broad when fully expanded; outer perianth-segments about 20, fringed with white hairs; inner perianth-segments about 40, in 2 rows.

Type locality: Arizona.

Distribution: Southeastern Arizona. It should be looked for in northern Sonora.

This species is very rare in living collections and in herbaria. When found in the field it is often associated with *Mammillaria grahamii* and *Coryphantha aggregata*, which has led to the suggestion that it might be a hybrid between these species.

The plant is named for General Timothy E. Wilcox, U. S. A., who collected extensively in Arizona, Oklahoma, Washington, and Alaska.

Illustration: Monatsschr. Kakteenk. 24: 23, as *Mammillaria wilcoxii*.

Plate XIII, figure 1, is from a photograph of a plant collected at Calabasas, Arizona, by Dr. Rose in 1908 (No. 11955).

129. *Neomammillaria mainae* (K. Brandegee).

Mammillaria mainae K. Brandegee, Zoc 5: 31. 1900.

Globose or somewhat depressed, 5 to 8 cm. broad; tubercles pale, green, naked in their axils; spines all puberulent, at least when young; radial spines about 10, widely spreading, yellowish or white except the brownish tips; central spines usually stout, yellowish except the strongly hooked tip; flowers from upper part of plant but in old axils, about 2 cm. long, with a broad open throat; outer perianth-segments with a brownish stripe, inner ones with a reddish central stripe with broad nearly white margins; acute inner perianth-segments more or less spreading; stamens purplish; style also purplish, stout, much longer than stamens; stigma-lobes 5 or 6, purplish, elongated, linear; fruit red, globose to obovate, not projecting beyond the tubercles; seeds dull black, obovate, 1 mm. long, punctate, with a narrow basal hilum.

Type locality: South of Nogales, Sonora, Mexico.

Distribution: Northern Sonora.

For a long time it was known only from material collected by Mrs. F. M. Main, near Nogales, Mexico. It has been offered in the trade under the name of *Mammillaria galeottii*, to which, according to Mrs. K. Brandegee, it is not at all related. It was observed by Rose, Standley, and Russell in two localities near Hermosillo, Sonora, Mexico, and living plants were sent to Washington, which flowered in August 1910. This is not very close to any of the other species. It was collected again in Sonora by C. R. Orcutt in 1922.

Illustration: Monatsschr. Kakteenk. 22: 19, as *Mammillaria mainae*.

Figure 172 is from a photograph of a specimen sent by Dr. Trelease from the Missouri Botanical Garden in 1910.

130. *Neomammillaria boedekeriana* (Quehl).

Mammillaria boedekeriana Quehl, Monatsschr. Kakteenk. 20: 108. 1910.

Globose to ovoid, but in collections becoming cylindric, dull green; tubercles cylindric; radial spines about 20, white; central spines 3, brownish black, one hooked; axils naked; flowers white with brownish stripes.

Type locality: Not cited.

Distribution: Doubtless Mexico, but range unknown.

This plant, which was for a long time in cultivation in Europe, has, according to Mr. Bödeker, entirely disappeared. He writes that it is a prolific bloomer and that once he had a plant with 32 flowers open at the same time. The species is named for Friederich Bödeker of Cologne, Germany. Quehl groups this species next to *Mammillaria wrightii*.

Illustration: Monatsschr. Kakteenk. 20: 109, as *Mammillaria boedekeriana*.

Figure 172a is from a photograph of a plant which had been in cultivation 14 years by Bödeker. The photograph was sent to us in 1923.






Fig. 172a.—*Neomammillaria boedekeriana*.

1



2




1. *Neomammillaria wilcoxii*, from Calabasas, Arizona.
2. *Neomammillaria gaumeri*, from Yucatan, Mexico.

131. *Neomammillaria microcarpa* (Engelmann).*Mammillaria microcarpa* Engelmann in Emory, Mil. Recon. 157. 1848.*Mammillaria grahamii* Engelmann, Proc. Amer. Acad. 3: 262. 1856.*Cactus grahamii* Kuntze, Rev. Gen. Pl. 1: 260. 1891.*Mammillaria grahamii arizonica* Quehl, Monatsschr. Kakteenk. 6: 44. 1896.*Coryphantha grahamii* Rydberg, Fl. Rocky Mountains 581. 1917.

Globose to cylindric, simple or budding either at base or near middle, often cespitose, but in small clusters, sometimes 8 cm. high; tubercles small, corky when old; axils of tubercles naked; radial spines 15 to 30, spreading, white, sometimes with dark tips, slender, rigid, glabrous, 6 to 12 mm. long; central spines 1 to 3, dark, when more than one the lower stouter, often 18 mm. long, hooked; flowers from near top of plant, 2 to 2.5 cm. long, broadly funnel-shaped; outer perianth-segments ovate, obtuse, short-ciliate; inner perianth-segments purplish, sometimes with whitish margins, obovate, acuminate; style longer than stamens, purplish; stigma-lobes 7 or 8, linear, green; fruit clavate, 2 to 2.5 cm. long, scarlet; seeds black, shining, pitted, globose, 0.8 to 1 mm. in diameter.

Type locality: "On the Gila, 3,000 to 4,000 feet above the sea."

Distribution: Southwestern Texas and Chihuahua to Arizona and Sonora; recorded from southern California and southern Utah.



FIGS. 173 and 174.—*Neomammillaria microcarpa*.

Neomammillaria microcarpa has long been a favorite in living collections under the name of *Mammillaria grahamii*, but it does not do well in cultivation and soon dies out.

This plant is generally known under the name of *Mammillaria grahamii*. The specific name must now give place to an older one, *microcarpa*. *Mammillaria microcarpa* was based on a drawing made by J. M. Stanly, the artist on W. H. Emory's famous expedition across the continent. This drawing was sent to Dr. George Engelmann by Colonel Emory, early in 1848, with the following note: "November 4, 1846, abundant." From Emory's narrative map of his journey published later, in 1848, we know that on that date his camp was on the eastern side of the Gila and only one day's trip by pack train from the mouth of the San Pedro. His camp was "in a grove of cacti of all kinds; among them being the huge pitahaya [*Carnegiea gigantea*], one of which was 50 feet high." For years we have been striving to have this plant re-collected from the type locality; in 1908 Dr. Rose made an unsuccessful attempt to reach Emory's station.

Finally, at Dr. Rose's request, Mrs. Ruth C. Ross, on June 11, 1922, visited the locality at which Emory's party was camped on November 4, 1846, where he had said that the little *Mammillaria* was abundant. The *Mammillaria* which Mrs. Ross found there, also in some abundance, was the plant which has long passed as *M. grahamii*. Mrs. Ross deserves

great credit for the enthusiasm which she has shown in visiting this remote locality and clearing up a botanical puzzle which had remained unsolved for 70 years.*

We have not seen any California or Utah plants and we suspect that the material so-named from those states may belong to the genus *Phellosperma*, which resembles this species in its hooked central spine. The plant is undoubtedly found in northern Mexico, but how far south it extends we are in doubt.

The variety *Mammillaria grahamii californica* has not been described.

Illustrations: Emory, Mil. Recon. 157. No. 3, as *Mammillaria microcarpa*; Cact. Mex. Bound. pl. 6, f. 1 to 8; Bol. Direccion de Estudios Biol. 2: f. 2; Rümpler, Sukkulenten 199. f. 112; Schelle, Handb. Kakteenk. 254. f. 176; Remark, Kakteenfreund 16, as *M. grahamii*; Cact. Journ. 1: 171, as *M. grayhamii*.

Figures 170 and 173 are from photographs of the plants collected by Dr. Rose from the northern end of the Tucson Mountains, Arizona, April 22, 1908; figure 174 is from a photograph of a plant collected by Mrs. Ross at the type locality.

CACTUS ESCHANZIERI Coulter, Contr. U. S. Nat. Herb. 3: 104. 1894.

"Depressed-globose, 3 cm. in diameter, simple; tubercles broader at base, 6 to 8 mm. long, with naked axils; spines all pubescent; radials 15 to 20, with dusky tips, the lateral 10 to 12 mm. long, the lower weaker, shorter, and curved, the upper shorter; solitary central spines reddish slender, somewhat twisted, usually hooked upwards, 15 to 25 mm. long; flowers red (?); fruit reddish (?), ovate, about 10 mm. long; seeds reddish, oblique-obovate, 1.2 mm. long, pitted, with subventral hilum."

It is stated at the original place of publication that the type collected by Eschanzier in 1901 was in the herbarium of Coulter, but it can not be found and is probably lost. Coulter says that it resembles *Cactus grahamii*, but judging from the description and its habitat it is not very near that species. It is evidently a *Neomammillaria*, possibly referable to one of the many species which have been described from San Luis Potosí.

132. *Neomammillaria milleri* sp. nov.

Globose to elongated cylindric, sometimes more than 2 dm. long and up to 8 cm. in diameter; tubercles closely set, rather thick, nearly 1 cm. long, the axils not bristly and seemingly always naked; radial spines about 20, widely spreading, 12 mm. long or less, white, with brownish tips; central spines 2 to 4, one or all hooked at apex, brown, about 2 mm. long; flowers campanulate, about 2 cm. long, the limb 2.5 cm. broad, purple to nearly pink; inner perianth-segments similar to the outer, oblong, the margins a little paler and somewhat undulate, the apex usually obtuse, often rounded, rarely acute; stamens pale purple; style white; stigma-lobes 7 to 9, linear, yellowish to cream-colored; fruit clavate, scarlet, 1.5 cm. long; seeds black.

Collected by Dr. Gerrit S. Miller jr., near Phoenix in 1921, and by Mrs. Bly near Kingman, June 29, 1921, and in 1922. It has been observed by C. R. Orcutt near Phoenix (No. 559a, type) and near Wickenburg (No. 559,) during the summer of 1922 and several fine specimens were sent in by him. He states that it has long been known as "*Mammillaria grahamii* var." and that it suggested at times *M. phellosperma*, *M. goodridgei*, and *M. grahamii*. It differs, however, from the first in its seeds, from the second in its naked axils, and from the last in its stouter habit and stronger central spines.

Figure 184a is from a photograph of the type, collected by Mr. Orcutt.

133. *Neomammillaria sheldonii* sp. nov.

Stems slender-cylindric, about 8 cm. high; axils of tubercles without setae; radial spines 12 to 15, pale with dark tips, the 3 or 4 upper ones darker, a little stouter and 1 or 2 of them subcentral, the true central erect or porrect, with upturned hook at end; outer perianth-segments ciliate; inner perianth-segments about 10, broad, acute, light purple with very pale margins; filaments and style light purple; stigma-lobes 6, green; fruit clavate, 2.5 to 3 cm. long, pale scarlet.

*Mrs. Ross's label bears this note: On upper terrace on right bank of Gila River in s. e. corner, section 15, t. 4 s. R. 16 E. (Christmas Triangle). From grove of cactus in which we believe Emory camped, Nov. 4, 1846.



M. E. Eaton del.

A. Hoen & Co. Baltimore

- | | |
|--|---|
| 1. Flowering plant of <i>Neomammillaria kunzeana</i> . | 5. Flowering plant of <i>Neomammillaria multiceps</i> . |
| 2. Flowering plant of <i>Neomammillaria bocasana</i> . | 6. Flowering plant of <i>Neomammillaria multiceps</i> . |
| 3. Flowering plant of <i>Neomammillaria decipiens</i> . | 7. Flowering plant of <i>Neomammillaria palmeri</i> . |
| 4. Top of flowering plant of <i>Neomammillaria armillata</i> . | 8. Flowering plant of <i>Neomammillaria wildii</i> . |

This plant is described chiefly from the specimens collected by Rose, Standley, and Russell, near Hermosillo, Sonora, Mexico (No. 12366, type), but it has also been collected in Sonora by C. R. Orcutt and by Charles Sheldon, for whom it is named.

The plant differs from the *Neomammillaria microcarpa* in its stouter redder spines, in its heavier and shorter central spine with the hook more uniformly turned upward, and in its flowers, which appear to be smaller.

Figure 175 shows a plant collected by Rose, Standley, and Russell, in Hermosillo in 1910 (No. 12366), which flowered in Washington.




FIG. 175.—*Neomammillaria sheldonii*.




FIG. 176.—*Neomammillaria carretii*.

134. *Neomammillaria armillata* (K. Brandegee).

Mammillaris armillata K. Brandegee, *Zoe* 5: 7. 1900.

In clusters of 3 to 12, cylindric, sometimes 30 cm. high; tubercles bluish green, somewhat angled; axils setose and slightly woolly; radial spines 9 to 15, 7 to 12 mm. long, yellowish; central spines x to 4, but usually 2, brownish, the lowest one hooked and a little longer than the others; flowers 10 to 12 mm. long, greenish to flesh-colored; stigma-lobes greenish, short; fruit red, clavate, 15 to 30 mm. long; seeds black, punctate, constricted just above the base.

Type locality: San José del Cabo.

Distribution: Southern Lower California and on islands adjacent to it.

This species is very common in southern Lower California near the coast. Dr. Rose in 1911 collected it both at the type locality (No. 16455), and at Cape San Lucas (No. 16374). Similar to this is his plant (No. 16877) from Cerralbo Island off the coast of Lower California.

Illustration: Grässner, *Haupt-Verz. Kakteen* 1914: 23, as *Mammillaria armillata*.

Plate XIV, figure 4, shows the top of a plant collected by Dr. Rose on Margarita Island, Lower California, in 1911 (No. 16302); plate XV, figure 2 shows a plant collected by Dr. Rose on Santa Maria Bay (No. 16276); figure 3 shows the top of a plant collected by Dr. Rose at San Esteban, Lower California; figure 4 shows another plant from the same island.

135. *Neomammillaria fraileana* sp. nov.

Stems elongated, cylindric, 1 to 1.5 dm. long; axils of tubercles naked or containing at most a single bristle; central spines dark brown, one of them strongly hooked; flowers rather large,

pinkish; inner perianth-segments acuminate, 2 to 2.5 cm. long, often lacerate towards the tip; filaments and style pinkish, the latter paler and much longer than the stamens; stigma-lobes 6, long and slender, rose-colored.

Collected by Dr. J. N. Rose on Pichilique Island, March 27, 1911 (No. 16508, type); on Cerralbo Island, April 19, 1911 (No. 16895); and on Catalina Island, April 16, 1911 (No. 16831).

136. *Neomammillaria swinglei* sp. nov.

Stems cylindric, 1 to 2 dm. long, 3 to 5 cm. in diameter; axils of tubercles more or less setose; radial spines rather stout for this group, spreading, dull white with dark tips; central spines 4, ascending, dark brown or black, the lowest one elongated (1 to 1.5 cm. long), hooked at apex or sometimes straight; outer perianth-segments greenish or sometimes pinkish; margins somewhat scarious; inner perianth-segments narrowly oblong, nearly white with a brown stripe down center; style pink, twice as long as the pink filaments; stigma-lobes 8, linear, pointed, green; fruit dark red, clavate, 14 to 18 mm. long; seeds 1 mm. in diameter, constricted below, black with a large elliptic basal hilum.

Common about Guaymas, Sonora; flowers and stems described from Rose's plant (No. 12568, type) and Johnston's plant (No. 3086), and the fruit and seeds from one collected by Swingle; also collected by Dr. W. S. W. Kew in 1920.

In cultivation the inodorous flowers remain open for several days (at least three).

Growing with this species (see Rose, No. 12569) were plants with all the central spines straight. This may be the plant from Guaymas which Scheer called "a very robust species of *Mammillaria sphaerica*." * Neither flowers nor fruit were seen.

137. *Neomammillaria dioica* (K. Brandegee).

Mammillaria dioica K. Brandegee, Erythea 5: 115. 1897.

Mammillaria fordii Orcutt, West Amer. Sci. 13: 49. 1902.

Either solitary or clustered, cylindric, 5 to 25 cm. high or even higher;† axils of tubercles woolly and short-setose; radial spines 11 to 22, white, the tips often brownish to black or rose-colored throughout, 5 to 7 mm. long, spreading; central spines 3 or 4, brownish, the lower one a little longer than the others and hooked; flowers borne towards top of plant, yellowish white with purplish mid-rib, 10 to 22 mm. long, incompletely dioecious; outer and inner perianth-segments usually 6 each; outer perianth-segments reddish, especially along midrib, the inner ones oblong, pale cream-, colored, notched or toothed near apex; style white; stigma-lobes 6, linear, bright yellow to brownish green; fruit scarlet, clavate, 10 to 25 mm. long; seeds black.

Type locality: West coast of Lower California.

Distribution: Southwestern California and northwestern Lower California. According to Mr. Orcutt, this plant extends east of the coastal mountains on the border of Imperial and San Diego Counties.

Although we have not seen the type of *Mammillaria fordii* we have referred it here on the advice of Mr. Orcutt, the author of this species.

Illustrations: Cact. Mex. Bound. pl. 8, f. 9 to 14, as *Mammillaria goodridgii*.

138. *Neomammillaria goodridgei* (Scheer).

Mammillaria goodridgei ‡ Scheer in Salm-Dyck, Cact. Hort. Dyck. 1849. 91. 1850.

Mammillaria goodridgii Scheer in Seemann, Bot. Herald 286. 1856.

Cactus goodridgii Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Stems clustered, erect, globose to cylindric, up to 10 cm. long, 3 to 4 cm. in diameter; axils of tubercles not setose; radial spines 12 to 15, spreading, white, sometimes with dark tips; central spines usually 1, white below, brown above, hooked; flowers perfect, rose-colored, 15 mm. long;

* Bot. Herald 286.

† In February 1922, Mr. C. R. Orcutt sent us a single plant from the Mason's Valley on the eastern side of the Coast Mountains in San Diego County, California, which was the largest solitary one we had ever seen, being more than 33 cm. long, 10 cm. in diameter, and weighed 3 lbs. 13 oz. Three small buds were produced near the middle of the plant.

‡ Given as *Mammillaria goodrichii*, in error.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering plant of *Neomammillaria bombycina*.
2. Flowering plant of *Neomammillaria armillata*.
3. Top of flowering plant of *Neomammillaria armillata*.
4. Flowering plant of *Neomammillaria armillata*.
5. Flowering plant of *Neomammillaria goodridgei*.

segments oblong, obtuse or retuse; fruit clavate, 1.5 to 2 cm. long, scarlet, naked; seeds black, punctate, with a narrow basal hilum.

Type locality: Cedros Island, off Lower California.

Distribution: Cedros Island and the adjacent mainland of Lower California.

This species was originally collected on Cedros Island, by Mr. J. Goodridge, surgeon on the Herald during its memorable trip to the western coast of the Americas. The plant, which was sent to Scheer and named by him, was sent to Prince Salm-Dyck, who described it without knowing the flowers or fruit. The name has been associated with *N. dioica*.

Several collectors have visited Cedros Island, but all failed to find *Mammillaria goodridgei* until Dr. Rose collected it in 1911 (No. 16171); he also found it on the nearby mainland at Abrejos Point (No. 16248). Recently a plant was sent in from near Mulegé by B. F. Hake.

Plate xv, figure 5, shows a plant collected by Dr. Rose at Mulegé, Lower California, in 1911, which flowered in the New York Botanical Garden, April 11, 1912.




FIG. 177.—*Neomammillaria zephyranthoides*.

139. *Neomammillaria zephyranthoides* (Scheidweiler).

Mammillaria zephyranthoides Scheidweiler, Allg. Gartenz. 9: 41. 1841.

Mammillaria fennelii Hopffer, Allg. Gartenz. 11: 3. 1843.

Cactus zephyranthodes Kuntze, Rev. Gun. Pl. 1: 261. 1891.

Depressed-globose to short-cylindric, up to 8 cm. high, 10 cm. in diameter; tubercles about 2 cm. long; radial spines 14 to 18, 8 to 10 mm. long, very slender, white; central spines 1 (sometimes 2), larger than the radials and hooked, at first purple, but in age yellowish at base; flowers 3 to 4 cm. broad with rotate limb; perianth-segments white with red stripes; fruit and seeds unknown.

Type locality: Oaxaca, altitude about 2,300 meters.

Distribution: Oaxaca, Mexico.

We have followed previous authors in referring here *Mammillaria fennelii* and Pfeiffer's illustration, based on his statement that the type plant was abnormal and much smaller than the one figured and with smaller tubercles.

The plant was in flower at Erfurt, Germany, where Dr. Rose studied it in 1912.

Illustrations: Pfeiffer, Abbild. Beschr. Cact. 2: pl. 8, as *Mammillaria zephyranthiflora*; Schelle, Handb. Kakteenk. 254. f. 175, as *Mammillaria zephyranthoides*.

Figure 177 is reproduced from the first illustration cited above.

140. *Neomammillaria carretii* (Rebut).

Mammillaria carretii Rebut in Schumann, Gesamtb. Kakteen 542. 1898.

Solitary, dull green, globose, depressed, small, 5 to 6 cm. in diameter; tubercles cylindric; axils of tubercles naked; radial spines 14, subulate, spreading, recurved, nearly clothing the plant, long, yellowish; central spine 1, slender, chestnut-brown, hooked; flowers 2.5 cm. long; inner perianth-segments white, streaked with rose; fruit and seeds unknown.




FIG. 178.—*Neomammillaria bombycina*.




FIG. 179.—*Neomammillaria occidentalis*.

Type locality: Not cited.

Distribution: Doubtless Mexico, but no definite locality known.

We have not seen this species and know it only from descriptions and illustrations.

It is related to *Neomammillaria saffordii* but radial spines are yellow, flowers white with a streak of rose, and probably larger throughout.

Illustrations: Grässner, Haupt-Verz. Kakteen 1912: 18; 1914: 24, as *Mammillaria carretii*.

Figure 176 is reproduced from a photograph sent us by L. Quehl in 1921.

141. *Neomammillaria jaliscana* sp. nov.

Cespitose, globose, 5 cm. in diameter, bright green; tubercles in 13 rows, 4 to 5 mm. high; radial spines 30 or more, at right angles to the tubercles; central spines 4 to 6, reddish brown, darker toward the tips, one of them strongly hooked; axils naked; flowers pinkish to purplish,

delicately fragrant, 1 cm. broad when fully expanded; outer segments ovate-oblong, acute or obtuse with a more or less serrulate margin; inner perianth-segments oblong, obtuse; filaments pinkish; stigma-lobes 3 or 4, white; fruit white, 8 mm. long; seeds black.

Collected by J. N. Rose at Rio Blanco, near Guadalajara, Mexico, in September 1903 (No. 858, type), by C. R. Orcutt near Guadalajara and by B. P. Reko from the same locality in 1922 (No. 4410).

Dr. Rose introduced this species into cultivation but his plants all died. It flowered with us in March '904 and again in 1923.

142. *Neomammillaria bombycina* (Quehl).

Mammillaria bombycina Quehl, Monatsschr. Kakteenk. 20: 149. 1910.

Cylindric, 15 to 20 cm. long, to 6 cm. in diameter; tubercles spiraled, obtuse; young areoles conspicuously white-woolly; radial spines numerous, acicular, widely spreading, short, 1 cm. long or less; central spines 4, elongated, a little spreading, those toward the top of plant erect, 2 cm. long, brown except at base, the lower one hooked; flowers from near top, light purple, about 1 cm. long; perianth-segments narrowly oblong; filaments and style pinkish; stigma-lobes 4, purplish.




FIG. 179a.—*Neomammillaria occidentalis*.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

We have had this plant in cultivation for a number of years. It is a very attractive plant, the top being covered by a mass of white hairs which come from the closely set young tubercles.

Mammillaria cordigera Heese resembles this species very much in its spines and form, but is described as with grooved tubercles, which would exclude it from this genus (see page 50).

Illustration: Monatsschr. Kakteenk. 20: 151, as *Mammillaria bombycina*.

Plate xv, figure 1, shows a plant received by Dr. Rose from M. de Laet in 1910 and probably from the type collection. Figure 178 is from a photograph of another plant from the same collection.

143. *Neomammillaria occidentalis* sp. nov.

Cespitose, the branches slender, cylindric, to cm. high, densely spiny; radial spines about 12, yellowish, spreading; central spines 4 or 5, reddish or brown, one of them longer and hooked; flowers small, 1 cm. long, pink; stigma-lobes 9, slender; fruit said to be red.

Collected by Dr. E. Palmer near Manzanillo, Colima, Mexico, December 1890 (No. 1053, type) and again from the same locality by Stephen E. Aguirre, American Vice-Consul-in-Charge, October 1922. Dr. Palmer's field notes say:

"A cactus quite plentiful among rocks in exposed places. Three flowers of a pink color and three red fruits were collected. The specimens of the plants collected were cut off close to the ground; they are a fair sample of plants of the average height and diameter, but in drying they shrink to three-fourths their original dimensions."

Figure 179 is from a photograph of a plant from the type collection; figure 179a is from a photograph of the plants referred to above, sent by Mr. Aguirre.

144. *Neomammillaria fasciculata* (Engelmann).

Mammillaria fasciculata Engelmann in Emory, Mil. Recon. 157. 1848.

Cactus fasciculatus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Mammillaria thornberi Orcutt, West Amer. Sci. 12: 101. 1902.

Forming clumps, often containing many plants (as many as 110 have been noted), slender-cylindric, usually 5 to 8 cm., but sometimes 30 cm. high; axils of tubercles naked; radial spines 13 to 20, slender, 5 to 7 mm. long, white, with dark brown or nearly black tips; central spine usually 1, sometimes 2 or 3, often much elongated and 18 mm. long, brownish or black, one (sometimes all) strongly hooked; flowers broadly funnel-shaped, purplish; inner perianth-segments broad, acute; fruit short-clavate, scarlet, 8 mm. long; seeds black.




Fig. 180.—*Neomammillaria fasciculata*.




Fig. 181.—*Neomammillaria longiflora*.

Type locality: Along the Gila River.

Distribution: Southern Arizona.

This plant was found by Emory, October 20, 1846, on the Gila River, 3,000 or 4,000 feet above the sea, and was afterwards described by Engelmann from the sketch made in the field; for more than 50 years afterwards the plant remained otherwise unknown. About 1902 it was rediscovered by Professor Thornber and Mr. Orcutt near Tucson. On this latter collection Mr. Orcutt based *Mammillaria thornberi*, but he afterwards referred it to *M. fasciculata*; he is now inclined to question this reduction and thinks that *M. fasciculata* may be a species of *Echinocereus*. Engelmann, however, pointed out, when he described this species, that the spines were not arranged in vertical ribs as in *Echinocereus*. While we have not been able to prove beyond doubt the identity of the two names, as there is only one plant of this habit known from southeastern Arizona, we have admitted only one species and have used for it the older name; if a second species is afterwards found it may then be necessary to revise our conclusions. The plant has been collected several times since 1902 but it is still rare.

Illustration: Emory, Mil. Recon. 157. f. 2, as *Mammillaria fasciculata*.

Figure 180 is from a photograph of a plant collected by F. E. Lloyd near Tucson in 1906.

145. *Neomammillaria nelsonii* sp. nov.

Globose, 5 cm. in diameter; tubercles numerous, small, terete, apparently not milky, 5 to 7 mm. long, their axils naked; radial spines about 15, acicular, white, 6 to 8 mm. long, spreading; central spines several, all like the radials; but one of them elongated, stouter and longer than the others, brown to black, strongly hooked, 12 to 15 mm. long; flowers unknown; fruit very slender, clavate, 3 cm. long or more, red, few-seeded; seeds globose, black, rugose, 2 mm. in diameter; hilum basal, triangular, white, depressed.

Collected by E. W. Nelson on cliffs at La Salada, Michoacán, Mexico, March 23, 1903 (No. 6932).

This plant in its form and in the color and shape of the fruit agrees with *Neomammillaria* but differs from all the species we know in its rather large rugose black seeds. It somewhat resembles *Neomammillaria zephyranthoides*.

Figure 182 shows the fruit, spine-cluster, and seed of the type.

146. *Neomammillaria longiflora* sp. nov.

Solitary or clustered, small, 3 cm. in diameter, apparently not at all milky; tubercles small, terete, not grooved on upper side, 5 to 7 mm. long, rather closely set and nearly hidden by the spines; radial spines about 30, acicular, 10 to 13 mm. long, yellow or straw-colored, somewhat spreading; central spines 4, reddish brown, much stouter than the radials, of them straight, about length of radials, 1 of them hooked at apex, twice as long as others; flowers several, even on small plants, borne near top, 2 cm. long or more, with a distinct narrow tube; perianth-segments pinkish, oblong, acute; ovary very small, ovoid, more or less sunken in the axils, thin above and perhaps opening by an operculum, the lower part with the seeds persisting for years; seed nearly globose, minutely pitted, 1 to 1.5 mm. in diameter, black with a prominent white hilum.




FIG. 182.—Fruit, spine-cluster, and seed of *N. nelsonii*.




FIG. 183.—Seed and spine-cluster of *N. longiflora*.

Collected at Santiago Papasquiario, Durango, by Dr. Edward Palmer in 1897 (No. 89).

We have repeatedly studied this curious plant during the last 25 years, but have never been able to identify it or reach a definite conclusion as to its relationship. Our material consists of a single plant split down one side, bearing several withered flowers, and two detached flowers. Recently, we were sent a photograph of a cactus from Mexico, labeled *Mammillaria* n. sp., Sierra de Cacaria S. de Ulama, which seemed to be Dr. Palmer's plant and led us to make a detailed study of it. One of the peculiarities was the absence of an exserted ovary, so conspicuous in all the *Neomammillaria*. The cut stem showed an exposed sunken ovary, and by mere chance an old fruit with ripe seeds, probably several years old, was found in the axils of one of the oldest tubercles. As described above, the seeds are very unlike those of any species of *Neomammillaria*.

Figure 181 is a reproduction of the photograph mentioned above; figure 183 shows the seed and spine-cluster of the type.

147. *Neomammillaria tacubayensis* (Fedde).

Mammillaria tacubayensis Fedde, Nov. Gen. Sp. Ind. 1905. 443. 1905.

Globose, 3 to 5 cm. in diameter; radial spines 35 to 40, white, 3 to 5 mm. long; central spines 1, black, 5 to 6 mm. long, hooked; flower 1. cm. long.

Type locality: Near Tacubaya, Mexico.

Distribution: Mexico, but range unknown.

We know the plant only from the original description and illustration.

Illustration: Gartenflora 53: 214. f. 33, as *Mammillaria stella de Tacubaya* (but legend placed under figure 32).

148. *Neomammillaria umbrina* (Ehrenberg).

Mammillaria umbrina Ehrenberg, Allg. Gartenz. 17: 287. 1849.

Cactus umbrinus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Simple or becoming cespitose, cylindric, 10 to 12.5 cm. high, dull green; tubercles conic; axils of tubercles naked; radial spines 22 to 25, spreading, white, 4 to 6 mm. long; central spines 4, 3 being 8 to 10 mm. long, one being 20 to 24 mm. long, hooked; flowers large, 2 cm. long; inner perianth-segments about 15, narrowly lanceolate, acute, purple; stamens numerous, described as connivent, white; style filiform, longer than the stamens; stigma-lobes 7, green.

Type locality: Mexico.

Distribution: Hidalgo, according to Schumann.

We know this species from description only; it is peculiar in having hooked spines and large flowers; it resembles somewhat *Neomammillaria zephyranthoides* but is undoubtedly distinct.

149. *Neomammillaria verhaertiana* (Bödeker).

Mammillaria verhaertiana Bödeker, Monatsschr. Kakteenk. 22: 152. 1912.

Solitary, short-cylindric; tubercles subconic, their axils setose; radial spines 20 or more yellowish, setaceous, 1 cm. long, glabrous; central spines 4 to 8, stouter than the radials, brown at tip, one of them hooked at apex; flowers white, 2 cm. long, appearing in a circle below top of plant; outer perianth-segments broadly lanceolate, yellowish white; anthers rose-colored; style rose; stigma-lobes 8 or 9.

Type locality: Mexico.

Distribution: Known only from the type locality.

We know the plant only from descriptions and illustrations and a few-spine-clusters sent us by L. Quehl. Bödeker placed it next to *Mammillaria spinosissima*, but unlike that species one of the central spines is hooked.

The species is named for François Verhaert.

Illustration: Monatsschr. Kakteenk. 22: 153, as *Mammillaria verhaertiana*.

150. *Neomammillaria xanthina* sp. nov.

Depressed-globose, 7 cm. high, 8 to 9 cm. broad, dull bluish green; axils of tubercles and spine-areoles densely white-woolly when young, glabrate in age; tubercles lactiferous, broader than high, the free part about mm. long, somewhat flattened dorsally, arranged in 34 spiral




FIG. 154.—*Neomammillaria xanthina*.

rows; spine-areole circular, small; radial spines 10 to 12, spreading, acicular, white, 4 mm. long or less; central spines 2, stouter, but not much longer than the radials, somewhat brownish, more or less erect; flowers from the top of the plant but in the axils of old tubercles, the tube not exerted and the limb appressed against the adjacent tubercles; perianth rotate, 16 mm. broad, its segments, stamens, and style pale lemon-yellow; outer perianth-segments oblong, obtuse with ciliate margins, the inner a little longer than the outer, usually entire, oblong, usually retuse at apex, sometimes apiculate.

Sent by B. P. Reko (No. 4401) but collected by A. Groeschner from the vicinity of Monte Mercado, Durango, Mexico, in 1922 and flowered in Washington in May 1923.

Figure 184 is from a photograph of the type specimen.

LITTLE-KNOWN SPECIES PROBABLY OF THIS GENUS.

MAMMILLARIA ALPINA Martius in Salm-Dyck, Cact. Hort. Dyck. 1849. 79. 1850.

Cactus alpinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

This plant has not been identified. Its large flowers, 2.5 cm. broad, suggest a species of *Coryphantha*. It was collected by Karwinsky in the state of Oaxaca.

MAMMILLARIA BELLATULA Förster, Allg. Gartenz. 15: 51. 1847.

Cactus bellatulus Kuntze, Rev. Gen. Pl. 1: 259. 1891.

Spherical, somewhat compressed, bright green; tubercles broadly cone-shaped, 4 mm. long, their axils naked; spine-areoles white-woolly when young; radial spines 12 to 16, whitish, bristle-like, spreading, 6 to 8 mm. long; central spines 2, straight, one pointing downward, the other upward, 12 to 16 mm. long, at first almost black, grayish brown in age; flowers and fruit unknown.

This species is said to have been grown from Brazilian seed; if this were true it would exclude it from this genus and for this reason Schumann questioned whether it might not be an *Echinocactus*. Judging from the description we believe that it is closely related to *Neomammillaria elegans* and is probably of Mexican origin.

MAMMILLARIA BERGII Miquel, Comment. Phytogr. 104. 1840.

Simple, subglobose, glaucous green; tubercles somewhat 4-angled at base, nearly terete above, woolly in the axils; spine-areoles woolly when young, becoming naked; spines 4, spreading, the uppermost one largest.

This plant is from Mexico.

MAMMILLARIA CAESPITITIA De Candolle, Mém. Mus. Hist. Nat. Paris 17: 112. 1828.

Mammillaria nitida Scheidweiler, Allg. Gartenz. 9: 42. 1841.

Cactus caespitilius Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Densely cespitose, the clump 10 cm. in diameter; joints globose, 2.5 cm. in diameter; tubercles small, ovate; spines straight, rigid, when young whitish yellow, in age gray; radial spines 9 or 10; central spines 1 or 2, longer than the radials, erect; flowers and fruit unknown.

Both Pfeiffer and Schumann overlooked this species and it is doubtful if it can ever be identified. The plant was collected by Thomas Coulter in Mexico.

MAMMILLARIA CONICA Haworth, Suppl. Pl. Succ. 71. 1819.

Tubercles large, conic; spines less than 10, all radial, red but paler at base; flowers and fruit unknown.

Neither Pfeiffer nor Schumann knew this species or its origin. The Index Kewensis refers it to South America. If from that region it must be a species of *Discocactus*, near *D. placentiformis*.

MAMMILLARIA DIACENTRA Jacobi, Allg. Gartenz. 24: 91. 1856.

Globose, about 7 cm. in diameter; tubercles milky, rhomboid at base, not setose in their axils; radial spines 5 or 6, white, with blackish tips; central spines 2, stouter and longer than the radials, grayish, with blackish tips, the lower centrals 2.5 cm. long or more; flowers small, reddish; style rose-colored: stigma-lobes 6.

This species was unknown to Schumann, and we are unable to group it; its origin is not recorded.

MAMMILLARIA FLAVESCENS Haworth, Suppl. Pl. Succ. 71. 1819.

Cactus mammillaris lanuginosus De Candolle, Pl. Succ. 111. 1799.

Cactus flavescens De Candolle, Cat. Hort. Monsp. 83. 1813.

Mammillaria straminea Haworth, Suppl. Pl. Succ. 71. 1819.

Cactus stramineus Sprengel, Syst. 2: 494. 1825, as to name.

Mammillaria simplex flavescens Schumann, Gesamtb. Kakteen 573. 1898.

This plant was first described in 1799 by De Candolle as "var. β " of *Cactus mammillaris* or *Cactus mammillaris lanuginosus* (Pl. Succ. pl. 111); at this time he referred to it certain citations of Plumier and Hermann which we now know belong to *Neomammillaria prolifera* and *N. mammillaris* respectively. This variety was raised to specific rank by De Candolle in 1813 as *Cactus flavescens* (Cact. Hort. Monsp. 83). From the more detailed description then given it is clear that *Cactus flavescens* can not be referred to either *N. prolifera* or *N. mammillaris*. It was transferred to the genus *Mammillaria* by Haworth in 1819, but he added little information except the statement that it had been in cultivation in the Chelsea Garden before 1811.

The question has been raised whether this plant is really West Indian. It is true that De Candolle does not state its origin, but it would be indicated that he believed that it was West Indian by his treating it as a variety of the common West Indian species and by his referring to it several West Indian descriptions when he later published it as a species. Pfeiffer states that it is tropical American. As *Neomammillaria mammillaris* is the only species known from South America it could not have come from that continent, and at that time no *Mammillaria* had been discovered in the United States or Mexico. Förster in 1846 says that it is West Indian, and this was Schumann's conclusion.

MAMMILLARIA FLAVICOMA Hortus in Förster, Handb. Cact. ed. 2. 298. 1885.

This species was described from garden plants of unknown origin. Schumann does not mention it in his monograph and it has remained unknown.

MAMMILLARIA GRISEA Salm-Dyck, Cact. Hort. Dyck. 1849. 110. 1850.

Cactus griseus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Stout, short-cylindric, 10 to 12.5 cm. high, 7.5 cm. in diameter; tubercles glaucous-green, somewhat 4-angled, their axils woolly and setose; radial spines 10 to 12, spreading, short, rigid, white; central spines 4 to 6, white, with brown or blackish tips, on greenhouse plants 10 to 15 mm. long, but on wild plants 5 cm. long or more; flower and fruit unknown.

This is perhaps different from *Mammillaria grisea* Galeotti (Förster, Handb. Cact. 219. 1846), which was never described.

MAMMILLARIA HEINEI Ehrenberg, Bot. Zeit. 2: 833. 1844.

Cactus heinei Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Schumann thought that this name was referable to *Mammillaria umbrina* but we have not been able to satisfy ourselves that the two are the same.

Much confusion is found in the spelling of the name; it sometimes appears as *M. haynii* and *M. haynei*. Salm-Dyck transfers two species of Ehrenberg to varieties of *M. haynii* but both are unknown to us. These varieties are as follows: var. *viridula* Salm-Dyck (Cact. Hort. Dyck. 1849. 10. 1850; *M. viridula* Ehrenberg, Allg. Gartenz. 16: 267. 1848),

and var. *minima* Salm-Dyck (Cact. Hort. Dyck 1849. 10. 1850; *M. digitalis* Ehrenberg, Allg. Gartenz. 16: 267. 1848).

MAMMILLARIA HELICTERES De Candolle, Mém. Mus. Hist. Nat. Paris 17: 31. pl. 5. 1828.

This name was based on Mociño and Sessé's drawing of a Mexican plant, which has never since been definitely identified. It was called by them *Cactus helicteres* (De Candolle, Prodr. 3: 460. 1828), but it was renamed *Mammillaria convoluta* by St. Lager (Ann. Soc. Bot. Lyon 7: 130. 1880). The published drawing indicates that the plant is of this genus.

MAMMILLARIA HEXACANTHA Salm-Dyck, Hort. Dyck. 344. 1834.

Cactus hexacanthus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Solitary, short-cylindric; tubercles somewhat compressed, light green; areoles ovate to oblong when young, white-tomentose, glabrate in age; radial spines 25 to 30, white, 4 mm. long; central spines 6, stouter than the radials, brown, the 4 lateral ones 8 mm. long, the uppermost ones a little longer, the lowermost ones 18 mm. long, somewhat deflexed; flowers and fruit unknown.

This plant, which is of Mexican origin, is unknown to us except from description; Schumann referred it to *Mammillaria coronaria*, but it has nothing to do with that plant.

MAMMILLARIA IRREGULARIS De Candolle, Mém. Mus. Hist. Nat. Paris 17: 111. 1828.

Cactus irregularis Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cespitose, 5 cm. high, with a subtuberos base; joints ovoid, 2.5 cm. in diameter; spines all radial, 20 to 25, spreading or somewhat reflexed; flowers and fruit unknown.

This plant was collected by T. Coulter (No. 31). It has never been re-identified. It was grown at the Botanical Garden at Geneva, Switzerland, at the time the description was published but, unfortunately, no specimens were preserved; other types based on Coulter's plants are similarly lost and can never be certainly identified.

MAMMILLARIA JOOSSENSIANA Quehl, Monatsschr. Kakteenk. 18: 95. 1908.

Simple, globose to cylindric, up to 5 cm. high, 3 cm. in diameter, pale green, slightly depressed at apex; young areoles white-woolly; radial spines 20, slender-subulate, straight, white, 12 mm. long; central spines 4, stouter than the radials, 15 mm. long or more, one of them often hooked; flowers small, yellow.

We know this plant, which is a native of Mexico, only from description and two small plants sent us by Frantz de Laet in 1922. Quehl places it in Schumann's classification just after *M. amoena*, although one of the central spines is hooked.

MAMMILLARIA LESAUNIERI Rebut in Schumann, Gesamtb. Kakteen 533. 1898.

Simple, globose, or a little longer than broad; tubercles conic, their axils naked; radial spines 11 to 13, slender, subulate, straight, white, 6 to 8 mm. long; central spines solitary, very short (5 mm. long or less), brownish, erect; flowers reddish, 2.5 cm. long.

Type locality: Described from cultivated plants.

Distribution: Supposed to be Mexico proper or Lower California.

This species is supposed to have the habit of *Mammillaria heyderi*.

Here probably belongs *Mammillaria lassonneriei* Rebut (Monatsschr. Kakteenk. 7: 29. 1897), a garden name of which we have found no accompanying description. The dealer, Grässner, in his Catalogue of Cacti for 1912 (p. 21) and 1914 (p. 33) has illustrated *M. lassaunieri*.

MAMMILLARIA LEUCOCENTRA Berg, Allg. Gartenz. 8: 130. 1840.

Cactus leucocentrus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Ovoid, about 10 cm. high; tubercles ovoid, their axils very white-woolly; young spine-areoles white-tomentose at first, becoming naked; radial spines spreading, numerous, setose, white; central spines 4 to 6, stouter and longer than the radials, white throughout or with black.

Recorded from Oaxaca, but not identified.

MAMMILLARIA LORICATA Martius in Pfeiffer, Enum. Cact. 13. 1837.

Echinocactus loricatus Poselger, Allg. Gartenz. 21: 107. 1853.

Coryphantha loricata Lemaire, Cactées 35. 1868.

Cactus loricatus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Solitary, simple, globose, 4 to 5 cm. in diameter, glaucous-green; tubercles short-ovate, 4-angled at base; radial spines 12, spreading, rigid, yellow, 6 to 8 mm. long; central spines 2, stouter than the radials, 8 to 10 mm. long, black at tip, the upper one straight, the lower one curved; flowers and fruit not described.

This plant is recorded as of Mexican origin, but we have found no description of it subsequent to the original and it may never be identified. Förster referred it to *Mammillaria polythele*, but Schumann did not know it.

Mammillaria heteracantha was referred here as a synonym by Pfeiffer (Enum. Cact. 13. 1837). This plant was mentioned by Martius (Verz. Konig. Bot. Gard. München 127. 1829), but so far as we can learn was never described.

MAMMILLARIA MONOCENTRA Jacobi, Allg. Gartenz. 24: 90. 1856.

Depressed-globose, up to 12 cm. high, about 8 cm. in diameter, umbilicate at apex; tubercles milky, somewhat rhomboid at base, a little flattened, not setose in their axils; radial spines 6, white with black tips, a little spreading; central spine solitary, stouter and longer than the radials, about 2.5 cm. long; flowers rather large, rose-colored; style rose-colored; stigma-lobes 6, reddish yellow.

Jacobi referred this plant, presumably of Mexican origin, to the group *Angulosae-tetragonae* of Salm-Dyck.

Schumann placed it among his list of little-known species; we know it from description only.

MAMMILLARIA NERVOSA CRISTATA Journ. Hort. Home Farm. III. 60: (?) 7. 1910.

We know this plant only from a brief description and an illustration on pages 7 and 8 of the journal here cited:

Mammillaria nervosus cristatus * grows in convoluted sinuous masses like a great brain-mass. The growths are covered with spiny mamillae (whence the name of the genus) and are of a dull olive-brownish hue. It, too, is Mexican."

We are not able to place this plant; it resembles the cristate form sometimes assumed by *Pediocactus simpsonii* and also resembles *Mammillaria bicolor* as shown by the illustration under *M. daedalea*.

Illustration: Journ. Hort. Home Farm. III. 60: 8 (or 7).

MAMMILLARIA NICHOLSONI Journ. Hort. Home Farm. III. 60: 7. 1910.

We know this species only from the illustration referred to below and the following brief note taken from the place of publication:

"*Mammillaria nicholsoni* resembles several of the Echinocactuses in external form. It was named we believe in honor of the late Mr. George Nicholson and came to Kew from the Swanley Collection. All our illustrations were secured at Kew where the collection is well cultivated. *M. nicholsoni* forms spherical masses with the typical protuberances or tubercles, these being tipped with sharp spines."

It is doubtless of Mexican origin.

Illustration: Journ. Hort. Home Farm. III. 60: 9.

MAMMILLARIA NUDA De Candolle, Prodr. 3: 460. 1828.

This is based on *Cactus nudus* (Mociño and Sessé, Pl. Mex. Sc. ined.), but has never been subsequently identified. It was also taken up by Otto Kuntze as *Cactus nudus* (Rev. Gen.

* This is the original spelling.

Pl. 1: 261. 1891). The original description was based on a drawing and calls for a cylindric, unbranched plant, bearing unarmed tubercles and rose-colored flowers.

MAMMILLARIA PICTA Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Cactus pictus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Globose to ovoid, dull green; tubercles cylindric, somewhat oblique, obtuse, their axils setose; spines pubescent; radial spines 12, yellowish at base, white near middle, above dark purple; central spines 1 (rarely 2), erect; flowers greenish white; stigma-lobes 3.

This species is known from the description only. It was recorded as from Mexico.

MAMMILLARIA PLECASTIGMA Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria plecastigma major Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Mammillaria plecastigma minor Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 27. 1858.

Cactus plecastigma Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Proliferous, the joints cylindric; tubercles cylindric, the apex oblique and rounded, with setae in their axils; radial spines 16 to 20, setaceous, white; central spines 3 or 4, at first yellow, becoming brown, one hooked at apex; flowers and fruit unknown.

Presumably of this genus and recorded as of Mexican origin; but not identified since it was described.

MAMMILLARIA PLINTHIMORPHA Jacobi, Allg. Gartenz. 24: 92. 1856.

Cespitose, forming clumps 15 cm. in diameter or more; joints globose; tubercles 4-angled, obtuse, bearing yellowish white wool in their axils; spines 4, subulate, somewhat angled, flesh-colored with blackish tips, the upper one the longest and sometimes more than 2.5 cm. long; flowers not known.

This plant was collected by Galeotti in Mexico in 1847; we do not know it and it was listed by Schumann among his little-known species.

MAMMILLARIA PULCHRA Haworth in Edwards's Bot. Reg. 16: pl. 1329. 1830.

Cactus pulcher Kuntze, Rev. Gen. Pl. 1: 261. 1891.

This species, which has yellow spines and dark-red flowers, was referred by Schumann, doubtfully, to *Mammillaria centricirrha*, and by Pfeiffer with doubt to *M. tentaculata*.

MAMMILLARIA RUTILA Zuccarini in Pfeiffer, Enum. Cact. 29. 1837.

Cactus rutilus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Simple, globose; axils of tubercles nearly naked; tubercles 1 cm. long, conic, dull green; areoles when young tomentose; radial spines 14 to 16, setiform, the upper ones smaller, 4 to 8 mm. long; central spines 4 to 6, spreading, rigid, 8 to 12 mm. long, curved, reddish brown, the lower one longest.

Type locality: Mexico.

This name is referred by Schumann to *M. coronaria*.

M. rutila pallidior Salm-Dyck (Cact. Hort. Dyck. 1849. 11. 1850) was never described, while *M. eugenia* (Salm-Dyck, Cact. Hort. Dyck. 1849. 11. 1850) is given as a synonym of *M. rutila*.

M. rutila octospina Scheidweiler (Bull. Acad. Sci. Brux. 6: 91. 1839) is briefly described.

MAMMILLARIA SAXATILIS Scheer, Bot. Herald 286. 1856.

Cactus saxatilis Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Plant small; spines brownish to straw-colored.

Only two plants were collected, somewhere in Mexico, by Potts and sent to Scheer; the flowers and fruit were not described. The species may never be identified.

MAMMILLARIA SCHMERWITZII Haage in Förster, Handb. Cact. ed. 2. 270. 1885.

Depressed-globose, 10 cm. in diameter, grassy green; radial spines 10 to 25, yellow, 4 to 5 mm. long; central spines 4 or 5, dark brown, 15 mm. long; flowers red.

This plant, recorded as of Mexican origin, was at one time offered for sale by A. Blanc and Company; we know it only from description and are unable to identify it.

MAMMILLARIA SEEMANNII Scheer in Seemann, Bot. Herald 288. 1856.

Cactus seemannii Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Hemispheric, stout, 10 cm. in diameter, 7.5 cm. high; tubercles somewhat ovoid, elongated, greenish, minutely punctate, their axils soon white-woolly; radial spines 11 to 13, nearly equal, less than 6 mm. long; central spines 1, shorter than the radials, subulate, straight, erect, blackish purple, becoming white.

This plant was sent to F. Scheer in 1850, who states that it came from Sonora or Durango. It is incompletely described and can not be identified. It may be a species of *Coryphantha*.

MAMMILLARIA SORORIA Meinshausen, Wöchenschr. Gärt. Pflanz. 1: 28. 1858.

Cactus sororius Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Depressed-globose, to 6 cm. high, 7.5 to 10 cm. in diameter, milky; tubercles angled, 12 mm. long, naked in their axils; radial spines 6, 4 mm. long; central spines 1, erect, stouter than the radials, blackish at apex; flowers greenish purple; stigma-lobes 4.

Recorded as of Mexican origin but otherwise unknown.

MAMMILLARIA SPINAUREA Salm-Dyck, Allg. Gartenz. 18: 59. 1850.

Cactus spinaureus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

Globose or becoming depressed; tubercles light green, somewhat 4-angled, gibbous at base, obtuse and oblique at apex, their axils woolly; radial spines about 12, slender, rigid, spreading; central spines 5 or 6, twice as long and stouter than the radials, recurved or reflexed, yellow.

The above was sent by John Potts from Chihuahua in 1850; Scheer thought that it might have been collected in Durango or Sonora. We have not been able to identify it.

MAMMILLARIA SUAVEOLENS Rümpler in Förster, Handb. Cact. ed. 2. 297. 1885.

About 4 cm. high; radial spines 13 to 15; central spines 4, brown; flowers and fruit unknown.

The above is unidentifiable from the brief description. It was grown in Germany from Mexican seed.

MAMMILLARIA TROHARTII Hildmann in Schumann, Gesamtb. Kakteen 586. 1898.

Simple or proliferous and densely cespitose, globose or somewhat depressed, glaucous-green, small (6 cm. in diameter); axils naked; areoles at first woolly, afterwards naked; tubercles very small, scarcely angled; radial spines 5, with brown tips; central spines solitary, dark brown, subulate; flowers and fruit unknown.

M. trohartii is of Mexican origin.

MAMMILLARIA UNISETA Quehl, Monatsschr. Kakteenk. 14: 128. 1904.

Solitary, globose, about cm. in diameter, somewhat depressed at apex; tubercles dark green, 4-angled; spines 6, about 3 mm. long, at first black, changing to gray; flowers and fruit unknown.

This plant was described from a specimen in the Botanical Garden at Halle of unknown origin, but doubtless from Mexico.

MAMMILLARIA VIPERINA J. A. Purpus, Monatsschr. Kakteenk. 22: 148. 1912.

Cespitose, decumbent, cylindric, 1.5 to 2 cm. in diameter; tubercles very short, sometimes nearly globular; spines numerous, mm. long, whitish brown to brownish black; flowers and fruit unknown.

This plant came from Rio de Zapotitlán, Puebla; we know it only from description and the very characteristic published illustration. Quehl, who had seen it, said that it was a form of *Mammillaria elongata*. We believe that it is near *M. sphacelata* and perhaps

a distinct species. The plant figured by Grässner (Haupt-Verz. Kakteen 38. 1914) shows nearly upright branches.

Illustration: Monatsschr. Kakteenk. 23: 21, as *Mammillaria viperina*.

MAMMILLARIA ZEYERIANA Haage jr. in Schumann, Gesamtb. Kakteen 574. 1898.

Simple, hemispheric to short-cylindric, up to 10 cm. high, pale glaucous-green; tubercles in 13 or 21 spirals, terete, 10 to 12 mm. long, their axils naked; spine-areoles elliptic, 3 mm. long; radial spines 10, white; central spines 4, the uppermost one curved, 15 mm. long, brownish; flowers and fruit unknown.

Described from Mexican plants; supposed to be of Mexican origin.

PLANTS KNOWN BY NAME ONLY.

Mammillaria acicularis Lemaire (Cact. Gen. Nov. Sp. 34. 1839) was described without the flowers, fruit, or native country being known and has not been identified; here belongs *Cactus acicularis* (Kuntze, Rev. Gen. Pl. 1: 261. 1891), but *C. acicularis* (Kuntze, Rev. Gen. Pl. 1: 260. 1891) based on some name of Lehmann we have not been able to find.

Mammillaria aulacantha, referred by Schumann and the Index Kewensis to De Candolle's Revision (Mém. Mus. Hist. Nat. Paris 1: 113. 1828), is not to be found at the place cited by them; here probably belongs *Cactus aulacanthus* Kuntze (Rev. Gen. Pl. 1: 260. 1891).

Mammillaria beneckeii Ehrenberg (Förster, Handb. Cact. 210. 1846; *Cactus beneckeii* Kuntze, Rev. Gen. Pl. 1: 260. 1891) was referred to *M. coronaria* by Schumann.

Mammillaria brandi is described in Blanc, Hints on Cacti, p. 67, as "a rare Mexican sort, with very long straw-colored spines deflecting from the plant. Flowers cream-colored and very fragrant."

Mammillaria centa is mentioned by C. A. Purpus in a short article in Die Gartenwelt (9: 249. 1905).

Mammillaria chrysantha is listed by De Candolle (Prodr. 3: 460. 1828) among species little known but not described. It is said to have been in the Berlin Botanic Garden.

Mammillaria circumtexta Martius (Hort. Reg. Monac. 127. 1829) seems never to have been described.

Mammillaria hochderferi is mentioned by C. A. Purpus in a short article in Die Gartenwelt (9: 249. 1905).

Mammillaria multiradiata (Martius, Hort. Reg. Monac. 127. 1829) is only a name.

Mammillaria nigra Ehrenberg (Allg. Gartenz. 17: 287. 1849) was referred to *M. coronaria* by Schumann; *Cactus niger* Kuntze (Rev. Gen. Pl. 1: 261. 1891) is a synonym of it.

Mammillaria parmentieri Link and Otto (Verh. Ver. Beförd. Gartenb. 6: 429. 1830), without description, was doubtfully referred to *M. flavescens*. It was supposed, however, to have come from Mexico.

The following species, briefly described by F. Schlumberger (Rev. Hort. IV. 5: 404. 1856), we do not know, nor do we find them mentioned elsewhere:

Mammillaria albiseta, with flowers like those of *M. spinosissima*.

Mammillaria bocasiana, with clear yellow flowers.

Mammillaria cunendstiana, with flowers like those of *M. clillifera*.

Mammillaria decholaria, with very small red flowers.

Mammillaria klenneirii, with rose-colored flowers.

Mammillaria roematactina, with abundant small rose-red flowers.

Mammillaria saluciana, flowers 1.5 cm. long and of the same diameter, flesh-colored.

The following names, without descriptions, appear in Förster's Handbuch (254, 255, 1846). Some of the names have been used subsequently, but so far as our observation goes they

are all still *nomen nudum*. *Mammillaria asteriflora* Cels, *M. binops* Haage, *M. cantera* Haage, *M. citrina* Scheidweiler, *M. contacta* Wendland, *M. coryphides* Forbes, *M. crinigera* Otto, *M. daedalea viridis* Fennel, *M. echinops* Fennel, *M. enneacantha* Otto, *M. heteracentra* Otto, *M. intricata* Otto, *M. miqueliana* Pfeiffer, *M. palmeri* Fennel, *M. pyrrhacantha* Pfeiffer, *M. pyrrhacantha pallida* Pfeiffer, *M. salmiana* Fennel, *M. stephani* Hortus, *M. suberecta* Pfeiffer, and *M. villosa* Fennel.

The following names appeared first, published by Forbes (Journ. Hort. Tour Germ. 147. 1837), but are so briefly described that they can not be identified: *Mammillaria cuneiflora* Hitchen, *M. cylindraca* Hitchen, *M. divaricata*, *M. flavescens* Hitchen, *M. grandis* Hitchen, *M. lutescens*, and *M. pulcherrima*. Some of these names were afterwards used, but whether they were applied to the same plants we can not tell.

The following names of *Mammillaria* listed by Haage (Cact. Knit. ed. 2. 1900) are without description: *brandtii* Haage jr., *bruennowii*, *celsiana longispina*, *de grandii*, *deleuili* Rebut, *desertorum*, *donkelaari*, *dubia* Hildmann, *fulvolanata*, *geminiflora*, *glabrescens*, *goeringii*, *grusonii similis*, *guebwilleriana* Haage jr., *hermantiana* Monville, *hevernicksii* Senke, *lapaixi* Rebut, *microdasys*, *monothele*, *morini* Rebut, *multicolor*, *nickelsi*, *nigerrima*, *numina*, *polia* Sieber, *quehlii*, *rebuti*, *roii* Rebut, *roessingii* Gruson, *semilonia*, *simonis*, *lellii*, *variimamma* Ehrenberg, *villa-lerdo*, *wegeneri cristata*, and *xanthispina*.

Schumann, at the close of his treatment of the genus *Mammillaria* (Gesamtb. Kakteen 599. 1898), lists 158 names which he had not been able to refer. Later, Otto Kuntze referred many of the names to the genus *Cactus*, thus making many useless synonyms. Some of these names of Schumann we have been able to refer more or less definitely to other species, but there still remain many which we can not place. Most of them were described without flower and fruit, and since the types were not preserved it is doubtful if many more can be ever identified. The residue is as follows:

MAMMILLARIA ACTINOPLEA Ehrenberg, Allg. Gartenz. 16: 266. 1848.

Mammillaria amabilis Ehrenberg, Allg. Gartenz. 17: 326. 1849.

Mammillaria albiseta Hortus in Förster, Handb. Cact. ed. 2. 354. 1885.

Cactus actinopleus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Cactus amabilis Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Mammillaria crebrispina nitida Monville (Labouret, Monogr. Cact. 75. 1853) is known only as a synonym.

MAMMILLARIA ARGENTA Fennel, Allg. Gartenz. 15: 66. 1847.

Cactus argenteus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA ATRORUBRA Ehrenberg, Allg. Gartenz. 17: 327. 1849.

Cactus atroruber Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA ATROSANGUINEA Ehrenberg, Allg. Gartenz. 17: 270. 1849.

Cactus atosanguineus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA BADISPINA Förster, Hamb. Gartenz. 17: 159. 1861.

MAMMILLARIA BARLOWII Regel and Klein, Ind. Sem. Hort. Petrop. 1860: 46. 1860.

Cactus barlowii Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA BERGENII Ehrenberg, Allg. Gartenz. 17: 326. 1849.

MAMMILLARIA BIFURCA A. Dietrich, Allg. Gartenz. 18: 186. 1850.

MAMMILLARIA BREVISETA Ehrenberg, Allg. Gartenz. 17: 251. 1849.

Cactus brevisetus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA CLOSIANA Roumey, Bull. Soc. Bot. France 2: 372. 1855.

MAMMILLARIA COROLLARIA Ehrenberg, Allg. Gartenz. 17: 294. 1849.

Cactus corallarius Kuntze, Rev. Gen. Pl. 1: 260. 1891.

MAMMILLARIA CORONATA Scheidweiler, Allg. Gartenz. 8: 338. 1840.

Cactus coronatus Kuntze, Rev. Gen. Pl. 1: 261. 1891.

MAMMILLARIA CURVISPINA Otto in Dietrich, Allg. Gartenz. 14: 204. 1846.

Cactus curvispinus Kuntze, Rev. Gen. Pl. 1: 260. 1891. Not Bertero, 1829.

- MAMMILLARIA DECORA Förster, Hamb. Gartenz. **17**: 159. 1861.
Mammillaria decora obscura Förster, Hamb. Gartenz. **17**: 159. 1861.
- MAMMILLARIA EBORINA Ehrenberg, Allg. Gartenz. **17**: 309. 1849.
Cactus eborinus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA EMUNDTSIANA Hortus in Förster, Handb. Cact. ed. 2. 341. 1885.
- MAMMILLARIA ERECTACANTHA Förster, Allg. Gartenz. **15**: 50. 1847.
Cactus erectacanthus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA EUCHLORA Ehrenberg, Allg. Gartenz. **16**: 266. 1848.
Cactus euchlorus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA FELLNERII Ehrenberg, Allg. Gartenz. **17**: 261. 1849.
Cactus fellneri Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA FLAVA Ehrenberg, Allg. Gartenz. **17**: 261. 1849.
Mammillaria tomentosa flava Salm-Dyck, Cact. Hort. Dyck. 1849. 12. 1850.
- MAMMILLARIA GEMINATA Scheidweiler, Allg. Gartenz. **9**: 42. 1841.
Cactus geminatus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- Illustration*: Möllers Deutsche Gart. Zeit. **25**: 475. f. 8, No. 20.
- MAMMILLARIA GIBBOSA Salm-Dyck, Hort. Dyck. 343. 1834.
Cactus gibbosus Kuntze, Rev. Gen. Pl. **1**: 261. 1891. Not Haworth, 1812.
- MAMMILLARIA GLABRATA Salm-Dyck, Cact. Hort. Dyck. 1849. 109. 1850.
Cactus glabratus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA GRANDICORNIS Mühlenpfordt, Allg. Gartenz. **14**: 372. 1846.
Cactus grandicornis Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA HAEMATACTINA Ehrenberg, Allg. Gartenz. **16**: 266. 1848.
Cactus haematactina Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA INCURVA Scheidweiler, Bull. Acad. Sci. Brux. **6**: 92. 1839.
Cactus incurvus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA JUCUNDA Ehrenberg, Allg. Gartenz. **17**: 250. 1849.
Cactus jucundus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA KLEINII Regel, Ind. Sem. Hort. Petrop. **1860**: 7. 1860.
Cactus kleinii Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA LAMPROCHAETA Jacobi, Allg. Gartenz. **24**: 82. 1856.
- MAMMILLARIA LEUCODASYS Salm-Dyck in Scheer, Seemann, Bot. Herald 286. 1856.
Cactus leucodasys Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- Mexico. Probably *M. micromeris* Engelmänn (*fide* Schumann).
- MAMMILLARIA LEUCODICTIA Linke, Allg. Gartenz. **16**: 330. 1848.
Cactus leucodictyus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA LIVIDA Fennel, Allg. Gartenz. **15**: 66. 1847.
Cactus lividus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.

Mammillaria farinosa (Fennel, Allg. Gartenz. **15**: 66. 1847) is referred to *M. livida* by the Index Kewensis.

- MAMMILLARIA MELANACANTHA Hortus in Förster, Handb. Cact. ed. 2. 386. 1885.
- MAMMILLARIA MICANS Dietrich in Linke, Allg. Gartenz. **16**: 330. 1848.
Cactus micans Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA MICRACANTHA Miquel, Linnaea **12**: 16. 1838.
Cactus micracanthus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA MUCRONATA Ehrenberg, Allg. Gartenz. **17**: 294. 1849.
Cactus mucronatus Kuntze, Rev. Gen. Pl. **1**: 260. 1891.
- MAMMILLARIA MULTISETA Ehrenberg, Allg. Gartenz. **17**: 242. 1849.
Cactus multisectus * Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA OBLIQUA Ehrenberg, Allg. Gartenz. **17**: 250. 1849.
Cactus obliquus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

* Kuntze's specific name is credited to Scheidweiler, but we do not find it.

- MAMMILLARIA OBVALLATA Otto in Dietrich, Allg. Gartenz. **14**: 308. 1846.
Cactus obvallatus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA OLORINA Ehrenberg, Allg. Gartenz. **17**: 326. 1849.
Cactus olorinus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA OOTHELE Lemaire, Cact. Gen. Nov. Sp. 37. 1839.
Mammillaria ovimamma Lemaire, Cact. Gen. Nov. Sp. 4. 1839.
Mammillaria ovimamma brevispina Salm-Dyck, Cact. Hort. Dyck. 1849. 108. 1850.
Mammillaria ovimamma oothele Labouret, Monogr. Cact. 85. 1853.
Cactus oothele Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Cactus ovimamma Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PERSICINA Ehrenberg, Allg. Gartenz. **17**: 250. 1849.
Cactus persicanus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PHAEOTRICA Monville in Labouret, Monogr. Cact. 39. 1853.
Cactus phaeotrichus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PLEIOCEPHALA Regel and Klein, Ind. Sem. Hort. Petrop. **1860**: 7. 1860.
Cactus pleiocephalus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA POLYMORPHA Scheer in Mühlenpfordt, Allg. Gartenz. **14**: 373. 1846.
Cactus polymorphus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PORPHYRACANTHA Jacobi, Allg. Gartenz. **24**: 81. 1856.
- MAMMILLARIA PROCERA Ehrenberg, Allg. Gartenz. **17**: 241. 1849.
Cactus procerus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PUGIONACANTHA Förster, Allg. Gartenz. **15**: 50. 1847.
Cactus pugionacanthus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA PUNCTATA Labouret in Förster, Handb. Cact. ed. 2. 293. 1885.
- MAMMILLARIA PURPURASCENS Ehrenberg, Allg. Gartenz. **17**: 260. 1849.
- MAMMILLARIA PURPUREA Ehrenberg, Allg. Gartenz. **17**: 270. 1849.
Cactus purpureus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA REGIA Ehrenberg, Allg. Gartenz. **17**: 269. 1849.
Cactus regius Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA ROSEA Scheidweiler, Hort. Belge **5**: 118. 1838.
Mammillaria rhodeocentra Lemaire, Cact. Gen. Nov. Sp. 52. 1839.
Mammillaria discolor nigricans Salm-Dyck in Walpers, Repert. Bot. **2**: 271. 1843.
Mammillaria rhodeocentra gracilispina Salm-Dyck, Cact. Hort. Dyck. 1849. 14. 1850.
Cactus roseus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
Cactus rhodeocentrus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- Salm-Dyck referred *Mammillaria rosea* to *M. rhodeocentra*, but the former is the older name.
- Illustration*: Hort. Belge **5**: pl. 7. as *Mammillaria rosea*.
- MAMMILLARIA RUFIDULA Ehrenberg, Allg. Gartenz. **17**: 295. 1849.
Cactus rufidulus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA RUFO-CROCEA Salm-Dyck, Cact. Hort. Dyck. 1849. 102. 1850.
Cactus rufo-croceus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA RUSCHIANA Regel, Ind. Sem. Hort. Turic. 4. 1830, in adnot.
Cactus rueschianus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA SEIDELII Terschek, Suppl. Cact. Verz. **1**.
Cactus seidelii Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA SEVERINI Regel and Klein, Ind. Sem. Hort. Petrop. **1860**: 46. 1860.
Cactus severinii Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA SPECIOSA De Vriese, Tijdschr. Nat. Geschr. **6**: 52. 1839. Not Gillies, 1830.
Cactus vrieseanus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA SPECTABILIS Mühlenpfordt, Allg. Gartenz. **13**: 346. 1845.
Cactus spectabilis Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA SUBULIFERA Ehrenberg, Allg. Gartenz. **17**: 242. 1849.
Cactus subulifer Kuntze, Rev. Gen. Pl. **1**: 261. 1891.
- MAMMILLARIA TECTA Miquel, Linnaea **12**: 12. 1838.
Cactus tectus Kuntze, Rev. Gen. Pl. **1**: 261. 1891.

- MAMMILLARIA TOMENTOSA Ehrenberg, Allg. Gartenz. 17: 262. 1849.
Cactus tomentosus Kuntze, Rev. Gen. Pl. 1: 261. 1891.
- MAMMILLARIA VARIMAMMA Ehrenberg, Allg. Gartenz. 17: 242. 1849.
Cactus varimamma Kuntze, Rev. Gen. Pl. 1: 261. 1891.
- MAMMILLARIA WEGENERI Ehrenberg, Bot. Zeit. 1: 738. 1843.
Cactus wegneri Kuntze, Rev. Gen. Pl. 1: 261. 1891.
- MAMMILLARIA ZEGSCHWITZII Terschek, Suppl. Cact. Verz. 1.
Cactus zegschwitzii Kuntze, Rev. Gen. Pl. 1: 261. 1891.
- MAMMILLARIA ZEPNICKII Ehrenberg, Bot. Zeit. 2: 835. 1844.
Cactus zepnickii Kuntze, Rev. Gen. Pl. 1: 261. 1891.

NAMES TO BE EXCLUDED FROM THIS GENUS.

The names *Mammillaria solitaria*, *M. spinosa*, *M. caudata*, *M. ambigua*, and *M. quadrata*, credited to G. Don, with the synonyms *Cactus solitarius*, *C. spinosus*, *C. caudatus*, *C. ambiguus* [Not Bonpland, 1813], and *C. quadratus*, credited to Gillies, respectively, each with a single word description, viz., solitary, spiny, tailed, ambiguous, quadrate, appeared in 1830 (Loudon, Hort. Brit. 194). As they all are said to come from Chile they can not be of this alliance.

Mammillaria brachydelphys Schumann (Just, Bot. Jahresb. 26: 343. 1898) seems to have been intended for *Maibuenia brachydelphys*.

Cereus caudatus Gillies (Sweet, Hort. Brit. ed. 3. 285. 1839) is probably the same as *M. caudata*.

Mammillaria corioides Bosch (Sweet, Hort. Brit. ed. 3. 281. 1839) was described as leather-like and native of South America. It can not be identified, but it is not of this relationship if it comes from South America. Schumann referred it to *Echinocactus*, but it does not belong to that genus as we now define it.

Mammillaria dichotoma (Sweet, Hort. Brit. ed. 3. 281. 1839), described only as forked, can not be identified.

Mammillaria mitis (De Candolle, Prodr. 3: 460. 1828), without description, is credited to Miller (Dict. Gard.), but Miller never used the generic name *Mammillaria*. Pfeiffer and Förster also refer this name to Miller. Steudel states that it is from South America. Kuntze also refers to the same as *Cactus mitis* (Rev. Gen. Fl. 1: 259. 1891). Schumann thought that it might be an *Echinocactus* and, if it really came from South America, as stated by the Index Kewensis, it is probably of the *Echinocactanae*.

Mammillaria speciosa Gillies (Sweet, Hort. Brit. ed. 2. 235. 1830), to which *Cactus speciosus* Gillies is referred as a synonym, is based upon some Chilean plant.

Mammillaria subulata Mühlenpfordt is listed both by Schumann and the Index Kewensis but the name intended was *Pereskia subulata*!

MAMMILLARIA CHILDSI Blanc, Illustr. Cat. 14. 1894.

"This line *Mammillaria* was sent out by us as *M. pectinata* before we bloomed it, from the fact that small plants answered the description exactly. After blooming, however, we discovered that it was a valuable new variety and named it as above. When small, the spines are regular, short and white; as the plant becomes older the spines also increase in size and assume a beautiful purple color. Flowers very numerous, even on small plants; color a clear pink."

We have not been able to identify this plant definitely. From the illustration, which shows large flowers from the center of the plant, we judge that it can not be referred to *Neomammillaria* nor to any of its near relatives. It may be a *Coryphantha*; in fact, at first it was taken for *C. pectinata*. The spines, however, are shown as arranged on vertical ribs, while the central spine is shown as erect; these two characters along with the central purple flowers suggest *Echinomastus erectocentrus*.

Illustration: Blanc, Illustr. Cat. 14.

MAMMILLARIA CORONARIA Haworth, Rev. Pl. Succ. 69. 1821, as to name.

Cactus coronatus Willdenow, Enum. Pl. Hort. Berol. Suppl. 30. 1813. Not Lamarck, 1783.

Judging from Willdenow's original descriptions of this plant it is not of this genus. He says that it is 5 feet long and a foot in diameter and that the central spine of the areole is hooked. Its geographical origin was not recorded and its flowers were not described. It was grown at Berlin prior to 1813 and later at the Chelsea Garden, London. Descriptions of this species are based largely on *Cactus cylindricus* Ortega, a very different plant.

Through the courtesy of N. E. Brown we have a photograph of *Mammillaria coronaria* from Haworth's collection with the date "Feb. 20, 1846." This photograph answers Haworth's brief description and differs from Willdenow's in having the spines all straight. Haworth's plants we would refer to *Neomammillaria*.

Cactus coronarius Willdenow, given by Haworth as a synonym of *Mammillaria coronaria*, is a mistake for *C. coronatus*.

The variety *Mammillaria coronaria minor* was briefly described by Förster (Handb. Cact. 212. 1846).

MAMMILLARIA FULVISPIA Haworth, Phil. Mag. 7: 108. 1830.

Cactus fulvispinus Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Mammillaria rhodantha fulvispina Schelle, Handb. Kakteenk. 257. 1907.


This plant was said by Haworth to come from Brazil and if so it is to be excluded from this relationship. Pfeiffer associated the name with a Mexican specimen which has led to its being referred by later writers to *M. rhodantha*. The varieties *M. fulvispina media* and *M. fulvispina minor* (Salm-Dyck, Cact. Hort. Dyck. 1844. 8. 1845) were not described.

MAMMILLARIA PICTURATA Labouret, Rev. Hort. IV. 4: 28. 1855.

Simple, cylindric, 8 cm. high, 5 cm. in diameter; radial spines 20, white, setiform, 4 mm. long; central spines 6, yellowish; flowers and fruit unknown.

Although Labouret stated that this plant came from Mendoza, Argentina, the Index Kewensis says Chile. If it is in southern South America, it does not belong to *Neomammillaria*.

The illustration (figure 1840) at the bottom of this page is of *Neomammillaria milleri* described on page 156.



Subtribe 7. EPIPHYLLANAE.

Mostly epiphytic and night-blooming cacti, generally growing on trees, but sometimes on the earth when this is rich in humus, rarely in the crevices of rocks, much branched, spineless (except *Eccremocactus* and some species of *Epiphyllanthus*); joints several or many, usually flat except at base, often thin, with the areoles borne along the margin (except in *Epiphyllanthus*); flowers regular (except in *Zygocactus* and *Epiphyllanthus*); perianth various; filaments usually long and slender; style long and slender; fruit spineless, usually red or purple, either naked or bearing a few scales (rarely many), these usually with naked axils; seeds small, black.

We recognize 9 genera, diverse both in the plant-body and in the flowers. While apparently not closely related among themselves, the genera forming this subtribe are not any more closely related to other genera, either in the *Cereanae* or in the *Rhipsalidanae*.

KEY TO GENERA.

- Plants branching dichotomously.
 - Perianth irregular.
 - Joints thin and leaf-like with toothed margin; areoles all marginal 1. *Zygocactus* (p. 177)
 - Joints thick, without teeth, bearing areoles all around 2. *Epiphyllanthus* (p. 180)
 - Perianth regular or nearly so; joints thin 3. *Schlumbergera* (p. 182)
- Plants branching irregularly.
 - Perianth-segments spreading or reflexed; flowers mostly large.
 - Tube of flower definitely longer than limb. 4. *Epiphyllum*. (p. 185)
 - Tube of flower not longer than limb.
 - Perianth campanulate, its segments few.
 - Stamens few; flowers small. 5. *Disocactus* (p. 201)
 - Stamens many; flowers large. 6. *Chiapasia* (p. 203)
 - Perianth short-funnel form, its segments many.
 - Outer perianth-segments short, obtuse or rounded, the inner white. 7. *Eccremocactus* (p. 204)
 - Outer perianth-segments acute or acuminate, the inner rose or red. 8. *Nopalxochia* (p. 204)
 - Perianth-segments erect; flowers small 9. *Wittia*. (p. 206)

1. ZYGOCACTUS Schumann in Martius, Fl. Bras. 4²: 223. 1890.

Stems dichotomously much branched, flattened, divided into short joints; flowers terminal, polychromic, irregular; ovary terete, smooth, gradually broadening from base, bearing minute scales at top; flower-tube abruptly bent just above the ovary, ending in a serrate mouth, bearing petaloid spreading scales scattered along its sides; stamens slender, white, arranged in 2 clusters; outer stamens borne along inside of flower-tube from near base to near middle; inner clusters of stamens about 20, arising from center and forming a short tube about base of style with an inner deflexed toothed membrane, upper part free, and all appressed against upper side of flower-tube and tipper perianth-segments; style purple, slender, as long as stamens and usually not surrounded by them; stigma-lobes linear, purple, erect and adhering (so far as we have seen); fruit purple, turgid, not at all angled; skin thin; seeds dark brown to nearly black, shining.

Type species: *Epiphyllum truncatum* Haworth.

This genus has passed for many years under the name of *Epiphyllum* but that name was wrongly applied to it. One species is here recognized, although several have been proposed by previous authors.

The generic name is from *ζυγόν* yoke and *κάκτος* cactus, referring, doubtless, to the peculiarly jointed stems.

1. Zygocactus truncatus (Haworth) Schumann in Martius, Fl. Bras. 4²: 224. 1890.

- Epiphyllum truncatum* Haworth, Suppl. Pl. Succ. 8. 1819.
- Cactus truncatus* Link, Enum. Pl. 2: 24. 1822.
- Cereus truncatus* Sweet, Hort. Brit. 272. 1826.
- Epiphyllum altensteinii* Pfeiffer, Enum. Cact. 128. 1837.
- Epiphyllum truncatum altensteinii* Lemaire, Cact. Gen. Nov. Sp. 76. 1839.
- Epiphyllum purpurascens* Lemaire, Hort. Univ. 2: 349. 1841.
- Epiphyllum truncatum violaceum* Morren, Belg. Hort. 16: 260. 1866.
- Epiphyllum truncatum spectabile* Morren, Belg. Hort. 16: 260. 1866.
- Zygocactus altensteinii* Schumann in Martius, Fl. Bras. 4²: 225. 1890.
- Epiphyllum delicatum* N. K. Brown, Gard. Chron. 111. 32: 411. 1902.
- Epiphyllum delicatulum* Schumann, Monatsschr. Kakteenk. 13: 9. 1903.
- Zygocactus delicatus* Britton and Rose, Contr. U. S. Nat. Herb. 16: 260. 1913.

Joints dark glossy green, about 3 cm. long, sharply serrate, with two prominent teeth at other-wise truncate apex; terminal areole broad and thin, filled with brown wool and bristles; flowers 6


to 7 cm. long; tube 2 cm. long; inner perianth-segments scarlet to white, oblong, obtuse to acute, reflexed; filaments white; style purple throughout; fruit obovoid, 1.5 to 2 cm. long.

Type locality: Brazil.

Distribution: Mountains, state of Rio de Janeiro, Brazil.

This species has been cultivated widely for many years under various names. It was introduced into cultivation about 1818 and, according to Edwards, flowered first in England in 1822 and has since been a great favorite as a household plant, blooming freely about the end of the year, hence the name Christmas cactus. It is also called crab cactus and ringent-flowered cactus.

Schumann gives as synonyms of this species *Epiphyllum salmoneum* and *E. spectabile*, referring them to Cels's Catalogue, which, however, we have not seen.



FIGS. 185 and 186.—Flowering branch and fruiting joint of *Zygocactus truncatus*.

Cereus truncatus altensteinii (Salm-Dyck, Hort. Dyck. 65. 1834) occurs in literature, sometimes attributed to Otto, but we have seen no description. We follow Löfgren, who refers *Zygocactus altensteinii* to *Z. truncatus*. The type came from the Organ Mountains near Rio de Janeiro; in 1915, Dr. Rose visited these mountains, where he found the true *Z. truncatus*.

There are many garden varieties, most of which are very beautiful. Among these are *Epiphyllum gibsonii*, introduced in 1886, with dark orange-red flowers, and *Epiphyllum guedeneyi*, of unknown origin, with large flowers, the outer segments white, tinged with sulphur, and the inner ones creamy white; the variety is referred by some to *Phyllocactus guedeneyi*. Nicholson (Dict. Gard. 1:517) describes some of the best as follows:

"*Bicolor*, white, edged with rose; *coccineum*, rich deep scarlet; *elegans*, bright orange-red, centre rich purple; *magnificum*, flowers large, white, tips bright rose-colored; *roseum*, bright rose; *ruckerianum*, deep reddish purple, with a rich violet centre; *salmoneum*, reddish salmon; *spectabile*, white, with delicate purple margin; *violaceum superbum*, pure white, rich deep purple edge."

Rümpel (Förster, Handb. Cact. ed. 2. 870, 871. 1885) described nine varieties, among which are *cruentum* and *tricolor*; *E. truncatum cruentum* was also briefly described by Morren (Belg. Hort. 16: 260. 1866). Among other varieties are *albiflorum*, *aurantiacum*, *grandidens*, *minus*, *purpureum*, and *vanhoutteanum*.

Epiphyllum ruckeri Paxton (Mag. Bot. 12: 46. 1846) was described from cultivated plants of unknown origin as an improved variety of *Epiphyllum truncatum*. It may have been a hybrid.

Epiphyllum truncatum multiflorum was given as a synonym of *Epiphyllum altensteinii* by Pfeiffer (Enum. Cact. 128. 1837).

Epiphyllum elegans Cels and *E. violaceum* Cels (Förster, Handb. Cact. 446. 1846) were supposed to be only varieties of *Epiphyllum truncatum*.

Schelle (Handb. Kakteenk. 223. 1907) lists more than fifty forms of *Epiphyllum truncatum*; the following not hitherto mentioned by us under *Epiphyllum* have the regular Latin form:

| | | | |
|---------------------|-----------------------|-----------------------|------------------------|
| amabile roseum | maximum | salmoneum brasiliense | spectabile superbum |
| carmineum | morellianum | salmoneum flavum | splendens |
| gracile | pallidum roseum | salmoneum marginatum | translucens |
| grandiflorum rubrum | purpureum | salmoneum rubrum | violaceum album |
| harrisonii | rubrum violaceum | snowi | violaceum grandiflorum |
| lateritium album | salmoneum aurantiacum | spectabile carmineum | superbum |
| makoyanum | | | |

Illustrations: Nov. Herb. Amat. pl. 83; Loudon, Encycl. Pl. 413. f. 6903; Loddiges, Bot. Gart. 13: pl. 1207; Curtis's Bot. Mag. 52: pl. 2562; Edwards's Bot. Reg. 9: pl. 696;




FIG. 187.—*Zygocactus truncatus*.

Reichenbach, Fl. Exot. pl. 325; Hooker, Exot. Fl. 1: pl. 20, as *Cactus truncatus*; Wiener Ill. Gart. Zeit. 18: 265. f. 55, as *Phyllocactus delicatus*; Blühende Kakteen 1: pl. 25; Cact. Journ. 1: 34, 114; Cycl. Amer. Hort. Bailey 2: f. 765; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 61, A, B, C; Schumann, Gesamtb. Kakteen f. 9, 43; Hort. Univ. 7: facing 132; Karsten, Deutsche Fl. 887. f. 501, No. 3; ed. 2. 2: 456. f. 605, No. 3; Förster, Handb. Cact. ed. 2. 129. f. 5; Rümpler, Sukkulente f. 87; Hort. Franc. 11. 4 pl. 3; Schelle, Handb. Kakteenk. 223. f. 145; Balt. Cact. Journ. 1: 49; Flörialia 42: 375; Gard. Chron. 1847: 324; 11. 6: 808. f. 148; 111. 7: 173. f. 29; 111. 19: f. 1; West Amer. Sci. 7: 172; Amer. Gard. 11: 534; Schelle, Handb. Kakteenk. 224. f. 146; 225. f. 147; Rother, Praktischer Leitfaden Kakteen 104; Belg. Hort. 16: pl. 257; also the vars. *spectabile*, *cruentum*, and *violaceum*; Deutsches Mag. Gart. Blumen. 1852: pl. 176. f. 2; Gartenwelt 4: 230; Goebel, Pflanz. Schild. 1: f. 55 (seedling); Garten-Zeitung 4: 182. f. 42, No. 2; Jacquin, Ecl. Pl. Rar. 2: pl. 142, as *Epiphyllum truncatum*; Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 28, as *E. altensteinii*; Gard. Chron. 111. 32: f. 140, as *E. delicatum*; Schumann, Gesamtb. Kakteen Nachtr. f. 9; Monatsschr. Kakteenk. 13: 7, as *E. delicatulum*; Deutsches

Mag. Gart. Blumen. 1852: pl. opp. 176, f. 1, as *E. truncatum elegans*; Arch. Jard. Bot. Rio de Janeiro 2: pl. 3, as *Zygocactus delicatus*; Van G  el, Sert. Bot. 1: pl. 117 as *Cactus truncatus*; Martius, Fl. Bras. 4²: pl. 46; Contr. U. S. Nat. Herb. 16: pl. 80; Stand. Cycl. Hort. Bailey 6: f. 4055.

Figure 185 shows a plant in the New York Botanical Garden which flowered December 15, 1911; figure 186 shows a fruiting joint collected by Dr. Rose in the Organ Mountains of Brazil in 1915 (No. 20819); figure 187 is from a photograph of a cultivated plant obtained by Dr. Rose in the Botanical Garden at Rio de Janeiro in 1915 (No. 20855).

2. EPIPHYLLANTHUS Berger, Rep. Mo. Bot. Gard. 16: 84. 1905.

Plants either epiphytic or growing in shade of rock in rich humus, often in clumps, more or less branched; joints globular, cylindric or much flattened; areoles scattered over surface of joints, circular, tomentose, either with or without spines; flowers zygomorphic, slender, purple to white; stamens somewhat exserted, arranged in 2 series, those forming the inner series united at base; style slender, a little longer than stamens; ovary angled, bearing a few small scales; fruit small.

Type species: *Epiphyllanthus obtusangulus* Berger.

The type of this genus has long been treated as a species of *Cereus*, although its dissimilarity to *Cereus* proper or to any of its immediate relatives must have been observed. It was left to Alwin Berger to call attention to its true alliance and to propose for it a new generic name; his statement regarding it is so clear that we quote from it as follows:

"This very strange little plant, still rare in cultivation, can not be considered either a *Cereus* or an *Epiphyllum*. But no doubt it is much more nearly allied to the latter than to the former genus. Schumann brought it into *Cereus* on account of its round and ribbed stems, but there exists no *Cereus* of a similar articulated growth; only with *Rhipsalis* and *Epiphyllum* can it be compared. The plant resembles somewhat a minute *Platyopuntia*. The joints are slightly flattened and have numerous little prominent areoles distributed spirally all over the surface. In this it differs greatly from *Epiphyllum* with which it agrees in all the characters of the flowers, the angular, nearly alate ovary, and especially in the inner stamens being united at the base into a small incurved membrane. Also, the fruit resembles more that of an *Epiphyllum* than that of a *Cereus*. The flowers rise from the top of the joints as in *Epiphyllum*. The plant is best considered as generically different from both, but must be placed with *Epiphyllum* and *Rhipsalis* among the *Inarmatae* of K. Schumann."

We recognize 3 species, all from central Brazil. All occur on the high mountain Itatiaya, province of Rio de Janeiro; what their actual relationships may be can be determined only by further field observations. They may all be referable to one variable species.

The generic name was given because of the resemblance of the flowers of the type species to those of *Epiphyllum truncatum* (*Zygocactus*).

KEY TO SPECIES.

- | | |
|---|------------------------------|
| Joints or some of them flattened, <i>Opuntia</i> -like. | 1. <i>E. obovatus</i> |
| Joints terete or obtusely angled. | |
| Flower purple to rose. | 2. <i>E. microsphaericus</i> |
| Flowers white. | 3. <i>E. candidus</i> |

1. *Epiphyllanthus obovatus* (Engelmann).

Epiphyllum obovatum Engelmann in Schumann, Gesamtb. Kakteen 224. 1897.

Epiphyllum opuntoides L  fgren and Dus  n, Arch. Mus. Nac. Rio de Janeiro 13: 49. 1905.

Zygocactus opuntoides L  fgren, Arch. Jard. Bot. Rio de Janeiro 2: 26. 1918.

Usually growing in shade of rocks, at first erect, becoming more or less decumbent, very much branched; joints usually . to 7 cm. long, obovate to oblong, more or less flattened, often suggesting small joints of some *Opuntia*, bearing scattered areoles and these often spinescent; old and lower joints often nearly terete, bearing large areoles with numerous short yellow spines; flowers 5 cm. long, purple; ovary naked.


Type locality: Brazil.

Distribution: Central Brazil.

Dr. Rose collected this species on Itatiaya, altitude about 2,300 meters, in July 1915 (No. 20495); the plant did not do well in cultivation with us and his specimens died.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 2: pl. 4, as *Zygocactus opuntioides*; Arkiv Bot. Stockholm 8: pt. 7. 10, as *Epiphyllum opuntioides*.

Figure 188 is reproduced from the first illustration cited above; figure 189 is reproduced from the second illustration cited above.



FIGS. 188 and 189.—*Epiphyllanthus obovatus*.

2. *Epiphyllanthus microsphaericus* (Schumann).

Cereus microsphaericus Schumann in Martius, Fl. Bras. 4²: 297. 1890.

Cereus parvulus Schumann in Martius, Fl. Bras. 4²: 297. 1890.

Cereus obtusangulus Schumann in Martius, Fl. Bras. 4²: 298. 1890.

Cereus anomalus Schumann, Keys Monogr. Cact. 16. 1903.

Epiphyllanthus obtusangulus Berger, Rep. Mo. Bot. Gard. 16: 84. 1905.

Zygocactus obtusangulus Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 28. 1918.

Low, at first erect, much branched and more or less prostrate, growing under rocks and perhaps epiphytic on trees; joints slender, terete or obtusely angled, somewhat spiny or often naked; flowers all terminal, purple to rose.

Type locality: Province of Rio de Janeiro, Brazil.

Distribution: Central Brazil.

Dr. Rose collected this species on Itatiaya, Brazil, in 1915 (No. 20494), growing at higher altitudes than *E. obovatus*.

Epiphyllum obtusangulum Lindberg (Martius, Fl. Bras. 4²: 198. 1890), usually referred here as a synonym, has not been published.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 2: pl. 5, as *Zygocactus obtusangulus*; Schumann, Gesamtb. Kakteen f. 30, as *Cereus obtusangulus*.

Figure 190 is reproduced from the first illustration cited above.

3. *Epiphyllanthus candidus* (Löfgren).

Zygocactus candidus Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 30. 1918.

Usually epiphytic on shrubs, but sometimes growing in the shade of large boulders; joints usually terete or nearly so, 2 to 4 cm. long, naked or sometimes bristly; flowers solitary, terminal, white; fruit globose, red.

Type locality: On Itatiaya, Brazil.

Distribution: Known only from the type locality.




FIG. 190.—*Epiphyllanthus microsphaericus*.

FIG. 191.—*Epiphyllanthus candidus*.

Dr. Rose collected this species on the very top of Itatiaya, growing in the shade of rocks (No. 20610) and in the deep cleft of the rock cap through which the ascent to the top is made.

Epiphyllum candidum Barbosa-Rodrigues (Arch. Jard. Bot. Rio de Janeiro 2: 30. 1918) is only a name.

Illustration: Arch. Jard. Bot. Rio de Janeiro 2: pl. 6, as *Zygocactus candidus*.

Figure 191 is reproduced from the illustration cited above.

3. *SCHLUMBERGERA* Lemaire, Rev. Hort. iv. 7 253. 1858.

Similar in habit to *Zygocactus*; stems much branched; joints short, crenate or serrate, mostly flattened; flowers purple to scarlet, regular; tube very short; stamens in 2 clusters, one scattered over the throat, the other forming a short tube at base of flower and surrounding style or free at base; ovary and fruit strongly 5-angled, naked or rarely bearing areole on one of the ribs and crowned by 5 more or less persistent, sepal-like scales; fruit hard, often remaining on plant for a long time.

Type species: *Epiphyllum russellianum* Hooker.

The taxonomic history of the two species here recognized is interesting. *Schlumbergera gaertneri* was at first supposed to be conspecific with *S. russelliana* and was made a variety

of that species by Regel. In 1890 Schumann considered them distinct species but congeneric; in 1897 he referred them to different genera. Both species are native of Brazil.

These plants have usually been associated with *Zygocactus truncatus* and all included in *Epiphyllum*. Although resembling *Zygocactus* very much in habit, they differ from it in flower and fruit characters. The flowers are nearly regular, not strongly oblique; are nearly rotate, not elongated; the stamens are of equal length and in a cylindric cluster shorter than the style, not of unequal lengths and in a flattened cluster, not extending beyond the style; the ovary and fruit are strongly angled, not terete.

Lemaire named the genus for Frederick Schlumberger, an amateur student of plants and a collector of cacti, begonias, and bromelias.

KEY TO SPECIES.

Flowers scarlet 1. *S. gaertneri*
Flowers purplish 2. *S. russelliana*

1. *Schlumbergera gaertneri* (Regel) Britton and Rose, Contr. U. S. Nat. Herb. 16: 260. 1913.

Epiphyllum russellianum gaertneri Regel, Gartenflora 33: 323. 1884.

Epiphyllum makoyanum W. Watson, Gard. and For. 2: 243. 1889.

Epiphyllum gaertneri Schumann in Martius, Fl. Bras. 4: 218. 1890.

Phyllocactus gaertneri Schumann in Rümpler, Sukkulente 147. 1892.




FIG. 192.—*Schlumbergera gaertneri*.

Branches spreading, the terminal ones often pendent; joints usually flattened, but sometimes 3 to 6-angled, fleshy, 5 cm. long or more by 2 cm. broad, dull green except the purplish crenate margins; areoles small, with short white wool and a few yellowish bristles; flowers 1 to 3, usually all at distal end of the terminal branches, 4 cm. long, dark scarlet; outermost perianth-segments usually 5, short, thick, triangular, drying separately from the others; outer perianth-segments spreading; innermost perianth-segments more erect, nearly distinct, acute; all of the segments, except the 5 outer ones, more or less coalesce and withering, remaining on top of ovary; ovary crowned by a slightly depressed disk or umbilicus with upturned margin, which passes into the flower-tube; on the margin are borne the free stamens, 1.5 cm. long, red; stigma-lobes 6, linear, cream-colored; ovary dark red, angled, 12 mm. long; fruit red, oblong, 15 mm. long, depressed at apex, in cultivation ripening in July.

Type locality: Minas Geraes, Brazil.

Distribution: Brazil.

While the joints are usually much flattened, yet they are sometimes strongly angled. In some cases too the juvenile growth is peculiar, forming short stubby joints with 6 ribs, with closely set areoles, each bearing a cluster of 7 or more bristly spines.

The plant flowers abundantly in Washington in April.

The two varieties *Epiphyllum gaertneri coccineum* and *E. gaertneri mackoyanum* (Monatsschr. Kakteenk. 7: 101. 1897) are doubtless forms of this species.

Illustrations: Wiener Ill. Gart. Zeit. 10: 136. f. 60; Rev. Hort. 59: pl. opt. 516; Blanc, Cacti 64. 1002 Cact. Journ. 1: 9, 114; Gartenflora 33: pl. 1172; 39: f. 96; Rev. Hort. Belg. 15: f. 23; pl. [19.] f. 2, opp. 229, as *Epiphyllum russellianum gaertneri*; Schelle, Handb. Kakteenk. 213. f. 141; Curtis's Bot. Mag. 117: pl. 7201; Gartenwelt 10: 559, as *Epiphyllum gaertneri*; Blühende Kakteen 1: pl. 21; Thomas, Zimmerkultur Kakteen 19;




FIG. 193.—*Schlumbergera gaertneri*.

Monatsschr. Kakteenk. 4: 107; Rümpler, Sukkulente 148. f. 80, as *Phyllocactus gaertneri*; Rev. Hort. Belg. 15: pl. [19.] f. 1, opp. 229; Journ. Hort. Home Farm. III. 18: 362. f. 58, as *Epiphyllum makoyanum*.

Figure 192 is from a photograph of a plant which flowered in the New York Botanical Garden in 1912; figure 193 shows a fruiting plant in the collections of the U. S. Department of Agriculture at Washington, D. C.

2. *Schlumbergera russelliana* (Gardner) Britton and Rose, Contr. U. S. Nat. Herb. 16: 261. 1913.

Cereus russellianus Gardner in Lemaire, Hort. Univ. 1: 31. 1839.

Epiphyllum russellianum Hooker in Curtis's Bot. Mag. 66: pl. 3717. 1840.

Phyllocactus russellianus Salm-Dyck, Cact. Hort. Dyck. 1844. 37. 1845.

Epiphyllum truncatum russellianum G. Don in Loudon, Encycl. Pl. ed. 3. 1378. 1855.

Schlumbergera epiphyllodes Lemaire, Rev. Hort. IV. 7: 253. 1858.

Epiphytic, growing on trees, rocks, or in humus, often found in dark crevices, 1 to 3 dm. long, either hanging or erect, much branched, divided into short joints; joints 1 to 2.5 cm. long; lower joints usually terete, covered with a brown epidermis; young joints green, flat, usually thin, with 1 or 2 small teeth on a side, 8 mm. broad or less, usually truncate at apex; areoles in axils of teeth, small, naked or bearing 1 or 2 bristles; flowers terminal, 4 to 5 cm. long, reddish purple; style slender, purple; ovary glabrous, sharply 4-angled, 1-celled; ovules numerous, arranged in 4 or 5 vertical double rows along walls of ovary; fruit described as red, 4-angled, or narrowly winded.

Type locality: Organ Mountains, Brazil.

Distribution: Brazil.

This plant was introduced into England in 1839. It was named by G. Gardner for the Duke of Bedford, who had sent him to Brazil to collect plants. The Duke of Bedford brought together at Woburn Abbey a very large and choice collection of cacti which became one of the finest in England. His gardener, Mr. James A. Forbes, published a catalogue of this collection in 1837.

Two varieties of this species are mentioned in horticultural works, namely, var. *rubra* and var. *superbum* under *Epiphyllum russellianum*.

Illustrations: Curtis's Bot. Mag. **66**: pl. 3717; Watson, Cact. Cult. 42. f. 9; Dict. Gard. Nicholson Suppl. 346. f. 370; Gartenflora **33**: pl. 1172; Förster, Handb. Cact. ed. 2. 873. f. 119; Rother, Praktischer Leitfaden Kakteen 106; Paxton's Mag. Bot. **10**: facing 245, as *Epiphyllum russellianum*; Hort. Univ. **1**: pl. 5, as *Cereus russellianus*; Rümpler, Sukkulente 146. f. 79, as *Phyllocactus russellianus*; Cycl. Amer. Hort. Bailey **2**: f. 766, as *Epiphyllum truncatum russellianum* (perhaps a hybrid); Contr. U. S. Nat. Herb. **16**: pl. 81.

SPECIES OF THIS RELATIONSHIP.

EPIPHYLLUM BRIDGESII Lemaire, Illustr. Hort. **8**: Misc. 5. 1861.

Epiphyllum truncatum bridgesii Rümpler in Förster, Handb. Cact. ed. 2. 870. 1885.

Epiphytic; joints green, flattened with 2 or more crenations on the side; areoles more or less setose, the setae yellowish brown; flowers terminal, 6 cm. long, nearly regular, purplish to crimson; perianth-segments oblong, acute; stamens long-exserted; style about as much exserted as stamens, purplish; ovary angled, angles sometimes bearing setose areoles.

Type locality: Not cited. Described from garden plants of unknown origin.

Distribution: Brazil or Bolivia or both.

This plant was described by Lemaire from a vegetative specimen seen in the collection of L. Desmet and from one in the collection of Schlumberger. He associated it with *Epiphyllum russellianum*, with which it must be allied, rather than with *E. truncatum*, to which it is referred as a synonym by the Index Kewensis.

Schlumberger had named the plant *Epiphyllum rueckerianum*, and here this name, often referred to in horticultural literature, should be referred.

It was briefly described by W. Watson (Gard. For. **2**: 243. 1889), who writes of its being awarded a first-class certificate at a flower show.

Schumann unfortunately describes the flower as zygomorphic, which may be an error; specimens recently sent to us from A. Berger have a regular flower. The ovary was originally described as angled and this is one of the differences between *Zygocactus* and *Schlumbergera*.

We do not know the origin of this plant. As it seems to have been introduced by Bridges it may have come from Bolivia, where he did much of his work.

This plant is sometimes called *Epiphyllum truncatum rueckerianum*.

Illustration: Dict. Hort. Bois 497. f. 347, as *Epiphyllum ruckerianum*.

4. EPIPHYLLUM (Hermann) Haworth, Syn. Pl. Succ. 197. 1812.

Phyllocactus Link, Handb. Erkenn. Gewächse **2**: 10. 1831.

Phyllocereus Miquel, Bull. Sci. Phys. Nat. Néerl. **112**. 1839.

Plants mostly epiphytic, the main stem often terete and woody; branches usually much flattened, often thin and leaf-like, sometimes 3-winged; areoles small, borne along the margins of the flattened branches; spines usually wanting in mature plants, but often represented in seedlings and juvenile forms by slender bristles; true leaves wanting; cotyledons rather large, sometimes persisting for a long time; flowers usually large, in some species nocturnal, in others diurnal, either odorless or very fragrant; flower-tube longer than the limb, in some species greatly elongated; filaments usually long, borne at top of tube or scattered over surface of throat; style elongated, white or

colored; stigma-lobes several, linear; perianth soon dropping from the ovary; fruit globular or short-oblong to narrowly oblong, often with low ridges, sometimes tubercled, red or purple, edible or insipid, when mature splitting down one side and exposing the white or crimson pulpy interior; seeds black, shining.

Type species: *Cactus phyllanthus* Linnaeus.

The generic name is from *ἐπί* upon, and *φύλλον* leaf, as it was supposed that the flowers were borne on leaves; it is a misnomer, for the flowers are not borne on leaves but on stems as in all other cacti.

In 1890 K. Schumann recognized 15 species; but, as a number of new ones were described soon afterward, he increased this number to 21 in his *Keys of the Monograph* published in 1903. In our treatment 16 species are recognized.

The name *Epiphyllum* is often used for a different group of cacti, that is, the crab cactus; the type species of *Epiphyllum* is, however, in the genus as we have here limited it. When Haworth published the genus he referred to it but one species, *Epiphyllum phyllanthus*, but he later added another species, *E. truncatum*, which, when it was found to belong to a different generic type, was erroneously allowed to retain the name *Epiphyllum*, while *Epiphyllum phyllanthus* became the type of the genus *Phyllocactus*, which, when first described in 1831, contained but a single species, so that *Epiphyllum* and *Phyllocactus* were based on the same type and *Phyllocactus* is a synonym of *Epiphyllum*. This is also true of *Phyllocereus*, which was based on the *Epiphyllum* of Haworth (*Syn. Pl. Succ.* 197. 1812), where only *E. phyllanthus* is described.




FIG. 194.—Top of fruiting branch of *Epiphyllum phyllanthus*. $\times 0.66$.

The pre-Linnaean species of this genus were usually referred to *Cereus* and, for it, the section *Alati* in *Cereus* was proposed by De Candolle (*Prodr.* 3: 469. 1828). Linnaeus, however, referred the only species which he recognized to *Cactus*, and Philip Miller referred the same species to *Opuntia*, but neither have had many followers.

Haworth (*Phil. Mag.* 6: 108, 109. 1829) followed by Don (*Hist. Dichl. Pl.* 3: 170. 1834) divides the genus into two sections, the *Nocturna* and the *Diurna*.

Phyllanthus Nicker (*Elam.* 2: 85. 1790) is generally supposed to be a synonym of this group but the genus is not typified; the *Index Kewensis* refers it to *Cereus* (?); Dr. E. L. Greene (*Leaflets* 1: 52) says that it applied to *Phyllanthus* and *Opuntia* of earlier authors: the *Phyllanthus* here referred to was *Cactus phyllanthus* Linnaeus.

Hermann (Par. Botavus Prodr. Add. 2. 1689) first used the name *Epiphyllum* when he listed the name *Epiphyllum americanum*. Haworth credited the name also to Hermann in 1812 when he established the genus.

KEY TO SPECIES.

- A. Perianth-tube 7 to 9 times as long as the limb. 1. *E. phyllanthus*
- AA. Perianth-tube 1½ to 3 times as long as the limb.
 - B. Ultimate joints acuminate.
 - Flowers 25 to 30 cm. long 2. *E. oxypetalum*
 - Flowers 10 to 15 cm. long.
 - Margins of joints crenate 3. *E. pumilum*
 - Margins of joints undulate 4. *E. caudatum*
 - BB. Ultimate joints acute, obtuse or rounded.
 - C. Joints deeply lobed.
 - Joints 2 to 7 cm. broad.
 - Lobes of joints spreading; outer perianth-segments lemon-yellow. 5. *E. darrabii*
 - Lobes of joints pointing forward; outer perianth-segments reddish yellow 6. *E. anguliger*
 - Joints very large, up to 25 cm. broad 7. *E. grandilobum*
 - CC. Joints crenate or nearly entire.
 - D. Joints deeply crenate, thick; perianth-tube bearing foliaceous scales 8. *E. crenatum*
 - DD. Joints low-crenate to nearly entire; perianth-tube without foliaceous scales.
 - E. Sinui of the joint-margins very narrow; flowers up to 20 cm. broad; stamens yellow 9. *E. macropterum*
 - EE. Sinui of the joint-margins open; flowers 15 cm. wide or less; stamens not yellow.
 - Ovary and fruit bearing linear scales 10. *E. lepidocarpum*
 - Ovary and fruit without linear scales.
 - Flowers 10 to 13 cm. long 11. *E. pittieri*
 - Flowers 15 to 28 cm. long.
 - Flowers about 28 cm. long; style orange 12. *E. guatemalense*
 - Flowers 15 to 25 cm. long; style white or pink.
 - Joints very stiff 13. *E. strictum*
 - Joints flexible or moderately stiff.
 - Joints very large, up to 1 meter long and 22 cm. wide 14. *E. stenopetalum*
 - Joints smaller, rarely over 7 cm. wide.
 - Joints shallowly crenate or subdentate; species of Costa Rica 15. *E. cartagense*
 - Joints deeply crenate; species of Tobago, Trinidad, and Venezuela 16. *E. hookeri*

1. *Epiphyllum phyllanthus* (Linnaeus) Haworth, Syn. Pl. Succ. 197. 1812.

- Cactus phyllanthus* Linnaeus, Sp. Pl. 469. 1753.
- Opuntia phyllanthus* Miller, Gard. Dict. ed. 8. No. 9. 1768.
- Cereus phyllanthus* De Candolle, Prodr. 3: 469. 1828.
- Phyllocactus phyllanthus* Link, Handb. Erkenn. Gewächse 2: 11. 1831.
- Rhipsalis macrocarpa* Miquel, Bull. Sci. Phys. Nat. Néerl. 1838: 49. 1838 (in most part).
- Rhipsalis phyllanthus* Schumann in Martius, Fl. Bras. 4: 298. 1890 (in part).
- Hariota macrocarpa* Kuntze, Rev. Gen. Pl. 1: 263. 1891.
- Phyllocactus phyllanthus paraguayensis* Weber, Dict. Hort. Bois 957. 1898.
- Phyllocactus phyllanthus boliviensis* Weber, Dict. Hort. Bois 957. 1898.
- Phyllocactus phyllanthus columbiensis* Weber, Dict. Hort. Bois 957. 1898.
- Epiphyllum gaillardae* Britton and Rose, Contr. U. S. Nat. Herb. 16: 240. 1913.
- Phyllocactus gaillardae* Vaupel, Monatsschr. Kakteenk. 23: 87. 1913.

Elongated and much branched; main branches narrow, terete or 3 or 4-angled, woody; terminal joints elongated, terete or 3-angled below, usually flat or thin, rarely 3-winged, bright green with a purple margin, sometimes 7 cm. broad, obtuse, the margin coarsely serrate, the teeth obtuse; flower slender, 25 to 30 cm. long, the slender tube very much longer than the limb, green, the limb greenish or white, its segments narrow, 2 to 2.5 cm. long; scales on flower-tube few, minute, spreading; style long, slender, pinkish (Schumann says white); filaments short; stigma-lobes short, white; fruit oblong, 7 to 9 cm. long, somewhat 8-ribbed, bright red; pulp white; seeds large, black, numerous.

According to De Candolle, the flowers are nocturnal and odoriferous.

Type locality: Brazil.

Distribution: Panama to British Guiana, Bolivia, Peru, and Brazil. Recorded from Paraguay.

The species has been recorded from the West Indies, apparently erroneously.

Our description is based on field notes made by Dr. Rose in Brazil in 1915 which differ slightly from published descriptions. This plant is common in the woods along the coast of eastern Brazil, often growing in inaccessible places high up in the great trees. Open flowers were not seen, but buds, fruit, and seeds were obtained. Living plants were collected and these have done well; one flower appeared in the collection of the Department of Agriculture during Dr. Rose's absence in Ecuador in 1918. The caretaker, Mr. Fraile, describes the flower as long and slender and very unlike other species of *Epiphyllum*, of which he has seen many (Rose, No. 19627). It fruited in the New York Botanical Garden in 1920. The plant is called for de baile or flower of the ball.

An *Epiphyllum* grows in the lowlands of Ecuador which we have tentatively referred here, although we have never seen its flowers or fruits. Dr. Rose collected it below Huigra, September 8, 1918 (No. 22614), and again above Santa Rosa near Limón Playó, October 17 (No. 23493).

Cactus phyllanthus of Linnaeus (Sp. Pl. 469. 1753) and *Epiphyllum phyllanthus* Haworth (Syn. Pl. Succ. 197. 1812) both contain references not only to this species but to *Epiphyllum phyllanthoides* also.

The variety *columbiensis* was described by both Weber and Schumann with a flower-tube only 6 cm. long.

Cereus phyllanthus marginatus Parmentier is mentioned by Lemaire (Cact. Gen. Nov. Sp. 76. 1839) but not described.

Illustrations: Contr. U. S. Nat. Herb. 16: pl. 68, as *Epiphyllum gaillardae*; Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 10, f. 1, as *Cereus phyllanthus*; Petiver, Gazoph. Dec. pl. 59, f. 10. 1709, as *Heliotropium*, etc.; Dillenius, Hort. Elth. pl. 64, as *Cereus scolopendrii*, etc.; De Candolle, Pl. Succ. Hist. pl. 145; Vellozo, Fl. Flum. 5: pl. 33 (except flower), as *Cactus phyllanthus*; Monatsschr. Kakteenk. 2: 73, as *Phyllocactus phyllanthus*; Martius, Fl. Bras. 4²: pl. 44.

Figure 194 is from a photograph of a fruiting branch borne on the specimens obtained by Dr. Rose in Brazil in 1915; figure 195 shows a seedling with its two large cotyledons, grown from seeds sent by Mrs. D. D. Gaillard from Panama.




FIG. 195.—Seedling of *Epiphyllum phyllanthus*. $\times 0.6$.

2. *Epiphyllum oxypetalum* (De Candolle) Haworth, Phil. Mag. 6: 109. 1829.

Cereus oxypetalus De Candolle, Prodr. 3: 470. 1828.

Cereus latifrons Pfeiffer, Enum. Cact. 125. 1837.

Phyllocactus oxypetalus Link in Walpers, Repert. Bot. 2: 341. 1843.

Phyllocactus latifrons Link in Walpers, Repert. Bot. 2: 341. 1843.

Phyllocactus grandis Lemaire, Fl. Serr. 3: 255b. 1847.

Phyllocactus guyanensis Brongnart in Labouret, Monogr. Cact. 416. 1853.

Epiphyllum acuminatum Schumann in Martius, Fl. Bras. 4²: 222. 1890.

Phyllocactus acuminatus Schumann, Gesamtb. Kakteen 213. 1897.

Phyllocactus purpusii Weingart, Monatsschr. Kakteenk. 17: 34. 1907.

Epiphyllum grande Britton and Rose, Contr. U. S. Nat. Herb. 16: 257. 1913.

Plants stout, 3 meters long or more, much branched; branches flat and thin, 10 to 12 cm. broad, long-acuminate, deeply crenate; flowers opening in the evening, drooping and limp after anthesis, fragrant; tube of flower 13 to 15 cm. long, rather stout, red, about 1 cm. thick, bearing distant narrow scales about 10 mm. long; outer perianth-segments narrow, reddish to amber, 8 to 10 cm. long; inner perianth-segments oblong, white; stamens numerous, white; style white, thick, 20 cm. long; stigma-lobes numerous, cream-colored, entire.

Type locality: Mexico.

Distribution: Mexico and Guatemala, Venezuela, and Brazil. Widely cultivated in the tropics and doubtless an escape in many places.

This species has long been cultivated and has always been a great favorite on account of the ease with which it is grown and the abundance of large flowers it furnishes. These begin to open in the early evening and are perfect about midnight.

According to Mr. Pittier, this plant is known as flor de baile in Venezuela.

Epiphyllum latifrons Zuccarini (Pfeiffer, Enum. Cact. 125. 1837) was given as a synonym of *Cereus latifrons* when that name was first published.

The name *Cactus oxypetalus* Mocino and Sessé was the first one given to this plant, but De Candolle (Prodr. 3: 470. 1828) published the species as a *Cereus*, citing the above name as a synonym.

The following hybrids were listed by Labouret (Monogr. Cact. 429. 1853) between *Phyllocactus latifrons* and some other species of *Epiphyllum* or related genera; *Phyllocactus longipes*, *P. lothii*, *P. londonii*, *P. macquianus*, *P. maelenii*, *P. maurantianus*, *P. mexicanus*, *P. roseus albus*, *P. roseus superbus*, *P. selloi*, *P. smoli*, and *P. smithii*.

Illustrations: Monatsschr. Kakteenk. 17: 35, as *Phyllocactus purpusii*; Meehans' Monthly 12: 188, as *Epiphyllum latifrons*; Mém. Mus. Hist. Nat. Paris 17: pl. 14, as *Cereus oxypetalus*; Förster, Handb. Cact. ed. 2. 849. f. 112, as *Phyllocactus oxypetalus*; Rother, Praktischer Leitfaden Kakteen 93; Monatsschr. Kakteenk. 20: 123, as *Phyllocactus grandis*; Martius, Fl. Bras. 4²: pl. 45, as *Epiphyllum acuminatum*; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 59, D, as *Phyllocactus acuminatus*; Gard. Chron. 1849: 788; Pfeiffer and Otto, Abbild. Beschr. Cact. 1: pl. 10, f. 2, 3; Curtis's Bot. Mag. 67: pl. 3813, as *Cereus latifrons*; Gartenwelt 10: 560; Cact. Journ. 1: 55; Goebel, Pflanz. Schild. 1: pl. 2, f. 6; Schelle, Handb. Kakteenk. 209. f. 139; 210. f. 10, as *Phyllocactus latifrons*.




FIG. 196.—*Epiphyllum pumilum*. $\times 0.5$.

3. *Epiphyllum pumilum* Britton and Rose, Contr. U. S. Nat. Herb. 16: 258. 1913.

Phyllocactus pumilus Vaupel, Monatsschr. Kakteenk. 23: 117. 1893.

At first erect or ascending but often becoming pendent, sometimes 5 meters long; main stems terete; branches of two types; some of them elongated, 8 to 15 dm. long, terete, whip-like, sometimes becoming flattened at tip; some broad and flattened, rarely 3-winged, except at base, usually acute or acuminate, 1 to 6 dm. long, 3 to 8.5 cm. broad, becoming thick when old, the margin remotely toothed; flowers small for the genus; tube . to 6 cm. long, greenish white to reddish, bearing a few very small ascending and appressed reddish scales; outer perianth-segments linear, greenish or reddish, acute; inner perianth-segments white, lanceolate, acuminate, 3 to 4 cm. long; stamens in two groups; style slender, white, oblong, 4 to 7 cm. long, 2 to 2.5 cm. in diameter; fruit brilliant cerise when ripe, 5 to 7-ridged, bearing a few very small reddish ascending scales; pulp of fruit white, edible, sweet; seeds minute, jet-black.

Type locality: Guatemala.

Distribution: Lowlands of Guatemala.

This species has frequently been collected in Guatemala and is usually called *Epiphyllum pittieri*, which it somewhat resembles in the size of the flower, but the style is always white.

The flowers are night-blooming and sweet-scented. The fruit is much sought after by the Guatemalan Indians, who call it pitahaya.

The above description is based on living specimens, full notes, and drawings, furnished by Harry Johnson, a very keen observer, at one time stationed in Guatemala.

Figure 196 is copied from pencil sketches made by Mr. Harry Johnson at Chamá, Alta Verapaz, Guatemala, in 1920.

4. *Epiphyllum caudatum* Britton and Rose, Contr. U. S. Nat. Herb. **16**: 256. 1913.

Phyllocactus caudatus Vaupel, Monatsschr. Kakteenk. **23**: 116. 1913.

Old stems terete and slender; lateral branches elongated-lanceolate, cuneately narrowed at base into a terete stalk, long-acuminate, 15 to 20 cm. long, 3 to 4 cm. wide, the margins low-undulate; flowers white, the tube slender, about 7 cm. long; inner perianth-segments about 6 cm. long; ovary and most of the flower-tube quite naked.




FIG. 197.—*Epiphyllum caudatum*.

Type locality: Near Comaltepec, Oaxaca, Mexico, altitude 540 to 900 meters.

Distribution: Known only from the type locality.

We have seen no specimens of this species except the type, but Dr. B. P. Reko, under date of June 28, 1919, wrote that he had seen the plant not only at Comaltepec, but at other places in the Sierra Juarez.

A plant sent from Chiapas, Mexico, by Dr. C. A. Purpus in 1920 has joints with similar acuminate tips, but the margins are indented. We do not know its flowers.

Figure 197 is from a photograph of the type specimen.

5. *Epiphyllum darrahii* (Schumann) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 256. 1913.

Phyllocactus darrahii Schumann, Gesamtb. Kakteen Nachtr. 69. 1903.

Stems much branched, often terete and woody below; joints rather thick, 2 to 3 dm. long, 3 to 5 cm. wide, deeply lobed, sometimes nearly to the midrib, the lobes usually obtuse; tube of flower 9 cm. long, somewhat curved, greenish; scales on tube and ovary small, linear, green, appressed; outer perianth-segments 10, linear, spreading or reflexed, acute, 4 cm. long, lemon-yellow; inner perianth-segments pure white, nearly as long as the outer, broader and more erect,



M. E. Eaton del.

1. Flowering plant of *Epiphyllum darrahii*.
2. Top of flowering plant of *Epiphyllum pittieri*.

A. Hoen & Co. Baltimore

short acuminate; filaments white, nearly as long as the perianth-segments; style overtopping the stamens, pure white; stigma-lobes 8, linear.

Type locality: Mexico.

Distribution: Mexico, but range unknown.

This species was named for Charles Darrah of Heaton Mersey near Manchester, England (1844-1903). His large and valuable collection of succulents, especially cacti, was presented to the Corporation of Manchester by his widow and family and is now housed in specially constructed houses in Alexander Park. In 1908 the late Robert Lamb published a catalogue of 129 pages of this collection.

The plant is cultivated in Mexico and is much prized as a potted plant for the patio; one of these was obtained by Dr. Rose in Ixmiquilpan in 1905 (No. 9091). Living specimens were sent home and these have repeatedly flowered in Washington and New York. It flowers abundantly, its blossoms giving off a most delicious honeysuckle-like fragrance; we have seen no specimens of wild plants.




FIG. 198.—*Epiphyllum darrahii*.

Illustration: Blühende Kakteen 2: pl. 91, as *Phyllocactus darrahii*.

Plate xvi, figure 1, shows the plant in flower, collected by Dr. Rose in Mexico in 1905, which flowered in the New York Botanical Garden in September 1917. Figure 198 is from a photograph of the plant collected by Dr. Rose at Ixmiquilpan, Mexico, in 1905 which afterwards flowered in Washington.

6. *Epiphyllum anguliger* (Lemaire) Don in Loudon, *Encycl. Pl. ed. 3.* 1380. 1855.

Phyllocactus anguliger Lemaire, *Jard. Fleur.* 1: pl. 92. 1851.

Phyllocactus serratus Brongnart in Labouret, *Monogr. Cact.* 417. 1853.

Much branched; stems and lower branches terete; upper branches flattened with deeply toothed margins, rather fleshy; areoles small, usually felted and sometimes bearing 1 or 2 white bristles; flower-tube stout, without scales, about 8 cm. long; outer perianth-segments brownish yellow, inner perianth-segments white, oblong, acuminate, about 5 cm. long; style slender, white.

Type locality: Near Matanejo, Mexico.

Distribution: Central and southern Mexico.

We know the species only from cultivated plants. When not in flower it is difficult to distinguish it from *Epiphyllum darrahii*.

This plant was first distributed by the Horticultural Society of London, which obtained it from the collector, T. Hartweg, in 1846, from southern Mexico, where it was found growing on oak trees.

Phyllocactus angularis occurs in the index of Labouret's Monograph (511), credited to Lemaire, and also is listed in the Index Kewensis. It may have been a manuscript name for this species.

Illustrations: Lemaire, Jard. Fleur. 1: pl. 92; Lindley and Paxton, Fl. Gard. 1: pl. 34; Curtis's Bot. Mag. 85: pl. 5100; Dict. Gard. Nicholson 3: f. 134; Amer. Gard. 11: 538; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 24; Cycl. Amer. Hort. Bailey 1: f. 306; Palmer, Cult. Cact. 167; Watson, Cact. Cult. 48. f. 11; ed. 3. f. 9; Florialia 42: 377, as *Phyllocactus anguliger*.

7. *Epiphyllum grandilobum* (Weber) Britton and Rose, Contr. U. S. Nat. Herb. 16: 257. 1913.

Phyllocactus grandilobus Weber, Bull. Mus. Hist. Nat. Paris 8: 464. 1902.

Branches bright green, very large, up to 25 cm. broad with the margins deeply lobed and with a thick midvein and obtuse or rounded apex; lobes rounded, 3 to 5 cm. long; flowers described as large, white, opening at night; fruit red without.

Type locality: La Hondura, Costa Rica.

Distribution: Costa Rica.

Weber speaks of this as a very remarkable species of which he had not seen flowers or fruit. His description was based on specimens collected by Wercklé in 1900 and also by Pittier in 1905.

Specimens of the type collection were obtained by Mr. Wm. R. Maxon from A. Brade in Costa Rica in 1906 (No. 13), but these have never flowered. In the New York Botanical Garden is a small specimen received from Wercklé in 1902 as *Epiphyllum grandilobum*; this shows one very deep lobe; a young joint shows shallow crenations and suggests *E. macropterum*. A plant of this relationship was collected by Mr. Pittier in Panama in 1911 (No. 4229) and is now growing in Washington, but has not flowered.

We believe that *Phyllocactus macrolobus* of Schumann's Keys belongs here, the specific name in error for *grandilobus*.

8. *Epiphyllum crenatum* (Lindley) G. Don in Loudon, Encycl. Pl. ed. 3. 1378, 1855.

Cereus crenatus Lindley in Edwards's Bot. Reg. 30: pl. 31. 1844.

*Phyllocactus crenatus** Lemaire, Hort. Univ. 6: 87. 1845.

Phyllocactus caulorrhizus Lemaire, Jard. Fleur. 1: Misc. 6. 1851.

Epiphyllum caulorrhizum G. Don in Loudon, Encycl. Pl. ed. 3. 1380. 1855.

Old stems woody and terete; branches glaucous, often rooting at the tips, rather stiff, 2 to 3 cm. broad, obtuse, erect, at least at first, with large deep crenations, cuneate at base, the midrib thick; areoles at base of stem and branches often bearing hairs or small bristles; flowers very fragrant, rather large, the limb 10 to 12 cm. broad, cream-colored to greenish yellow, tube 10 to 12 cm. long, slender, bearing linear scales 2 to 3 cm. long; inner perianth-segments oblanceolate, 6 cm. long; filaments yellow; style white; stigma-lobes narrow; ovary scaly, some of the scales 2 cm. long, somewhat spreading.

Type locality: Honduras.

Distribution: Honduras and Guatemala.

This species has long been a favorite with gardeners, and many hybrids with it have been produced; the flowers, which are delicately fragrant, are diurnal and remain expanded for several days.

Among hybrids with other species are *Phyllocactus crenatus amaranthinus*, *P. elegans*, *erleri*, *haageanus*, *lateritius*, *roseus*, *splendens*, *superbus*, and *vogelii*.

Illustrations: Edwards's Bot. Reg. 30: pl. 31, as *Cereus crenatus*; Blühende Kakteen 3: pl. 180, as *Phyllocactus crenatus vogelii*; Gartenflora 40: pl. 1347; Garten-Zeitung 4: 182. f. 42,

* This name was also published by Walpers in 1843 (Repert. Bot. 2. 820).



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. End of branch of *Epiphyllum macropterum*.
2. Base of branch of same.

No. 4; Rother, Praktischer Leitfaden Kakteen 80, as *Phyllocactus crenatus*; Loudon, Encycl. Pl. ed. 3. 1379. f. 19401.

Figure 199 is from a photograph showing the base and tip of a branch of this species sent from Guatemala.




FIG. 199.—*Epiphyllum crenatum*.

9. *Epiphyllum macropterum* (Lemaire).

- Phyllocactus macropterus* Lemaire, Illustr. Hort. 11: Misc. 7. 1864.
Phyllocactus thomsonianus Schumann, Monatsschr. Kakteenk. 5: 6. 1895.
Phyllocactus costaricensis Weber, Bull. Mus. Hist. Nat. Paris 8: 463. 1902.
Phyllocactus macrocarpus Weber, Bull. Mus. Hist. Nat. Paris 8: 464. 1902.
Epiphyllum costaricense Britton and Rose, Contr. U. S. Nat. Herb. 16: 256. 1913.
Epiphyllum thomsonianum Britton and Rose, Contr. U. S. Nat. Herb. 16: 259. 1913.

Plants up to 2 meters long, the joints weak, sometimes 10 cm. broad, thin, their margins horny; areoles distant (to 6 cm. apart) along the slightly indented margins; flower very large for genus, long, curved as in *Epiphyllum oxypetalum*; scales of ovary small, green, spreading, with long hairs in their axils; scales on tube longer (10 to 12 mm. long), less spreading but similar to those on ovary, acute; outer perianth-segments narrow, salmon-colored or with yellow tips, 10 cm. long; inner perianth-segments pure white, 8 to 9 cm. long, 2 to 3 cm. broad; tube of the flower 10 to 12 cm. long; throat 5 to 6 cm. long, funnelform, narrow below, 3 cm. broad at top; stamens lemon-yellow, slender, in 2 definite clusters, a single continuous row at top of throat, the second cluster scattered all over throat except for intervals of 2 cm. below upper one; style stout, 20 cm. long, pure white.

Type locality: Not cited.

Distribution: Costa Rica.

According to Mr. Fraile, the flower always comes out on the under side of the joint and lies appressed to it, instead of standing out free from it as in other species of the genus.

A vigorous plant in greenhouse cultivation but it flowers only sparingly.

Illustrations: Monatsschr. Kakteenk. 5: pl. [1]; Blühende Kakteen 1: pl. 41, as *Phyllocactus thomsonianus*.

Plate xvii, figure 1, shows a branch of a plant sent by Dr. Wm. R. Maxon from San José, Costa Rica, which flowered in the New York Botanical Garden in 1912; figure 2 shows

the base of the branch. Figure 200 is from a photograph showing the top and base of a joint.

- 10. *Epiphyllum lepidocarpum*** (Weber) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 257. 1913.
Phyllocactus lepidocarpus Weber, Bull. Mus. Hist. Nat. Paris **8**: 462. 1902.

Old and lower part of stems woody, cylindric; upper branches usually flattened, sometimes 3-winged, thickish, but not very stiff, 2 to 3 cm. wide; margins cut "stair-like," the areole closed by a small scale bearing in its axil short wool and a few bristles; flowers 20 cm. long, white and night-blooming; stamens in 2 rows; style white; fruit 9 cm. long by 4 cm. in diameter, violet-red, covered with long scales, at first erect, but finally becoming reflexed; flesh described as white, * acidulous, somewhat agreeable to the taste.




FIG. 200.—*Epiphyllum macropterum*.

Type locality: Near Cartago, Costa Rica.


Distribution: Known only from the type locality.

Our description is based on that of M. Weber. The very scaly fruit should be characteristic, but plants received from Costa Rica under the name *Phyllocactus lepidocarpus* produced smooth fruits at the New York Botanical Garden.

- 11. *Epiphyllum pittieri*** (Weber) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 258. 1913.
Phyllocactus pittieri Weber, Dict. Hort. Bois 957. 1898.

Stem usually terete below, much divided, 2 to 3 meters long; branches flat and thin, mostly cm. wide or less, the margins coarsely toothed; flowers rather small, the tube about 8 cm. long, white to greenish white, bearing a few red, ascending scales; outer perianth-segments 4 to 4.5 cm. long, narrow, yellowish green, or some of the lower ones tinged with red, acute; inner perianth-segments white, a little shorter than the outer; stamens white, erect, in 2 series, longer than the style; style white above, red or purplish below; ovary with a few red scales; fruit dark red, 2 cm. long; seeds dull black.

*Mr. Wercklé, who discovered this species, states in a letter (September 22, 1921) that the flesh is crimson.



Epiphyllum pittieri, from Costa Rica.

Type locality: Costa Rica.

Distribution: Costa Rica.

This species is an abundant bloomer, flowering in cultivation usually in January but also at other times of the year; its flowers are the smallest of the genus.

Plate xvi, figure 2, shows a flowering branch from the specimen sent by Mr. Pittier from Zent, Costa Rica, in 1904; plate xviii shows another plant of the same collection which flowered in Washington.

12. *Epiphyllum guatemalense* Britton and Rose, Contr. U. S. Nat. Herb. **16**: 257. 1913.

Phyllocactus guatemalensis Vaupel, Monatsschr. Kakteenk. **23**: 116. 1913.

Plant rather stout, in cultivation a meter long or longer; old stem woody, with gray bark, terete; branches green, flat, 4 to 8 cm. broad, narrowed at base and there terete, coarsely crenate, obtuse at apex flower-bud pointed flowers nocturnal, about 2.8 cm. long; tube about 1.5 cm. long,




FIG. 202.—*Epiphyllum guatemalense*.

straight or nearly so, green or yellowish green, somewhat angled, at least below, bearing only a few red-tipped scales; inner central part of tube densely pilose; outer perianth-segments scale-like with red reflexed tips; inner pure white, narrow, 8 to 9 cm. long, acuminate; stamens borne on whole surface of rather short throat and therefore in more than one series; filaments pure white; style 2.5 cm. long, somewhat glossy, orange; stigma-lobes orange; ovary pale, bearing a few spreading scales.

Type locality: Guatemala.

Distribution: Guatemala, but range unknown.

Two very distinct forms occur in this species which are hard to explain. They are so different that it seemed at first they must represent two distinct species, as they occur on separate plants. In one (it may be simply the juvenile form) the joints are rather thin and broad (to 8 cm. broad), the margins soft, with low broad undulations separated by a narrow, nearly closed sinus; in the other (it may perhaps be the adult form) the joints are stiff and narrow, the margins horny, the undulations with an open triangular sinus.

Illustration: Contr. U. S. Nat. Herb. **16**: pl. 78.

Figure 201 is from a photograph of the type plant.

- 13. *Epiphyllum strictum*** (Lemaire) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 259. 1913.
Phyllocactus strictus Lemaire, Illustr. Hort. **1**: Misc. 107. 1854.

Plant up to 2 meters long; joints linear, green, 5 to 8 cm. broad, coarsely serrate, stiff; tube of flower 13 to 15 cm. long, slender, green, bearing a few distant scales 8 to 12 mm. long; outer perianth-segments pink, the inner white, narrow, acuminate, 6 to 8 cm. long; filaments white; style pink or red; stigma-lobes yellow; fruit globose, 4 to 5 cm. in diameter; seeds black.

Type locality: Cuba, but the plant was grown there from seed.

Distribution: Southern Mexico and Guatemala to Panama.

The plant was found in the wild state in Honduras by Mr. Percy Wilson in 1902. All the other specimens studied by us are from cultivated plants. The species is common in collections.

Illustrations: Schumann, Gesamtb. Kakteen f. 41; Monatsschr. Kakteenk. **6**: 183; Thomas, Zimmerkultur Kakteen 18, as *Phyllocactus strictus*.




FIG. 202.—*Epiphyllum stenopetalum*.

- 14. *Epiphyllum stenopetalum*** (Förster) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 259. 1913.
Phyllocactus stenopetalus Förster, Handb. Cact. 441. 1846.

Described as with the habit of *Epiphyllum latifrons* but with different flowers, these delicately fragrant; flower-tube 2 to 5 cm. long, bearing small, spreading, rose-colored scales; outer perianth-segments rose-colored to reddish green; inner perianth-segments white, elongated, linear (to 8 cm. long, very narrow, 4 to 7 mm. broad), spreading or recurved; stamens somewhat exserted; style slender, pink or purplish; stigma-lobes 12 to 14, yellow; fruit unknown.

Type locality: Not cited.

Distribution: Oaxaca, Mexico.

This plant is a night-bloomer but the flowers are late in closing, sometimes remaining partially open as late as 9 o'clock in the morning.

The above description is compiled from that of Salm-Dyck with reference to a plant at the New York Botanical Garden, received from Paris in 1909.

It resembles *E. strictum* but the joints are more flexible and broader and it has somewhat larger flowers than that species; we have a herbarium specimen identified by Schumann which was collected by P. Sintenis from a cultivated plant grown in Porto Rico.

In 1911 C. Conzatti sent us from Coyula, Cuicatlán, Oaxaca, cuttings of what we now take to be this species. These grew into vigorous plants 3 meters long and flowered in Washington in 1921 and 1922.

Illustration: Goebel, Pflanz. Schild. 1: f. 56, as *Phyllocactus stenopetalus* (seedling).

Figure 202 is from a photograph showing the top and base of a branch from Professor Conzatti's plant.

15. *Epiphyllum cartagense* (Weber) Britton and Rose, Contr. U. S. Nat. Herb. 16: 256. 1913.

Phyllocactus cartagensis Weber, Bull. Mus. Hist. Nat. Paris 8: 462. 1902.

Phyllocactus cartagensis refractus Weber, Bull. Mus. Hist. Nat. Paris 8: 462. 1902.

Phyllocactus cartagensis robustus Weber, Monatsschr. Kakteenk. 15: 180. 1905.

Plants 2 to 3 meters long, usually more or less flattened in the lower and older parts; joints short or elongated, 4 to 5 cm. broad, coarsely toothed or crenate, green; flowers opening at night, the slender tube 10 to 15 cm. long, reddish, bearing a few short distant scales; outer perianth-segments pink to yellowish; inner segments 5 to 7 cm. long, white; stamens in one series; filaments white; style pink to white; stigma-lobes yellow; fruit oblong, 7 to 8 cm. long, 3 cm. in diameter, red without, white within.

Type locality: Near Cartago, Costa Rica.

Distribution: Costa Rica.

A species apparently composed of several races, differing in margins of the joints, in size of flowers, and in color of style. It is called in Costa Rica platanillo de monte.

16. *Epiphyllum hookeri* Haworth, Phil. Mag. 6: 108. 1829.

Cereus hookeri Link and Otto, Cat. Sem. Hort. Berol. 1828.

Cereus marginatus Salm-Dyck, Hort. Dyck. 340. 1834. Not De Candolle, 1828.

Phyllocactus hookeri Salm-Dyck, Cact. Hort. Dyck. 1841. 38. 1842.

Plants usually 2 to 3 meters long, but sometimes 7 meters long; joints 5 to 9 cm. broad, rather thin, light green, deeply crenate; flowers inodorous, the tube slender, 11 to 13 cm. long, greenish, bearing a few narrow, slightly spreading, rose-tipped scales; outer perianth-segments narrow, greenish pink, sometimes rose-colored at tip, the inner pure white, narrow, 5 cm. long; stamens in a single series, attached at top of throat; filaments white; style carmine, except yellowish base and pinkish top, smooth in upper half, papillose in lower half; stigma-lobes yellow; ovary green, somewhat angled, 2 cm. long, bearing a few small spreading scales; fruit oblong, 8 cm. long, red, somewhat angled, bearing a few scattered scales; seeds numerous, black, shining, reniform.

Type locality: Cited as Brazil, presumably in error.

Distribution: Tobago, Trinidad, and northern Venezuela.

This plant when it first flowered in cultivation in 1826 was taken for *Cactus phyllanthus* and was so figured and described in the Botanical Magazine, but it was soon discovered to be very different from that species.

While Brazil is cited as the type locality for this species we have seen no specimens from any point south of Venezuela. The plant is and has been widely cultivated in tropical America, commonly under the erroneous name, *Epiphyllum phyllanthus*. In Trinidad it forms great masses on trees and on coastal cliffs, ascending the trees to a length of 10 meters or more, branching profusely, and is very floriferous.

Phyllocactus marginatus Salm-Dyck (Cact. Hort. Dyck. 1844. 37. 1845) doubtless belongs here.

Illustrations: Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 5, as *Cereus hookeri*; Curtis's Bot. Mag. 53: pl. 2692; Loudon, Encycl. Pl. 413. f. 6901, as *Cactus phyllanthus*; Addisonia 5: pl. 192.

Plate XIX shows a flowering branch from a specimen sent by W. E. Broadway from the Island of Tobago in 1909.

HYBRIDS.

EPIPHYLLUM ACKERMANNII Haworth, Phil. Mag. 6: 109. 1829.

Cactus ackermannii Lindley in Edwards's Bot. Reg. 16: pl. 1331. 1830.

Cereus ackermannii Otto in Pfeiffer, Enum. Cact. 123. 1837.

Phyllocactus ackermannii Salm-Dyck, Cact. Hort. Dyck. 1841. 38. 1842.

Branches weak, flat, and thin with crenate margins; areoles felted, often bristly or with weak spines, especially on the young growth; flowers day-blooming, very large, sometimes 1.5 to 2 dm. broad, crimson; inner perianth-segments oblong, acute; filaments long, weak, declined; style more or less declined, pinkish; stigma-lobes white; ovary more or less bristly.

Type locality: Mexico.

Distribution: Mexico.

This species was originally described as from Mexican plants sent to Haworth from Ackermann and, supposedly, from wild plants, but the general belief now is that the plant is of hybrid origin. The flowers are so much like those of *Heliocereus* that this genus probably furnished one of its parents (see Botanical Magazine, pl. 3598).

On the other hand, E. A. Goldman collected in Chiapas a series of specimens which seems to represent more than one species, but all the flowers are similar to those of *Epiphyllum ackermannii* and one of the specimens may represent the wild state of that species. The plants all have flat joints bearing clusters of spines in their areoles.

Many garden varieties and artificial hybrids have been obtained from this plant, some described under English and others under Latin names.

Illustrations: Edwards's Bot. Reg. 16: pl. 1331, as *Cactus ackermannii*; Curtis's Bot. Mag. 64: pl. 3598, as *Cereus ackermannii*; Blühende Kakteen 1: pl. 49; Cycl. Amer. Hort. Bailey 3: f. 1773; Dict. Gard. Nicholson 3: f. 133; Karsten, Deutsche Fl. 887. f. 501, No. 6; ed. 2. 2: 456. f. 605, No. 6; Förster, Handb. Cact. ed. 2. 841. f. 111; Rümpler, Sukkulente 149. f. 81; Watson, Cact. Cult. 47. f. 10; Rother, Praktischer Leitfaden Kakteen 97; ed. 3. f. 8; Amer. Gard. 11: pl. opp. 445; Gartenflora 32: 374, as *Phyllocactus ackermannii*; Loudon, Encycl. Pl. 1202. f. 17368 Encycl. Britannica ed. 11. 4: 926. f. 3, as *Phyllocactus*; Rev. Hort. 1861: 226. f. 44; Stand. Cycl. Hort. Bailey 2: f. 1402.

Cactus hybridus was described and illustrated by P. C. Van Gél (Sert. Bot. 1: pl. 115. 1832). He states that it is known in Great Britain as *C. ackermannii*.

EPIPHYLLUM HYBRIDUM Hortus in Pfeiffer, Enum. Cact. 121. 1837.

This was given as a synonym of *Cereus speciosissimus lateritius*, which is briefly mentioned in volume 2 (p. 128) of this work.

EPIPHYLLUM JENKENSII G. Don, Gen. Hist. Dichl. Fl. 3: 170. 1834.

Epiphyllum speciosum jenkinsii G. Don in Loudon, Encycl. Pl. ed. 2. 1202. 1841.

This plant is an artificial hybrid raised from *Heliocereus speciosissimus*, impregnated by the pollen of *Epiphyllum phyllanthoides*; it has branches 3-angled at base but flattened above, with areoles very prominent and spiny; flowers large, 10 cm. broad and deep scarlet; fruit nearly globular, purple, 2.5 cm. in diameter, its areoles bearing a few spines and bristles. We have had it to flower and fruit in cultivation.


EPIPHYLLUM SPLENDIDUM Paxton, Mag. Bot. 1: 49. 1834.

Cereus splendidus Steudel, Nom. ed. 2. 1: 336. 1840.

Epiphyllum aitoni Steudel, Nom. ed. 2. 1: 561. 1840.

Epiphyllum hitcheni Steudel, Nom. ed. 2. 1: 561. 1840.

We know this plant only from a colored illustration (Paxton, Mag. Bot. 1: pl. facing 49). The flower is very large, 10 inches broad, red, tinged with orange; flower-tube much shorter than limb, and suggests a relationship with *Epiphyllum ackermannii*. Branches flat and strongly crenate. It is



Epiphyllum hookeri, from Tobago, West Indies.

said to be a native of Mexico, but probably is of hybrid origin. It is described as having one of the largest flowers among cacti, rivaling *Selenicereus grandiflorus* and *Helicocereus speciosus*.

Epiphyllum splendens Hortus (Ann. Fl. Porn. 343, 1839) is referred here by the Index Kewensis. It is, however, described on page 345 and illustrated on plate 44. This illustration is very different from that of Paxton.

PHYLLOCACTUS ALBUS GRANDIFLORUS.

Illustration: Cact. Journ. 1: 37.

PHYLLOCACTUS ALBUS SUPERBUS.

Illustration: Blanc, Cacti 88, No. 2511.

PHYLLOCACTUS COOPERI Regel, Gartenflora 33: 218. 1884.

This is a hybrid between *Epiphyllum crenatum* and *Selenicereus grandiflorus*. It has large yellowish flowers.

Illustrations: Gartenflora 33: pl. 1176, as *Phyllocactus crenato grandiflorus*; Cassell's Dict. Gard. 2: 192.

PHYLLOCACTUS EREBUS.

This is a large red-flowered hybrid.

Illustration: Blühende Kakteen 3: pl. 160.

PHYLLOCACTUS HAAGEI.

This is doubtless a garden hybrid related to *Epiphyllum ackermannii*. It has large flowers, 12.5 cm. broad, at first flesh-colored, becoming carmine.

Illustrations: Dict. Gard. Nicholson 4: 590. f. 8; Watson, Cact. Cult. 54. f. 13.

PHYLLOCACTUS HIBRIDUS GORDONIANUS.

Illustration: Blühende Kakteen 1: pl. 36.

PHYLLOCACTUS HIBRIDUS WRAYI.

This is said to be a cross between *Selenicereus grandiflorus* and *Epiphyllum crenatum*.

Illustration: Blühende Kakteen 2: pl. 6.

PHYLLOCACTUS HILDMANNII.

Illustration: Gartenflora 44: pl. 1421, f. 2.

PHYLLOCACTUS MARSUS.

Illustration: Dict. Gard. Nicholson Suppl. 598. f. 631.

PHYLLOCACTUS PFERSDORFFII.

Illustrations: Cact. Journ. 1: 38; Rümpler, Sukkulente f. 85; Schelle, Handb. Kakteenk. 221. f. 144.

PHYLLOCACTUS ROSEUS GRANDIFLORUS Watson, Cact. Cult. 55. 1889.

This was figured and described by Watson with flowers 15 cm. long and broad, nodding and white (!); doubtless of hybrid origin; it may be the same as *Phyllocactus roseus grandisimus* (Monatsschr. Kakteenk. 19: 182. 1909).

Illustrations: Dict. Gard. Nicholson 4: 591. f. 59; Förster, Handb. Cact. ed. 2. 857. f. 117; Watson, Cact. Cult. 55. f. 14.

PHYLLOCACTUS RUESTII Weingart, Monatsschr. Kakteenk. 24: 123. 1914.

We have not seen this plant and Mr. Weingart, who described it, says that he does not possess either living or herbarium material but that it is still growing at Halle, Germany.

PHYLLOCACTUS TRIUMPHANS.

Illustration: Monatsschr. Kakteenk. 20: 3.

Epiphyllum speciosum lateritium Henslow (Loudon, Encycl. Pl. ed. 2. 1202. 1841), an English hybrid, produced in 1828, is described as having brick-colored flowers.

Phyllocactus tonduzii Weber is mentioned by Schumann (Monatsschr. Kakteenk. 10: 127. 1900).

Phyllocactus tuna is a name used by Wercklé (Monatsschr. Kakteenk. 15: 180. 1905) for a Costa Rican plant, without description.

Phyllocactus weingartii Berger (Monatsschr. Kakteenk. 30: 33. 1920) is related to *Epiphyllum ackermannii*.

Charles Simon in 1893 published a list of 62 names of *Epiphyllum*, most of which are undoubtedly hybrids and some are referable to *Zygocactus*. The following binomials and trinomials are in the usual Latin form for specific and varietal names and are not recorded elsewhere:

| | | | |
|-----------------|-----------------|----------------------|------------------------|
| album violaceum | grandiflorum | palidum roseum | translucens |
| amabile | grande superbum | purpureum | tricolor |
| aurantiacum | harrissoni | roseum | violaceum elegans |
| brasiliense | hercule | rubrum violaceum | violaceum grandiflorum |
| carminatum | latetium album | ruckerianum superbum | violaceum rubrum |
| cruentum | maximum | salmoneum marginatum | violaceum speciosum |
| gracilis | multiflorum | spectabile coccineum | violaceum superbum |

There are many Latin names of *Phyllocactus* in catalogues, representing hybrids. We give below only those which have been used more or less in general botanical works, either as binomials or trinomials in regular Latin form:

| | | | |
|----------------------|----------------|-------------------------|-------------------------|
| acutifrons | buestii | germania | lunus |
| agatha | campmannii | guentneri | mexicanus * |
| alatus | caparti | hamburgiensis | purpureus |
| albus superbiens | capelleanus | hauffii | ruelcheri |
| alexandrinae | carolus magnus | helenus | speciosissimus feltonii |
| amarantinus | castneri | hempelii | superbus |
| arnoldi | chico | hibridus | ulbrechtii |
| aurantiacus superbus | coccineus | incomparabilis minuatus | victoria-regia |
| belgicus | demouline | jenkinsonii | vogelii |
| bergen | dolores | kerthii | wippermannii |
| bleindlii | epirus | laarsenii | wrayi |
| boehmii | fuertii | lorenzii | zarka |

In 1897 Charles Simon, of Saint-Ouen, Paris, published a list of 370 names of *Phyllocactus*, most of which were probably hybrids. The following binomials and trinomials are in the usual Latin form for specific and varietal names and are not recorded elsewhere:

| | | | |
|----------------------|---------------------|-------------------------|------------------------|
| ackermannii hybridus | crenatus hirsutis | funkii | lorentzii |
| ackermannii major | crenatus lateralis | gloriosus | ludmani |
| alatus major | crenatus latifolius | gordonianus | ludwigi |
| albus grandidissimus | crenatus luteus | grandidissimus | maigretii |
| albus perfectus | crenatus ruber | grandiflorus | magnificus |
| albus superbissimus | crispielsi | grandiflorus albiflorus | makoyi |
| amabilis | curtissi | grandiflorus ruber | mayanus |
| amabilis perfectus | dangeli | guebwillerianus | meyerianus |
| atrosanguineus | decumbens | guedeneyi | muehlenpfordtii |
| aurantiacus | deveauxi | hansii | mulhousiausus |
| bergei | dieffenbachianus | havermansii | multiflorus |
| billiardieri | dumoulini | hitchensis | neubertii |
| binderi | edwardsii | ignescens | niedtii |
| blindii | elegans | jenkinsonii superbus | niger |
| boliviensis | erectus perfectus | johnsonii | nitens |
| bolivillerianus | erectus superbus | jordanis | nymphoea beata |
| bothii | ernesti | kampmannii | paraguayensis |
| brongnarti | erubescens | kermesimus magnus | pentneri |
| burmeisteri | fastuosus | kiardi | phyllantoides |
| colmariensis | feasti | kranzii | phyllantoides crenatus |
| colombiensis | felonis | krausei | potstachianus |
| courantii | feltoni | laetingii | poullertianus |
| crassuliefolius | floribundus | laloyi | pressleri |
| crenatus amarantinus | formosus | laudowi | |
| crenatus caulorhizus | franzii | leopoldii | |

* A hybrid referred by Index Kewensis to *Cereus mexicanus*.

pulcherrimus
quilliardetti
raveaudii
rebuti
reichei
reineckii
roseus carmineus
roseus carneus
roseus floribundus
roseus grandidissimus

roseus miniatus
roseus splendidus
roydii
ruber
ruber perfectus
ruber violaceus
sarniensis
schaffieri
schallerianus
schlimmi

schmidtii
sellowii
specillimus
speciosissimus
speciosissimus grandiflorus
speciosus roseus
splendens
splendidus
stenesi
superbissimus

tettani
tricolor
undiflorus
vandesii
vonhoffini
vitellinus
warszewiczii
wittmackianus

CACTUS ENSIFORMIS Biden, Gard. Chron, II. 20: 53. 1883.

This is evidently some *Epiphyllum* hybrid. It was sent to H. B. Biden from Manchester, England, in 1883 and flowered the same year. Its flowers were described as 6 inches across, white, richly scented, and remaining open for 3 days.

Cactus speciosus grandiflorus (Monatsschr. Kakteenk. 14: 11. 1904) is supposed to be some hybrid *Epiphyllum*.




FIG. 203.—Tip of branch with flower of *Disocactus biformis*. $\times 0.8$.

5. DISOCACTUS Lindley in Edwards's Bot. Reg. 31: pl. 9. 1845.

Disisocactus Kunze, Bot. Zeit. 3: 533. 1845.

Irregularly branching, spineless epiphytes, the stem terete; branches numerous, flattened; areoles marginal; flowers diurnal, borne near tips of branches, nearly regular; tube shorter than limb; perianth-segments few, elongated, spreading; ovary small, cylindric, elongated, bearing a few minute scales; fruit globular to ovoid, not at all angled.



FIG. 104.—*Disocactus biformis*.

Type species: *Cereus biformis* Lindley.

We recognize two species, both from Central America.

The name is from *δύς* twice, and *κάκτος* cactus, and was given because the perianth-segments of the inner and outer series were equal in the type specimens.

KEY TO SPECIES.

Slender with linear lateral branches, their margins slightly toothed; style and stamens about length of perianth-segments. 1. *D. biformis*
 Spreading with oblanceolate lateral branches, their margins crenate; style and stamens long-exserted 2. *D. eichlamii*

1. *Disocactus biformis* Lindley in Edwards's Bot. Reg. 31: pl. 9. 1845.

Cereus biformis Lindley in Edwards's Bot. Reg. 29 Misc. 51. 1843.

Disocactus biformis Kunze, Bot. Zeit. 3: 533. 1845.

Phyllocactus biformis Labouret, Monogr. Cact. 418. 1853.

Epiphyllum bifforme G. Don in Loudon, Encycl. Pl. ed. 3. 1378. 1855.

Plant 2 dm. long or longer; branches linear, 5 to 8 cm. long, 1 to 2 cm. broad, with serrate margins; flower-bud elongated, curved upward, pointed; tube of the flower about 1 cm. long, the segments 8 (rarely 9), magenta, about 3 cm. long, the outer 4 or 5 spreading or curved backward, linear, the inner 3 or 4 broader and more erect; stamens 10 to 12, slightly exserted, borne in 2 series at top of tube; style slender, purple; stigma-lobes 4, white; ovary short-oblong, green, somewhat tubercled, with a few areoles subtended by small ovate scales; fruit ovoid, 1.5 cm. long, turgid, wine-colored.

Type locality: Honduras. The species described from a garden specimen, introduced into England in 1839.




FIG. 205.—*Disocactus eichlamii*.

Distribution: Honduras and Guatemala.

We have had this plant under observation for a number of years. It is rather a shy bloomer with us, although we get one or two flowers each spring; the flowers open in the night or early morning and remain open all day; they begin to wither the second morning. The perianth-segments are more widely spreading in the morning than in the afternoon. The flower is almost horizontal and the tube proper is about the length of the ovary. The fruit matures very slowly. In 1920 we had a plant flower in April, but the fruit did not mature until July 8.

Illustrations: Förster, Handb. Cact. ed. 2. 876. f. 120; Rümpler, Sukkulenten f. 86, as *Disocactus biformis*; Blühende Kakteen 1: pl. 54; Curtis's Bot. Mag. 101: pl. 6156; Dict. Gard. Nicholson 3: f. 135; Monatsschr. Kakteenk. 9: 141; Watson, Cact. Cult. 50. f. 12, as *Phyllocactus biformis*; Loudon, Encycl. Pl. ed. 3. 1379. f. 19403, as *Epiphyllum bifforme*; Edwards's Bot. Reg. 31: pl. 9; Palmer, Cult. Cact. 175.

Plate xxxii, figure 2, shows a branch of a fruiting plant sent to Dr Rose by Robert Lamb of Manchester, England, in 1912. Figure 203 shows the flower of the same plant; figure 204 is from a photograph of the same plant in flower.

2. *Disocactus eichlamii* (Weingart) Britton and Rose, Contr. U. S. Nat. Herb. **16**: 259. 1913.*Phyllocactus eichlamii* Weingart, Monatsschr. Kakteenk. **21**: 5. 1911.

Branching near the base; branches oblong, 2 to 3 dm. long, 3 to 5 cm. broad, cuneate at base, obtuse, thickish, strongly crenate; flowers several at the uppermost areoles, slender, 4 cm. long; stamens and style exserted; stigma-lobes; fruit red, 1.5 cm. in diameter with white pulp; seeds 1.5 mm. long.

Type locality: Guatemala.

Distribution: Guatemala.

This plant we know only from the collection of F. Eichlam; living specimens were sent to Washington by him which soon afterward flowered, but these have since died. Eichlam wrote that the flowers were a brilliant red. The species was named for Federico Eichlam, who lived in Guatemala at the time of his death in 1911.

Illustration: Contr. U. S. Nat. Herb. **16**: pl. 79.

Figure 203 is from a photograph of the type plant in flower.

6. CHIAPASIA gen. nov.

An epiphytic spineless cactus, the branches flattened, crenate, with slender terete bases, the large flowers borne at upper areoles; perianth narrowly campanulate; tube about half as long as limb, bearing a few small triangular scales; segments about 8, linear, recurved, spreading; ovary ovoid, shorter than tube, also with a few small scales; filaments about 20, not longer than perianth-segments; stigma-lobes few.

Type species: *Epiphyllum nelsonii* Britton and Rose.

A monotypic genus, its name taken from that of the Mexican state in which it grows.

1. *Chiapasia nelsonii* Britton and Rose.

Epiphyllum nelsonii Britton and Rose, Contr. U. S. Nat. Herb. **16**: 257. 1913.

Phyllocactus nelsonii Vaupel, Monatsschr. Kakteenk. **23**: 116. 1913.

Phyllocactus chiapensis J. A. Purpus, Monatsschr. Kakteenk. **28**: 118. 1918.




FIG. 206.—*Chiapasia nelsonii*.

Branches 6 to 12 dm. long, slender and terete below, flat and thin above, 3 to 4 cm. broad; margin low, crenate; flowers light rose-red; tube 2 to 3 cm. long; segments about 6 cm. long, narrow, acute.

Type locality: Near Chicharras, Chiapas, Mexico, altitude 900 to 1,800 meters.

Distribution: Known only from Chiapas.

Our first description of this plant was based on an herbarium specimen, but considerable additional information is now known regarding its habit. A very fine plant was grown at Darmstadt by J. A. Purpus, a photograph of which we have, showing that the main branches are long and terete while the lateral branches are broad and thin, often pendent, with 1 to 3 flowers near the end; the flowers are horizontal, with the perianth-segments more or less recurved; the stamens and style are slender and long-exserted.

Illustration: Monatsschr. Kakteenk. 28: 119, as *Phyllocactus chiapensis*.

Figure 206 is from a photograph of the type specimen.

7. *ECCREMOCACTUS* Britton and Rose, Contr. U. S. Nat. Herb. 16: 261. 1913.

Plants epiphytic, pendent (erect or ascending in cultivation), several-jointed, the joints flat and thickish with spine-bearing marginal areoles or in cultivation often spineless; flowers solitary at upper areoles, funnellform, the short, nearly cylindric tube bearing small somewhat spreading scales, but no spines; perianth withering-persistent, its segments obtuse, rounded, or the innermost acutish; stamens and style white, included, slender, declinate; fruit carmine-red, oblong, with a few spineless areoles; seeds numerous, minute, black.

Type species: *Eccremocactus bradei* Britton and Rose.

Only one species is known, a native of Costa Rica. We have had the plant in cultivation for a number of years; it is a shy bloomer.

The generic name is from *ἐκκρεμής* hanging from, and *κακτος* cactus.

1. *Eccremocactus bradei* Britton and Rose, Contr. U. S. Nat. Herb. 16: 262. 1913.

Phyllocactus bradei Vaupel, Monatsschr. Kakteenk. 23: 118. 1913.

Epiphytic on trees; joints 15 to 30 cm. long, 5 to 10 cm. broad, light dull green, flat, but the central axis somewhat elevated on both sides, the margins shallowly crenate, with small spine-bearing areoles in the sinuses; spines solitary or in 2's or 3's, dark brown, 6 mm. long or less; flowers developing very slowly, 6 to 7 cm. long, slightly asymmetrical; outermost perianth-segments thick, shining, pinkish; outer ones oblong, thinner, pinkish white; inner perianth-segments oblong, obtuse, 3 to 3.5 cm. long; flower-tube 1 cm. long; throat broad, short, covered with stamens; filaments very slender, delicate, white, strongly declined; style slender, nearly white, slightly pinkish above, elongated, glabrous; stigma-lobes 8; ovary angled by the elongated tubercles; its areoles bearing a line of short hairs, subtended by thick ovate purple scales; seeds 1.5 mm. long.

Type locality: Cerro Turriwares, near Orotina (formerly Santo Domingo de San Mateo), Costa Rica.

Distribution: In dense forests at low altitudes, Costa Rica.

Our attention was first called to this plant by Dr. Maxon, who obtained specimens from Mr. Alfredo Brade in 1906; these bloomed in June 1911, but good flowers were not obtained. In 1913 Otón Jiménez sent specimens to Dr. Rose which flowered in 1918.

The flowers open in the night and are closed on the following morning. The branches of wild plants bear clusters of spines at the areole, but our cultivated plants are spineless and in the vegetative state resemble those of a turgid *Epiphyllum*. When the plant sent by Mr. Jiménez from Costa Rica (No. 905) flowered in 1921 seven flower-buds were produced from the seven uppermost areoles.

Illustration: Contr. U. S. Nat. Herb. 16: pl. 83.


Plate xx is from a photograph of a plant sent to Washington by Otón Jiménez in 1913, which flowered in May 1921.

8. *NOPALXOCHIA* gen. nov.

A flat-jointed, spineless epiphytic cactus; joints crenate, the rather large, short-funnelform, rose or red flowers, solitary at lateral marginal areoles; flower-tube about as long as limb, bearing several narrow scales; outer perianth-segments short, acute, reflexed or spreading; inner spreading or connivent, acute; stamens numerous.

Type species: *Cactus phyllanthoides* De Candolle.

A monotypic Mexican genus, the name taken from the Aztec of Hernández.



Eccremocactus bradei, from Costa Rica.

1. Nopalxochia phyllanthoides (De Candolle).

Cactus phyllanthoides De Candolle, Cat. Hort. Monsp. 84. 1813.

Cactus speciosus Bonpland, Descr. Pl. Rar. Malm. 8. 1813. Not Cavanilles, 1803.

Epiphyllum speciosum Haworth, Suppl. Pl. Succ. 84. 1819.

Cactus elegans Link, Enum. 2: 25. 1822.

Epiphyllum phyllanthoides Sweet, Hort. Brit. 172. 1826.

Cereus phyllanthoides De Candolle, Prodr. 3: 469. 1828.

Phyllocactus phyllanthoides Link, Handb. Gewächs. 2: 11. 1831.

Opuntia speciosa Steudel, Nom. ed. 2. 2: 222. 1841.

Stems somewhat woody, branching, the branches terete at base, flattened and thin above, sometimes 5 cm. broad, green; margin of branches coarsely crenate; flowers diurnal, the tube 2 cm. long; inner perianth-segments oblong, more or less spreading; filaments and style elongated, slender; stigma-lobes 5 to 7.

Type locality: Mexico.

Distribution: Mexico or Colombia, but known only from cultivated plants.

The distribution of this species is assigned to Mexico, but both Edwards and Sims state definitely that it was first observed by Humboldt and Bonpland near the village of Turbaco, which is a few leagues south of Cartagena, Colombia. From seeds collected at that time, plants were grown in the garden of La Malmaison; one of these flowered in 1811 and was described and illustrated as *Cactus speciosus* in 1813.

This is one of the oldest known species of cacti; it was figured by Hernández in 1651 and by Plukenet in 1691. It has long been in cultivation, perhaps in prehistoric times.

It is often hybridized with other species. The following hybrids with it are given: *Phyllocactus phyllanthoides albiflorus*, *striatus*, *striatus multiflorus*.

Salm-Dyck (Hort. Dyck. 65. 1834) lists four varieties as follows: *Cereus phyllanthoides curtisii*, *C. phyllanthoides guillardieri*, *C. phyllanthoides jenkinsonii*, and *C. phyllanthoides vandesii*. Pfeiffer (Enum. Cact. 124. 1837) also mentions *Cereus phyllanthoides albiflorus*.

Epiphyllum vandesii Don (Gen. Hist. Dichl. Pl. 3: 170. 1834) is a hybrid produced by placing the pollen of *Epiphyllum phyllanthoides* on the stigmas of *Heliocereus elegantissimus*.

Illustrations: Plukenet, Phyt. pl. 247, f. 5, as *Phyllanthos*; Loudon, Encycl. Pl. 413. f. 6902; Curtis's Bot. Mag. 46: pl. 2092, as *Cactus phyllanthoides*; Schumann, Gesamtb. Kakteen 217. f. 42; Monatsschr. Kakteenk. 7: 87; Wiener Ill. Gart. Zeit. 28: f. 39; Gartenwelt 4: 560; 5: 6 and pl. facing 6; Ann. Rep. Smiths. Inst. 1908: f. 24; Möllers Deutsche Gärt. Zeit. 11: 61; Goebel, Pflanz. Schild. 1: f. 13, 52, 54, as *Phyllocactus phyllanthoides*; Bonpland, Descr. Pl. Rar. pl. 3; Edwards's Bot. Reg. 4: pl. 304; Herb. Génér. Amat. 4: pl. 244, as *Cactus speciosus*; Loudon, Encycl. Pl. ed. 2. 1202. f. 17367, as *Epiphyllum speciosum*; Pfeiffer, Abbild. Beschr. Cact. 2: pl. 17, as *Cereus phyllanthoides* var. *stricta*; Ann. Inst. Roy. Hort. Fromont 2: pl. 1, f. E, as *E. phyllanthoides*; Hort. Ripul. pl. 10; Van Géel, Sert. Bot. pl. 111, as *Cactus alatus*.

Figure 207 is a reproduction of Bonpland's illustration as *Cactus speciosus*.



FIG. 207.—*Nopalxochia phyllanthoides*.

9. **WITTIA** Schumann, Monatsschr. Kakteenk. 13: 117. 1903.

Epiphytic, branching cacti, pendent from trees and rocks; joints elongated, flattened or somewhat thickened, spineless, the margins more or less crenate; flowers small for this group, not fugacious, with a definite tube, much longer than limb; perianth-segments short, erect; style (so far as known) slender, white, a little exserted; fruit small, berry-like.

Type species: *Wittia amazonica* Schumann.

In vegetative characters this genus is similar to some of the *Rhipsalidanae*, but the flower has a tube longer than the limb.

The genus is named for N. H. Witt, who made valuable collections in Brazil.

We recognize two species, natives of Panama and northern South America.

KEY TO SPECIES.

Fruit roughened by small tubercles..... 1. *W. amazonica*
Fruit smooth..... 2. *W. panamensis*




FIG. 208.—*Wittia amazonica*.

1. **Wittia amazonica** Schumann, Monatsschr. Kakteenk. 13: 117. 1903.

Branches flattened except at base, 15 to 40 cm. long, 4.5 to 9 cm. broad, often with constrictions, cuneate at base, coarsely crenate, obtuse or acute at apex; flowers 2.5 cm. long; perianth wine-colored, 2 cm. long, cylindric, somewhat curved; perianth-segments 10, erect, in 2 series; stamens included, in 2 series; style 18 mm. long; stigma-lobes 5; ovary strongly tuberculate; scales on ovary 3-angled; fruit 12 to 17 mm. long, deeply umbilicate at apex.

Type locality: Near Laetitia and Tarapoto, Peru.

Distribution: Northeastern Peru, not far from the Brazilian border.

We know the plant from description and illustration only.

Illustration: Monatsschr. Kakteenk. 13: 119.

Figure 208 is reproduced from the illustration cited above.

2. **Wittia panamensis** Britton and Rose, Contr. U. S. Nat. Herb. 10: 241. 1913.

Branches much flattened, up to 1 meter long, 4 to 7 cm. wide, low-crenate; flowers sometimes 15 or more on a branch but solitary at areoles on upper half of joint, purple, becoming straight, 2.5 to 3.5 cm. long, 5-angled, stiff; outer perianth-segments 10, in 2 series, equal, obtuse; outermost ones angled on back; inner perianth-segments, similar to outer but thinner, not angled or only slightly so, a little longer, all erect; innermost segments 10 or 11, thinner, paler, and much smaller than outer, apiculate, sometimes toothed above; tube proper 5 to 6 mm. long, the throat 10 mm. long; stamens many, in 2 series, one on base of throat on long filaments, one on top of throat on short filaments, all included; stigma-lobes, white, but remaining in a close cluster, the top exserted beyond perianth-segments; ovary globular, purple, bearing a few scarious scales.


Type locality: Mountains above Chepo, Panama.

Distribution: Panama, Colombia, and perhaps Venezuela.

Mr. Henri Pittier has collected in Venezuela a plant which is closely related to this species (No. 7656).

Illustrations: Contr. U. S. Nat. Herb. 16: pl. 73; Curtis's Bot. Mag. 145: pl. 8799.

Figure 209, shown below, is from a photograph showing the plant collected by Mr. Pittier in 1912 which afterwards flowered in Washington.



Subtribe 8. RHIPSALIDANAE.

Mostly epiphytic cacti, generally growing on trees but sometimes clambering over rocks or pendent from them, much branched; branches alternate or often in whorls, slender, terete, angled or flat and thin, spineless, except in *Pfeiffera* and *Acanthorhipsis*; flowers regular, mostly small, rotate, and without any tube or with a very short tube; stamens usually few, attached to disk or near base of flower-tube; style usually short; fruit a small juicy berry, white, red, or purple; seeds minute.

We have placed this subtribe at the end of our monograph because it appears to us to represent the most extreme differentiation within the family. It is indeed difficult to explain to most people that its species are really cacti.

We recognize eight closely related genera.

KEY TO GENERA.

- Flowers with a short definite tube.
 Joints terete. 1. *Erythrorhipsis* (p. 208)
 Joints flattened, ribbed, or angled.
 Joints and flowers terminal 2. *Rhispalidopsis* (p. 209)
 Joints and flowers normally lateral.
 Joints with spiny areoles; ovary and fruit with areoles subtended by scales.
 Joints ribbed; fruit-areoles spiny 3. *Pfeiffera* (p. 250)
 Joints flattened or 3-winged; fruit-areoles not spiny 4. *Acanthorhipsis* (p. 211)
 Joints not spiny; fruit mostly without areoles.
 Upper joints normally flattened; areoles not pilose 5. *Pseudorhipsis* (p. 253)
 Upper joints flattened or 3-angled; areoles long-pilose 6. *Lepismium* (p. 215)
 Flowers without tube.
 Petals erect; ends of same joint unlike; flowers and branches always terminal. 7. *Hattiora* (p. 216)
 Petals usually widely spreading; ends of same joint usually similar; flowers and branches lateral or terminal 8. *Rhipsis* (p. 219)

1. ERYTHRORHIPSALIS Berger, Monatsschr. Kakteenk. 30: 4. 1920.

Epiphytic, with slender terete stem and branches, often pendent; branches dichotomous or sometimes in whorls of to 6; areoles scattered, small, all bearing several bristles; flowers terminal, regular, diurnal, white to rose-colored with a short but definite tube; ovary and fruit bristly, the latter red; seeds much larger than in *Rhipsis*.

Type species: *Rhipsis pilocarpa* Löfgren.

The generic name is from *ἐρυθρός* red, and *Rhipsis*, referring to the red fruit and to the resemblance of this genus to *Rhipsis*. Only one species is known.

The genus resembles in habit some of the species of *Rhipsis* with round stems but has a distinct flower-tube, on the top of which the stamens are borne. It also differs from *Rhipsis* in having a long exserted style, exserted even in the bud; in its slowly opening flower (requiring several days to expand); in its very fragrant flower; in having its ovary and fruit bearing areoles, each with a cluster of bristles; and in its larger seeds.

Löfgren, when he described *Rhipsis pilocarpa*, was inclined to think that it might belong to *Pfeiffera*. In his latest treatment of it (Arch. Jard. Bot. Rio de Janeiro 1: 68) he referred it and *Pfeiffera ianthothele* to *Rhipsis* under the subgenus *Pfeiffera*.

At the place cited above, Berger proposed that *Rhipsis pilocarpa* should be regarded as a new subgenus of *Rhipsis* but at the same time he incidentally made it the type of a new genus. Mr. Berger, who has written most interestingly of it, says in part:

"In 1903, Löfgren made known *Rhipsis pilocarpa* (Monatsschr. Kakteenk. 13: 52 to 57) which formerly had a fairly wide distribution in our collections. I received it from various sources, the finest specimens coming from the Botanic Garden in Bremen, from which place it was sent for naming. The plant grew well in my hothouse but appeared to prefer it cooler and sunnier. The habit picture in the Monatsschrift, above cited, is not exactly right. The plant is striking because of its beautiful bristles; it is very odd. In general it does not differ from the rest of the species of *Rhipsis*. Because of its beautiful bristles one is persuaded to put it into *Ophiorhipsis*. Meantime the habit, the flowers, and the fruit show themselves to be a fundamental obstacle.

"In all the species of *Rhipsis* which I have had experience with, the ovary and later the fruit are entirely naked; at the most there is at times a little scale. In these plants, however, the ovaries, which in form remind one of those of *Cereus*, bear a mass of small tubercles with little



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Fruiting branch of *Rhipsalis grandiflora*.
2. Flowering branch of *Rhipsalis lindbergiana*.
3. Fruiting branch of *Rhipsalis shaferi*.
4. Flowering and fruiting branch of *Rhipsalis lindbergiana*.
5. Flowering plant of *Erythrorhipsalis pilocarpa*.
6. Flowering branch of *Rhipsalis grandiflora*.

scales, in whose axils are a large number of projecting white bristles. Still more different is the fruit, which Löfgren did not know. It is striking because of its size, about 10 to 12 mm. by 10 to 12 mm., and while the rest of the *Rhipsalis* fruits in size, form, and color resemble mistletoe berries or are rarely yellow or pale rose, these are strongly wine-red and beset with numerous bristles bearing small areoles, forming a wreath on the umbilicus of the fruit, like *Cereus* and especially *Opuntia*, only the bristles are white and not pricking. In cross-sections the fruit is also red but has a watery sap and a larger number of seeds, coiled on the placenta in the middle of the fruit. The seeds are about double the size of those of *Rhipsalis*."

1. *Erythrorhpsalis pilocarpa* (Löfgren) Berger, Monatsschr. Kakteenk. 30: 4. 1920.

Rhipsalis pilocarpa Löfgren, Monatsschr. Kakteenk. 13: 52. 1903.

Stems dark green to purple, at first erect, sometimes 4 dm. long and unbranched, terminated by 2 to 4 branches in a whorl, the ultimate branches often only 1 cm. long, in time the whole plant becoming pendent; joints clustered, when withering somewhat angled, tipped by yellow bristles; areoles filled with long setose hairs or bristles subtended by ovate scarious bracts; flowers at ends of terminal branches, very fragrant, opening slowly, up to 2 cm. broad; flower-tube 2 mm. long, reddish on the inside; outer perianth-segments 5 or 6, triangular, rose-colored; inner perianth-segments 10 to 15, spreading or sometimes recurved, lanceolate, acuminate, 10 mm. long, white or cream-colored with pinkish tips; stamens numerous, red at bases; ovary with several areoles, bearing as many as 10 bristles, subtended by small scarious scales and surrounded by purple spots; style exserted in the bud; stigma-lobes 4 to 8, white, spreading apart the second day after the appearance of the style and before the stamens appear.

Type locality: Ytu and Ypanema, São Paulo, Brazil.

Distribution: States of São Paulo and Rio de Janeiro, Brazil.

Pfeiffera rhpsaloides Löfgren (Monatsschr. Kakteenk. 13: 54. 1903) was another name suggested for this plant when it was first described.

Illustrations: Blühende Kakteen 2: pl. 99; Monatsschr. Kakteenk. 13: 55; Rev. Centr. Sci. Campinas No. 4, Opp. 188; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 11, 20; Arch. Jard. Bot. Rio de Janeiro 1: pl. 1, as *Rhipsalis pilocarpa*.

Plate XXI, figure 5, is of a plant in the New York Botanical Garden which flowered in April 1919 and was obtained by Dr. Shafer from Dr. Löfgren at Rio de Janeiro in 1917.

2. RHIPSALIDOPSIS gen. nov.

Somewhat shrubby, erect, reclining or pendulous, the joints 3 to 5-angled; branches usually several, terminal; areoles small, sometimes bearing setae; flowers terminal, with a broad rotate limb and a very short tube; stamens erect; style slender; fruit unknown.

Type species: *Rhipsalis rosea* Lagerheim.

One species is known, native of southern Brazil.

This plant was originally described as *Rhipsalis*, but it has a much larger flower and the perianth-segments are united into a short tube. In habit it resembles some of the species of *Epiphyllanthus* but has a regular flower. We have placed it near *Pfeiffera*, but we do not believe that it is close to that genus, for it has a rotate flower and the flowers and branches are terminal, as in *Zygocactus*.

The generic name is given on account of its resemblance to some of the species of *Rhipsalis*.

1. *Rhpsalidopsis rosea* (Lagerheim).

Rhpsalis rosea Lagerheim, Svensk Bot. Tidskr. 6: 717. 1912

Branches short, 1 to 3, strongly 4-angled or sometimes 5-angled, with concave sides; buds red; flowers 3.7 cm. broad, fragrant; perianth-segments few, rose-colored; stamens 11 mm. long, rose-colored; style 13 mm. long, rose-colored; stigma-lobes, white, 3 mm. long.



Fig. 210.—*Rhpsalidopsis rosea*.

Type locality: Woods near Caiguava, state of Parana, Brazil, altitude 1,100 to 1,300 meters.

Distribution: Southern Brazil.

Illustrations: Svensk Bot. Tidskr. 6: pl. 28; Arch. Jard. Bot. Rio de Janeiro 2: pl. 14, 15; Monatsschr. Kakteenk. 32: 121, as *Rhipsalis rosea*.

Figure 210 is reproduced from the first illustration above cited.

3. PFEIFFERA Salm-Dyck, Cact. Hort. Dyck. 1844. 40. 1845.

Epiphytic, with a woody base; branches in wild state hanging, mostly 4-angled, not emitting aerial roots; spines several, acicular; flowers regular, diurnal, pale yellow to rose-colored (sometimes described as purple-red), small, the segments united at base into a very short tube; stamens included, some borne on flower-tube and some on disk; ovary and fruit spiny; seeds black, oblong.

Type species: *Cereus ianthothele* Monville.

Only one species is known, and this was first described as a *Cereus* and afterwards referred to *Rhipsalis*. We agree with the author in regarding it as a distinct genus.

The genus was named for Dr. Ludwig Pfeiffer, a physician by profession and one of the most distinguished authorities on the Cactaceae. He visited Cuba in 1838-1839. Dr. Pfeiffer was born July 4, 1805, at Kassel, Germany, and died in 1877.




FIG. 211.—*Acanthorhipsalis micrantha*.

1. Pfeiffera ianthothele (Monville) Weber, Dict. Hort. Bois 944. 1898.

Cereus ianthothele * Monville, Hort. Univ. 1: 218. 1839.

Pfeiffera cereiformis Salm-Dyck, Cact. Hort. Dyck. 1844. 41. 1845.

Rhipsalis cereiformis Förster, Handb. Cact. 454. 1846.

Hariota cereiformis Kuntze, Rev. Gen. Pl. 1: 262. 1891.

Rhipsalis ianthothele K. Brandegee, Cycl. Amer. Hort. Bailey 4: 1514. 1902.

Stem weak, spreading or pendent, 3 to 6 dm. long, 2 cm. in diameter or less; joints 8 to 12 cm. long, 3 to 5-ribbed, 10 mm. in diameter, light green, spiny; ribs tuberculate; areoles 10 mm. apart; spines 6 or 7, 5 to 7 mm. long, yellowish; flowers including the ovary about 15 mm. long; inner perianth-segments 5, pale yellow to cream-colored, acute, erect or slightly spreading at tip; stamens numerous, shorter than the perianth-segments, included; style longer than stamens; stigma-lobes 8, linear, spreading; ovary strongly tuberculate, purplish, its areoles bearing white bristly spines; fruit globose, 12 to 16 mm. in diameter, rose-red, spiny; seeds numerous, black.

* The specific name is sometimes spelled *ianthothele*; it was originally given as *ianthothelus*.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Pfeiffera ianthothele*.
2. Flowering and fruiting branch of *Lepismium cruciforme*.
3. Top of fruiting branch of *Pfeiffera ianthothele*.
4. Flowering and fruiting branch of *Rhipsalis jamaicensis*.

5. Flowering branch of *Pseudorhipsalis alata*.
6. Flowering and fruiting branch of *Pseudorhipsalis himantoclada*.
7. Flowering branch of *Rhipsalis grandiflora*.

Type locality: Montevideo is cited in the original description, but this must be wrong.

Distribution. Northwestern Argentina, especially in the states of Salta, Tucuman, and Catamarca.

Illustrations: Goebel, Pflanz. Schild. 1: 45, B; Palmer, Cult. Cact. 191; Förster, Handb. Cact. ed. 2. 895. f. 122; Pfeiffer, Abbild. Beschr. Cact. 2: pl. 9; Garten-Zeitung 4: 182. f. 42, No. 10, as *Pfeiffera cereiformis*; Schumann, Gesamtb. Kakteen 611. f. 97, A, B; Blühende Kakteen 3: pl. 152.

Plate XXII, figures 1 and 7, shows flowering branches from a plant collected by Dr. Shafer in Argentina in 1917 (No. 71), which flowered in April 1919; figure 3 shows the mature fruit.

4. ACANTHORHIPSALIS (Schumann) gen. nov.

Small branching cacti, more or less epiphytic, growing on forest trees or creeping over rocks; joints flattened or sometimes 3-winged, short or elongated, their margins crenate or serrate; areoles spiny; flowers solitary from lateral areoles; perianth-segments united into a short tube; ovary bearing on its surface small scales with tufts of felt in their axils, at least in typical species; seeds small, black, narrowed at base.

The type is *Cereus micranthus* Vaupel and to this genus we have also referred two little-known species of *Rhipsalis*, both of which have flattened joints and spiny areoles. In their habit and armament they resemble *Acanthorhipsalis micrantha* more than they do the true species of *Rhipsalis*. The plants are native of Peru, Bolivia, and Argentina.

KEY TO SPECIES.

Joints crenate.

Joints about 2 cm. broad; spines 5 to 15 mm. long.....1. *A. micrantha*

Joints usually 4 to 6 cm. broad; spines 4 mm. long or less.....2. *A. crenata*

Joints serrate.....3. *A. monacantha*

1. *Acanthorhipsalis micrantha* (Vaupel).

Cereus micranthus Vaupel, Bot. Jahrb. Engler 50: Beibl. 111: 19. 1913.

Stems much branched; joints 2 or 3-winged, about 2 dm. long and 2 cm. broad, yellowish green, at least when dry; areoles 6 to 10 mm. apart; spines 3 to 10, 5 to 15 mm. long, brown to blackish, straight or a little curved; flower, including the ovary, 22 mm. long.




FIG. 212.—*Acanthorhipsalis crenata*.

Type locality: Sandia, southeastern Peru, altitude 2,100 meters.

Distribution: Known only from the type locality.

This plant was described by Dr. Vaupel as a species of *Cereus*, but as he writes us under date of October 20, 1920, it is of course not a *Cereus* in the stricter sense, but is more

nearly related to *Rhipsalis*. This view was taken by Schumann, who had labeled it *Rhipsalis peruviana* Schumann (Vaupel, Bot. Jahrb. Engler 50: Beibl. 111: 19. 1913).

The plant was collected by A. Weberbauer July 31, 1902 (No. 1353); a fragment of the type, which is in the Berlin Herbarium, was sent us by Dr. Vaupel in 1920.

Figure 211 is from a photograph of a part of the type specimen now in the National Herbarium at Washington.

2. *Acanthorhipsalis crenata* (Britton).

*Hariota crenata** Britton, Bull. Torr. Club 18: 35. 1891.

Branches lateral, narrowly oblong, very flat, obtuse, 20 to 30 cm. long, 3 to 6 cm. broad, strongly crenate, with a stout central axis; areoles between crenations rather large, filled with wool and bearing 3 to 8 spines, these 2 to 4 mm. long; flowers red, lateral, small; berry 7 mm. in diameter.

Type locality: Yungas, Bolivia.

Distribution: Known only from the type locality.

When first described, this species was thought to be nearest the Brazilian *Rhipsalis platycarpa*, which it resembles, but that species has no spines.

Figure 212 is from a photograph of Dr. Rusby's herbarium specimen (No. 2047).




FIG. 213.—*Acanthorhipsalis monacantha*.

3. *Acanthorhipsalis monacantha* (Grisebach).

Rhipsalis monacantha Grisebach, Abh. Ges. Wiss. Göttingen 24: 140. 1879.

Hariota monacantha Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Epiphytic, branching; branches flat and thin, linear-oblong, 2 cm. broad, sometimes 8 dm. long, obtuse, cuneate at base; serrate (acuminate says Schumann, but figured by him as obtuse); areoles white-felted and spiny, spines 1 to 6, but usually only 1 or 2, 5 to 10 mm. long, yellow; flowers solitary at the areoles, lateral, white, 1 cm. long; fruit globular, 8 to 10 mm. in diameter, white; seeds blackish, pitted, obovoid.

Type locality: Oran, near San Andrés, Argentina.

Distribution: Northern Argentina.

* This name is printed *H. cinerea* in the Index Kewensis.

Illustration: Schumann, Gesamt. Kakteen 633. f. 98, H, as *Rhipsalis monacantha*.

Figure 213 is from a photograph of a herbarium specimen collected at Calilegua, Jujuy, Argentina, by J. A. Shafer in 1917 (No. 56).

5. PSEUDORHIPSALIS gen. nov.

Epiphytic, much branched, and elongated cacti, at first erect, but soon prostrate or hanging; branches flattened, rather thin, serrate or crenate; flowers numerous, borne solitary at the lateral areoles, narrowly campanulate; segments united into a short but definite tube; ovary and fruit globular, bearing several scales; seeds black.

Two species are here included, of which *Cactus alatus* Swartz is made the generic type. These plants in their habit and branches resemble certain species of *Rhipsalis*, especially *R. ramulosa* and its relatives, but differ from all the species of *Rhipsalis* in having united perianth-segments and more scaly ovary and fruit.

KEY TO SPECIES.

Ovary and outer perianth-segments reddish 1. *P. himantoclada*
Ovary and outer perianth-segments greenish or yellowish green. 2. *P. alata*

1. *Pseudorhipsalis himantoclada* (Roland-Gosselin).

Rhipsalis himantoclada Roland-Gosselin, Bull. Soc. Bot. France 55: 694. 1908.

Wittia costaricensis Britton and Rose, Contr. U. S. Nat. Herb. 16: 261. 1913.




FIG. 214.—*Pseudorhipsalis himantoclada*.

Stems 4 to 5 dm. long, erect or curved, flat, 1 to 3 cm. broad, with horizontal branches narrowed at base, pointed, the margin low-serrate; areoles 12 to 15 mm. apart; ovary, tube, and sepals purplish; ovary 3 mm. long, bearing a few very short scales; tube of flower about 8 mm. long; inner perianth-segments white, obtuse, spreading; stamens erect; style white; stigma-lobes 4.

Type locality: Pozo Azul, Costa Rica.

Distribution: Costa Rica.

We are told by Mr. Otón Jiménez that Mr. Wercklé, who first collected the plant, would refer *Wittia costaricensis* here. He states also that it is very luxuriant and when growing wild becomes so large that one man can not carry a single plant.

Illustration: Contr. U. S. Nat. Herb. 16: pl. 82, as *Wittia costaricensis*.

Plate xxii, figure 6, shows a flowering branch collected by Wercklé in 1907 which flowered in the New York Botanical Garden, December 20, 1911. Figure 214 is from a photograph of a terminal branch; figure 215 shows a flowering branch; figure 216 shows a flower cut longitudinally.

2. *Pseudorhipsalis alata* (Swartz).

Cactus alatus Swartz, Prodr. 77. 1788.

Cereus alatus De Candolle, Prodr. 3: 470. 1828.

Epiphyllum alatum Haworth, Phil. Mag. 6: 109. 1829. Not Haworth, 1819.

Rhipsalis swartziana Pfeiffer, Enum. Cact. 131. 1837.

Hariota swartziana Lemaire, Cact. Gen. Nov. Sp. 75. 1830.

Rhipsalis alata Schumann in Martius, Fl. Bras. 4: 288. 1890.

Hariota alata Kuntze, Rev. Gen. Pl. 1: 262. 1891.


Rhipsalis harrisii Gürke, Monatsschr. Kakteenk. 18: 180. 1809.

Pendent from trees and rocks, up to 5 meters long, branched; joints broadly linear to lanceolate or linear-oblong, 2 to 4 dm. long, 3 to 6 cm. broad, obtuse, the margin crenate-undulate; flowers yellowish white, 15 mm. long; flower-tube 4 mm. long; perianth-segments 10, lanceolate, acute; stamens numerous, about half as long as perianth; style slender; stigma-lobes; ovary somewhat tubercled, bearing several broad scales; fruit ovoid, 1 cm. long, yellowish green; seeds obovate, black, bearing depressed tubercles; hilum oblique.

Type locality: Jamaica.

Distribution: Mountains of Jamaica.

This plant has usually passed as a *Rhipsalis*, but its definite flower-tube and somewhat tubercled and scaly ovary exclude it from that genus. This species has long been known in Jamaica; it was mentioned by Sloane as a spineless *Opuntia* and it is also referred to by Patrick Browne.



FIGS. 215 and 216.—Top of flowering branch and longitudinal section of flower of *Pseudorhipsalis himantoclada*. $\times 2$.

FIGS. 217 and 218.—Section of flower and flowering branch of *Pseudorhipsalis alata*. $\times 0.66$.

Cactus dentatus Ruiz (Martius, Fl. Bras. 4²: 288. 1890) was given as a synonym of *Rhipsalis alata* by Schumann, but better referred to *R. ramulosa* (see page 241).

Cereus alatus crassior Salm-Dyck (Hort. Dyck. 66. 1834) is only a name, which may or may not refer to the Jamaican plant.

Illustration: Torrey 9: 157. f. 2, as *Rhipsalis alata*.

Plate XXII, figure 5, shows a plant collected by Dr. Britton in Jamaica in 1907, which flowered in the New York Botanical Garden, November 8, 1912. Figure 218 shows a flowering branch (natural size); figure 217 shows half of a flower with tube, perianth-segments, and stamens.

6. LEPISMIUM Pfeiffer, Allg. Gartenz. 3: 315, 380. 1835.

Saxicolous or epiphytic cacti, usually much branched and elongated, the branches flat, angled or 3-winged, the margins strongly crenate; areoles in the crenations producing a tuft of long white hairs; flowers 1 to 5 at an areole, white to pinkish; perianth-segments united at base into a short tube; filaments slender, adnate to flower-tube; stigma-lobes 4 or 5; fruit globose, smooth, turgid, purple; seeds minute; cotyledons broad, acuminate.

Type species: *Lepismium commune* Pfeiffer.

We recognize but one species, which has been described under many names. The generic name is from *λεπίς*, a scale, referring to the small scales subtending the areole.

1. Lepismium cruciforme (Vellozo) Miquel, Bull. Néerl. 49. 1838.

- Cactus cruciformis* Vellozo, Fl. Plum. 207. 1825.
Cereus tenuispinus Haworth, Phil. Mag. 1: 125. 1827.
Cereus myosurus Salm-Dyck in De Candolle, Prodr. 3: 469. 1828.
Cereus tennis De Candolle, Prodr. 3: 469. 1828.
Cereus squamulosus Salm-Dyck in De Candolle, Prodr. 3: 469. 1828.
Cereus setosus Loddiges, Bot. Cab. 19: pl. 1887. 1832.
Lepismium tenue Pfeiffer, Allg. Gartenz. 3: 315. 1835.
Lepismium commune Pfeiffer, Allg. Gartenz. 3: 315. 1835.
Lepismium knightii Pfeiffer, Allg. Gartenz. 3: 380. 1835.
Lepismium myosurus Pfeiffer, Enum. Cact. 139. 1837.
Cereus cruciformis Steudel, Nom. ed. 2. 1: 333. 1840.
Rhipsalis myosurus Förster, Handb. Cact. 455. 1846.
Rhipsalis mittleri Förster, Handb. Cact. 455. 1846.
Rhipsalis knightii Förster, Handb. Cact. 456. 1846.
Lepismium myosurus knightii Salm-Dyck in Labouret, Monogr. Cact. 445. 1853.
Lepismium myosurus laevigatum Salm-Dyck in Labouret, Monogr. Cact. 446. 1853.
Lepismium radicans Vöchtling, Jahrb. Wiss. Bot. Leipzig 9: 399. 1873.
Lepismium cavernosum Lindberg, Gartenflora 39: 251. 1890.
Rhipsalis brevibarbis Schumann in Martius, Fl. Bras. 4: 268. 1890.
Rhipsalis squamulosa Schumann in Martius, Fl. Bras. 4: 280. 1890.
Rhipsalis macropogon Schumann in Martius, Fl. Bras. 4: 282. 1890.
Hariota cruciformis Kuntze, Rev. Gen. Pl. 1: 262. 1891.
Hariota squamulosa Kuntze, Rev. Gen. Pl. 1: 263. 1891.
Hariota knightii Kuntze, Rev. Gen. Pl. 1: 263. 1891.
Hariota knightii tenuispinis Kuntze, Rev. Gen. Pl. 1: 263. 1891.
Rhipsalis anceps Weber, Rev. Hort. 64: 427. 1892.
Rhipsalis cavernosa Schumann, Monatsschr. Kakteenk. 3: 24. 1893.
Rhipsalis radicans Weber, Dict. Hort. Bois 1047. 1898.
Rhipsalis radicans anceps Weber, Dict. Hort. Bois 1047. 1898.
Rhipsalis radicans ensiformis Weber, Dict. Hort. Bois 1047. 1898.
Lepismium cavernosum ensiforme Weber in Roland-Gosselin, Rev. Hort. 70: 108. 1899.

Usually creeping over rocks, freely rooting, appressed, somewhat branching; branches foliaceous, usually flat, sometimes 3, 4, or even 5-angled, linear-lanceolate, 2 cm. broad, narrowed at base, more or less purplish, especially on edges; margins somewhat repand; areoles sunken in margins; flowers white, 2 to , or even more from an areole, 12 to 13 mm. long; fruit globular, juicy, purplish to red, translucent, 6 to 12 mm. in diameter; seeds light brown to black, 1.8 mm. long.

Type locality: Coast of Brazil.

Distribution: States of Rio de Janeiro and Minas Geraes, Brazil.

Rhipsalis radicans rosea Weber (Dict. Hort. Bois 1047. 1898) has small rose-colored flowers which, according to Weber, resemble those of *R. myosurus*.

Schumann (Gesamtb. Kakteen 649. 1898) gives *Lepismium anceps* Weber (in Hort. Paris) as a synonym for *Rhipsalis anceps*. Here belongs also *R. ensiformis* Weber (Dict. Hort. Bois 1047. 1898).

Some of the plants now in cultivation are not so broadly winged as is shown in the illustration in Curtis's Botanical Magazine referred to below. This illustration was based upon specimens which were supposed to have come from Prince de Salm-Dyck and, therefore, presumably are typical.

Cereus knightii Parmentier (Pfeiffer, Enum. Cact. 139. 1837) is given as a synonym of *Lepismium knightii*.

Cactus tenuis Schott (De Candolle, Prodr. 3: 469. 1828) was cited as a synonym of *Cereus tenuis*.

Schumann cited *Lepismium mittleri* as a synonym of *Rhipsalis squamulosa*, referring it to Förster (Handb. Cact. 455. 1846), but the plant is there described as *Rhipsalis mittleri*.

Cereus elegans Hortus appeared first (Pfeiffer, Enum. Cact. 138. 1837) as a synonym of *Lepismium commune*, while the Index Kewensis refers it to *Rhipsalis mittleri*.

Cereus myosurus tenuior Salm-Dyck (Hort. Dyck. 65. 1834) is only a name.

Lepismium cavernosum minus Lindberg is a name mentioned by Roland-Gosselin (Rev. Hort. 70: 108. 1899).

Lepismium duprei, the name mentioned by Salm-Dyck (Cact. Hort. Dyck. 1844. 41. 1845) and by Förster (Handb. Cact. 456. 1846) as in the collections at Paris, was never described.

Lepismium laevigatum Salm-Dyck (Cact. Hort. Dyck. 1844. 41. 1845) is without description, nor do we find it listed in the Index Kewensis.

Illustrations: Fl. Flum. 5: pl. 29, as *Cactus cruciformis*; Loddiges, Bot. Cab. 1: pl. 1887; Loudon, Encycl. Pl. ed. 2. 1202. f. 17365, as *Cereus setosus*; Palmer, Cult. Cact. 195, as *Lepismium*; Curtis's Bot. Mag. 66: pl. 3755; Garten-Zeitung 4: 182. f. 42, No. 3; Loudon's Encycl. Pl. ed. 3. 1380. f. 19411, as *Lepismium myosurum*; Monatsschr. Kakteenk. 3: 41, as *L. knightii*; Abh. Bayer. Akad. Wiss. München 2: pl. 7, f. 1; Curtis's Bot. Mag. 66: pl. 3763; Förster, Handb. Cact. ed. 2. 898. f. 123 (in error 103); Loudon, Encycl. Pl. ed. 3. 1380. f. 19412; Nov. Act. Nat. Cur. 19¹: pl. 16, f. 12, as *L. commune*; Goebel, Pflanz. Schild. 1: pl. 2, f. 3, 4, as *L. radicans* (seedling); Gartenwelt 16: 633; Schumann, Gesamtb. Kakteen f. 98, C, D, as *Rhipsalis cavernosa*; Gartenflora 39: f. 38, as *Lepismium cavernosum*; Martius, Fl. Bras. 4²: pl. 55, f. 2, as *Rhipsalis macropogon*; Arch. Jard. Bot. Rio de Janeiro 1: pl. 25, as *Rhipsalis radicans*; Arch. Jard. Bot. Rio de Janeiro 1: pl. 24, as *R. myosura*; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 19, as *R. squamulosa*; Rev. Hort. 85: f. 152, as *R. anceps*.

Plate XXII, figure 2, shows the plant obtained by Dr. Rose in Brazil in 1915 which flowered November 18 of that year.

LEPISMIUM RAMOSISSIMUM Lemaire in Förster, Handb. Cact. ed. 2. 899. 1885.

Rhipsalis ramosissima Schumann in Martius, Fl. Bras. 4²: 299. 1890.

Hariota ramosissima Kuntze, Rev. Gen. Pl. 1: 263. 1891.

This is a very uncertain species which we know only from descriptions. It is from Brazil.

7. HATIORA Britton and Rose, Stand. Cycl. Hort. Bailey 3: 1432. 1915.

Hariota De Candolle, Mém. Cact. 23. 1834. Not Adanson, 1763.

Unarmed, slender, branched cacti; branches terete, short, arising in 2's or 3's from tops of older ones, smooth, leafless and spineless,* bearing several small areoles along their sides and each a large, woolly, terminal one from which the flower and succeeding branches arise; sepals usually in 2 series, outer ones broader and short, inner ones larger and more petal-like; petals distinct, narrowed toward base; stamens distinct, erect, borne on disk; stigma-lobes 4 or 5, erect or a little spreading, white; ovary globular, naked or nearly so.

Type species: *Rhipsalis salicornioides* Haworth.

Some six or seven species have been described; we recognize three.

The genus *Hariota* was named for Thomas Hariot, a botanist of the 16th century, *Hatiora* being an anagram. It is closely related to *Rhipsalis*, with which it is often united.

The flowers open only in bright sunlight and are rotate or nearly so. In the United States the plants flower under glass, usually in the winter from December to February, but sometimes as late as April.

*Sometimes peculiar lateral branches are produced which are made up of short, rounded joints with numerous areoles bearing several bristles or hairy spines. See illustrations of Schumann (Gesamtb. Kakteen f. 97, I) and Loddiges (Bot. Cab. 4: 369). In cases which we have observed these occur on stunted or starved plants, the areoles arranged in 6 rows forming low angles on the branchlets.

KEY TO SPECIES.

- Lower part of joints slender, pedicel-like 1. *H. salicornioides*
 Joints only slightly narrowed below or not narrowed.
 Joints clavate 2. *H. bambusoides*
 Joints cylindric 3. *H. cylindrica*

1. *Hatiora salicornioides* (Haworth) Britton and Rose, Stand. Cycl. Hort. Bailey 3: 1433. 1915.

Rhipsalis salicornioides Haworth, Suppl. Pl. Succ. 83. 1819.

*Cactus salicornioides** Link and Otto, Icon. Pl. Select. 49. 1822.

Cactus lyratus Vellozo, Fl. Flum. ed. 2. 4: 205. 1825.

Hariota salicornioides De Candolle, Mém. Cact. 23. 1834.

Rhipsalis salicornioides strictior Salm-Dyck, Cact. Hort. Dyck. 1849. 230. 1850.

Hariota salicornioides strictior Gürke, Blühende Kakteen 2: under pl. 95. 1907.

Stems to 2 meters long with a jointed cylindric trunk; branchlets club-shaped, the lower part very slender and pedicel-like, 1.5 to 3 cm. long, green or purplish; areoles of cultivated specimens without setae; flowers 8 to 10 mm. long, salmon-colored, the outer sepals short and obtuse; inner petals somewhat crenate, obtuse; filaments yellowish, at top appressed against style, shorter than petals; style yellowish; stigma-lobes 4 or 5, white.

Type locality: Recorded originally from the West Indies in error.

Distribution: Southeast Brazil.

These plants grow quite differently in the woods from the way they do in greenhouses. The following note was made by Dr. Rose in 1915 while collecting at Rio de Janeiro:

The plant grows on trunks of trees, its roots long and fibrous, dm. long or more and wrapped about the trunk of the tree; at first it is erect, then spreading and finally pendent; it is then a meter long or more and very much branched; main stem and branches 5 to 10 mm. in diameter, made up of short terete joints (2 to 5 cm. long); branches in whorls of 2 to 6.

A very remarkable form, if not a distinct species, was obtained by Dr. Rose in the forest of Itatiaia, altitude 1,200 meters, in July 1915 (No. 20585). The terminal joints are 1 to 2 cm. long, the lower half slender, pedicel-like, the upper half twisted and contorted. This is well shown in our illustration (figure 219).

Rhipsalis salicornioides gracilior (Salm-Dyck, Cact. Hort. Dyck. 1844. 40. 1845; *Hariota salicornioides gracilior* Gürke, Blühende Kakteen. 2: under pl. 9. 1907) is only a name.

The following varieties of *Rhipsalis salicornioides* of Weber are probably to be referred here: var. *gracilis* Weber (Dict. Hort. Bois 1048. 1898; *Rhipsalis gracilis* Weber and *Hariota gracilis* Weber, Dict. Hort. Bois 1048. 1898) and var. *stricta* Weber (Dict. Hort. Bois 1048. 1898; *Rhipsalis stricta* Cels, Dict. Hort. Bois 1048. 1898). The name *Rhipsalis stricta* seems never to have been published. Weber cited it as above, referring it to Cels as the author. Schumann uses the name earlier where he states that it was used in France for *Hariota salicornioides* (Monatsschr. Kakteenk. 4: 74. 1894). Pfeiffer refers here as a synonym *Opuntia salicornioides* (Enum. Cact. 141. 1837), attributing the name to Sprengel, who, however, used it as *Cactus* (*Opuntia*) *salicornioides*. *Hariota stricta* has been used (Monatsschr. Kakteenk. 5: 22. 1895). The variety *ramosior* Salm-Dyck (Pfeiffer, Enum. Cact. 142. 1837) may or may not belong to this species.




FIG. 219.—Unusual form of *Hatiora salicornioides*. $\times 0.8$.

* This name is often credited to Sprengel (Syst. 2: 497. 1825).

Rhipsalis schottmuelleri Hortus is given by Schelle (Handb. Kakteenk. 227. 1907) as a synonym of *Hariota salicornioides schottmuelleri*, an unpublished variety.

Hariota villigera (Schumann in Martius, Fl. Bras. 4²: 265. 1890; *Rhipsalis salicornioides villigera* Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 85. 1915) we know from description only; it seems to be stouter than *salicornioides* but may belong here. It was based on Sellow's specimen from São Paulo, but its flowers are unknown.

Illustrations: Loddiges, Bot. Cab. 4: pl. 369; Cact. Journ. 1: 180; Curtis's Bot. Mag. 51: pl. 2461; Blanc, Cacti 90. No. 1013; Balt. Cact. Journ. 1: 122; Gard. Chron. II. 6: 731. f. 134; Amer. Gard. 11: 463; Goebel, Pflanz. Schild. 1: pl. 4, f. 5, 6; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 17; Arch. Jard. Bot. Rio de Janeiro 1: pl. 12; Gartenwelt 13: 117, as *Rhipsalis salicornioides*; Link and Otto, Icon. Pl. Select. pl. 21, as *Cactus salicornioides*; Schumann, Gesamtb. Kakteen f. 97, C, D; Martius, Fl. Bras. 4²: pl. 52; Monatsschr. Kakteenk. 5: 23; Schelle, Handb. Kakteenk. 227. f. 148, as *Hariota salicornioides*; Rev. Hort. 1861: 110. f. 23, as *Rhipsalis salicorne*; Fl. Flum. 5: pl. 21, as *Cactus lyratus*.

Plate XXIII, figure 4, shows a plant in the New York Botanical Garden which flowered February 2, 1912. Figure 219 shows a peculiar form collected by Dr. Rose in Brazil in 1915.




FIG. 220.—*Hattiora bambusoides*.

2. *Hattiora bambusoides* (Weber).

Rhipsalis salicornioides bambusoides Weber, Rev. Hort. 64: 429. 1892.

Hariota salicornioides bambusoides Schumann, Gesamtb. Kakteen 613. 1898.

Rhipsalis bambusoides Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 41. 1918.

Stems becoming 2 meters high and stouter than those of *H. salicornioides*; joints clavate, 3 to 5 cm. long, 4 mm. in diameter at the top; flowers orange; sepals obtuse; petals usually erect but sometimes spreading.

Type locality: Brazil.

Distribution: State of Rio de Janeiro, Brazil.



M. E. Eaton del.

A. Hoern & Co. Baltimore

1. Flowering branch of *Hatiora cylindrica*.
2. Fruiting branch of *Rhipsalis heteroclada*.

3. Fruiting branch of *Rhipsalis cribrata*.
4. Flowering branch of *Hatiora salicornioides*.

Introduced into Jardin des Plantes, Paris, France, from Brazil.

We have not seen this type material, but if plate 95 in the *Blühende Kakteen* is typical, our identification is correct. In the description accompanying this plate it is stated that the drawing was made from a plant sent by Mr. Weber to the Berlin Botanical Garden.

Hariota bambusoides Weber (Dict. Hort. Bois 1048. 1898) was given as a synonym but was never described.

Illustrations: *Blühende Kakteen* 2: pl. 95, as *Hariota salicornioides bambusoides*; *Gartenwelt* 13: 117, as *Rhipsalis salicornioides bambusoides*.

Figure 220 is reproduced from the first illustration cited above.

3. *Hatiora cylindrica* sp. nov.

Forming dense masses one meter in diameter or more; joints cylindric, 3 cm. long or less, pale green, becoming spotted or finally red throughout; flowers usually solitary, 12 mm. long; sepals ovate, short, red; petals orange to yellow, oblong, obtuse.

Collected by J. N. Rose in company with Dr. Löfgren and Señor Porto at Ilha Grande, Distrito Federal, near Rio de Janeiro, July 22 to 24, 1915.

Illustration: Arch. Jard. Bot. Rio de Janeiro 2: pl. 13, ad *Rhipsalis bambusoides*.

Plate xxiii, figure 1, shows the plant collected by Dr. Rose on Ilha Grande, near Rio de Janeiro, which flowered in the New York Botanical Garden, December 18, 1918.

8. RHIPSALIS Gaertner, Fruct. Sem. 1 137. 1788.

**Hariota* Adanson, Fam. Pl. 2: 243. 1763.

†*Cassytha* Miller, Gard. Dict. ed. 8. 1768. Not Linnaeus, 1753.

Cacti sometimes growing in humus, but usually epiphytic and hanging from trees, sometimes erect, sometimes clambering over rocks, more or less rooting or, when hanging, irregularly producing aerial roots; roots always fibrous; stems usually much branched (often heteromorphic), terete, angled or much flattened and leaf-like, very slender and thread-like or stout and stiff; leaves wanting or represented by minute bracts; areoles borne along margin of flat-branched forms, along ribs or scattered irregularly in other forms, usually small, bearing hairs, wool, bristles, and flowers; flowers usually solitary, but sometimes several from a single areole, opening night or day and remaining open for 1 to 8 days, small for the family; perianth-segments distinct, few, sometimes only 5, usually spreading, sometimes reflexed; filaments few or numerous, erect, slender, borne on outer margin of disk in one or two rows; style erect; stigma-lobes 3 or more, usually slender, spreading; ovary small, sometimes depressed or sunken in branch; fruit globular or oblong, sometimes angled when immature, but finally turgid, juicy, white or colored, usually naked (setose at areoles in 1 or 2 species) or sometimes bearing a few scales; seeds small, few to many.

Type species: *Rhipsalis cassutha* Gaertner.

The generic name is from *πίψ* wicker-work, referring to the slender, pliable branches of the typical species.

We recognize 57 species, although more than 115 names have been published.

The species range from Florida, Mexico, and the West Indies through continental America to Argentina; only 2 species are found in Mexico; 1 in Florida; 2 are known in the West Indies; a very few in northern South America; 3 or 4 only on the west coast of South America; and , or 6 in Argentina. The center of distribution is in the states of Rio de Janeiro, São Paulo, and Minas Geraes, in southern Brazil. In the little state of Rio de Janeiro and chiefly about the city of the same name, Dr. Rose collected 15 species in 1915.

The occurrence of species of *Rhipsalis*, in the wild state, in tropical Africa and in Ceylon, forms the only possible exception to the American natural distribution of cacti. Eight sup-

* No species was cited by Adanson for his genus *Hariota* but it was based on Burmann's plate of Plumier (pl. 197, f. 2), which has been identified as *Cactus parasiticus* Lamarck, not Linnaeus. The type of *Cactus parasiticus* Linnaeus is a species of *Vanilla*, probably *V. claviculata* Swartz.

† Miller, in his *Gardeners' Dictionary* of 1768, described *Rhipsalis cassutha* under the name of *Cassytha filiformis*, a name which had already been published by Linnaeus for a wholly different plant. Miller's generic name, *Cassytha*, therefore, being a misidentification, should not be treated as a synonym proper of *Rhipsalis*, although usually so cited.

posedly distinct species have been described by authors from tropical Africa, and *R. cassutha* has long been known to exist in Ceylon. M. Roland-Gosselin, a diligent French student of cacti, after an investigation of these Old World plants, published in 1912 a very interesting paper,* giving his conclusion that they are really all American species, their seeds having been transmitted to the Old World by migratory birds, and he referred them all to known American species. We have followed him in these reductions but we have not been able in all cases to study authentic specimens. It raises the interesting question if the Old World plants should be regarded as native or introduced.

In stem structure some of the species, such as *Rhipsalis elliptica*, approach very closely *Zygocactus truncatus*, while certain forms of *Epiphyllanthus* are easily mistaken for a *Rhipsalis*.

As we have treated the genus here, the flowers and fruits are fairly uniform. The stem structures are various and parallel in a way those of *Opuntia*, ranging from slender and terete to broad and thin; in some species they are leaf-like as in *Epiphyllum*, or 3-angled, suggesting *Hylocereus*. The areoles are usually small and bear only a small tuft of wool, but in some species they bear hairs or bristles. The flowers may open at any time of the day and in most species do not close at the approach of night; they are not readily affected by shade or direct sunlight and open but once.

KEY TO SPECIES.

- A. Joints terete, ribbed or angled, none of them flat.
 - B. Joints terete or young ones angled, smooth, or areoles bristly or hairy.
 - C. Joints short, oblong, not more than 5 times as long as thick; areoles of young joints with a few long hairs.
 - Lateral joints simple; flowers lateral. Series 1, *Mesembryanthemoides* 1. *R. mesembryanthemoides*
 - Lateral joints much branched; flowers terminal. Series 2, *Cereusculae* 2. *R. cereuscula*
 - CC. Joints cylindric, rarely clavate, slender, short or elongated.
 - D. Flowering areoles small, not very woolly, not depressed.
 - E. Ultimate joints slender, about 2.5 mm. thick or less, relatively short.
 - F. Young joints or some of them angled, their areoles bearing hairs. Series 3, *Prismaticae*.
 - Species of Brazil or Madagascar 3. *R. prismatica*
 - Species of Costa Rica 4. *R. simmleri*
 - FF. All joints cylindric to clavate. Series 4, *Capilliformes*.
 - Joints clavate 5. *R. clavata*
 - Joints cylindric.
 - Ultimate joints up to 2.5 mm. thick; petals 9 mm. long 6. *R. campos-portoana*
 - Ultimate joints about 1.5 mm. thick; petals 6 mm. long or less.
 - Plant stiff; areoles red; flowers white 7. *R. heteroclada*
 - Plant weak; areoles not red.
 - Flowers greenish white or yellowish, to 6 mm. wide. 8. *R. capilliformis*
 - Flowers white or nearly white, about 8 mm. wide.
 - Pendent; secondary branches 2 to 3-chotomous 9. *R. burchellii*
 - Spreading or diffuse; upper branches subverticillate 10. *R. cribrata*
 - EE. Ultimate joints stouter, mostly 3 to 10 mm. thick and elongated.
 - F. Scale subtending the areoles inconspicuous or none.
 - G. Ultimate joints 3 to 6 mm. thick. Series 5, *Cassuthae*.
 - H. Ultimate joints definitely shorter than others, often verticillate.
 - Plants weak, pendent. 11. *R. cassutha*
 - Plants stiffer, not strictly pendent.
 - Ultimate joints slender 12. *R. virgata*
 - Ultimate joints stout 13. *R. teres*
 - HH. Ultimate joints not definitely shorter than others, simple or dichotomous, rarely verticillate.
 - I. Areoles without bristles or with spreading bristles.
 - Petals about 4 mm. long.
 - Fruit naked
 - Petals pink 14. *R. lindbergiana*
 - Petals white. 15. *R. shaferi*
 - Fruit with scales and these setose in axils 16. *R. fasciculata*
 - Petals 8 to 12 mm. long.
 - Flowers purplish red; areoles not bristly. 17. *R. pulchra*
 - Flowers white; areoles somewhat bristly 18. *R. lumbricoides*

* Bull. Soc. Bot. France 59: 97-102. 1912. Translation in Torrey 13: 151-156. 1913.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Rhipsalis leucorhaphis*.
2. Fruiting branch of same.
3. Fruiting branch of *Rhipsalis megalantha*.
4. Fruiting branch of *Rhipsalis neves-armondii*.
5. Fruiting branch of same.

6. Fruiting branch of *Rhipsalis pittieri*.
7. Flowering and fruiting branch of *Rhipsalis shaferi*.
8. Flowering branch of *Rhipsalis aculeata*.
9. Flowering of *Rhipsalis mesembryanthemoides*.
10. Flowering of same.

KEY TO SPECIES—continued.

- II. All areoles with appressed bristles 19. *R. aculeata*
- GG. Ultimate joints 8 to 12 mm. thick. Series 6, *Grandiflorae*.
 Flowers 2 to 2.5 cm. broad 20. *R. grandiflora*
 Flowers up to 4 cm. broad 21. *R. megalantha*
- FF. Scale subtending the areole white, scarious, appressed, conspicuous. Series 7, *Leucorhaphes*.
 Joints to 8 mm. thick; areoles with deciduous bristles 22. *R. leucorhaphis*
 Joints 3 mm. thick; areoles without bristles 23. *R. loefgrenii*
- DD. Flowering areoles large, very woolly, depressed. Series 8, *Floccosae*.
 Ultimate joints much shorter than others; verticillate; plants stiff. 24. *R. neves-armondii*
 Ultimate joints not much shorter than others; plants weak.
 Main branches stout, 8 to 10 mm. in diameter.
 Fruit pure white; Venezuelan species. 25. *R. pittieri*
 Fruit red or tinged with red; Brazilian and Argentine species.
 Fruit bright red 26. *R. pulvinigera*
 Fruit whitish but when mature tinged with red or purple.
 Fruit 5 mm. in diameter; flowers white, becoming yellowish 27. *R. floccosa*
 Fruit 8 to 10 mm. in diameter; flowers tinged with red, larger than the last 28. *R. tucumanensis*
 All branches slender, 3 to 6 mm. in diameter.
 Flowers white 29. *R. gibberula*
 Petals with red tips. 30. *R. puniceo-discus*
- BB. Joints ribbed or angled, at least when old.
 C. Some joints bristly, others unarmed, ribbed when old. Series 9, *Dissimiles*. 31. *R. dissimilis*
- CC. All joints unarmed, angled or winged.
 Joints 5-angled or 5-winged.
 Joints 5-winged; wings crenate. Series 10, *Pentapterae*. 32. *R. pentaptera*
 Joints 5-angled; ribs nearly continuous. Series 11, *Sulcatae*. 33. *R. sulcata*
 Joints 3-angled.
 Angles of joints continuous, wingless. Series 12, *Trigonae*. 34. *R. trigona*
 Angles of joints interrupted, winged. Series 13, *Paradoxae*. 35. *R. paradoxa*
- AA. At least some of joints flat, on same plant or on different plants.
 B. Joints deeply serrate; flowers nodding. Series 14, *Houlletianae*. 36. *R. houlletiana*
- BB. Joints repand, entire or crenate; flowers not nodding.
 C. Joints linear to linear-lanceolate. Series 15, *Lorentzianae*.
 D. Joints both flat and 3-angled or rarely 4-angled, mostly narrowly linear.
 Fruit black.
 Joints 10 mm. wide or less; fruit to 6 mm. in diameter 37. *R. warmingiana*
 Joints 10 to 15 mm. wide; fruit about 10 mm. in diameter 38. *R. gonocarpa*
 Fruit white to reddish.
 Joints long-acuminate 39. *R. linearis*
 Joints blunt.
 Joints scarcely crenate; Ecuadorean and Peruvian species 40. *R. micrantha*
 Joints definitely crenate; Costa Rican species 41. *R. tonduzii*
- DD. All joints flat, linear-lanceolate to oblong.
 Joints deeply crenate, the lobes rounded 42. *R. boliviana*
 Joints repand, low-crenate, or nearly entire.
 Joints coriaceous, distinctly crenate 43. *R. lorentziana*
 Joints thin in texture, merely repand or low-crenate.
 Bolivian species 44. *R. ramulosa*
 West Indian and Central American species.
 Larger joints 3 cm. wide 45. *R. purpusis*
 Joints 0.5 to 2.5 cm. wide.
 Perianth-segments 7 to 8 mm. long, greenish white or pinkish 46. *R. coriacea*
 Perianth-segments to 6 mm. long, green 47. *R. jamaicensis*
- CC. Joints elliptic to oblong. Series 16, *Crispatae*.
 Joints thick, coriaceous.
 Joints oblong, cuneate at base.
 Flowers and fruit usually solitary at areoles 48. *R. platycarpa*
 Flowers several at an areole 49. *R. russellii*
 Terminal joints short-oblong to elliptic.
 Fruit red. 50. *R. elliptica*
 Fruit white. 51. *R. pachyptera*
- Joints thin.
 Brazilian species.
 Joints purplish green, obovate. 52. *R. rhombea*
 Joints bright green or reddish.
 Joints reddish green, the margins much crisped. 53. *R. crispimarginata*
 Joints bright green, the margins slightly crisped.
 Ultimate joints broad, elliptic to obovate. 54. *R. crispata*
 Ultimate joints narrow, oblong to narrowly obovate. 55. *R. oblonga*
 Bolivian species 56. *R. cuneata*
- AAA. Species not grouped 57. *R. roseana*

1. *Rhipsalis mesembryanthemoides** Haworth, Rev. Pl. Succ. 71. 1821.*Rhipsalis salicornioides* * (variety B) Haworth, Suppl. Pl. Succ. 83. 1819.*Hariota mesembrianthemoides* † Lemaire, Cact. Gen. Nov. Sp. 74. 1839.

Branches very dissimilar; main branches elongated, slender, terete, more or less setose, often bearing aerial roots, covered with short stubby branchlets: these sometimes also bearing short joints, usually less than 1 cm. long, more or less angled, often with short setae from the small areoles; flower-buds small, pinkish; flowers solitary at areoles of the branchlets, opening in early morning, rather large, 1.5 cm. broad, white or light pink; petals 5, spreading, acute; stamens about 20, erect, white; style white; stigma-lobes 3, white; fruit short-oblong, 5 mm. long, white or tinged with red.

Type locality: Not cited where published.

Distribution: Rio de Janeiro, Brazil.

The plant is common in cultivation; in nature it grows in dense masses on trunks of trees. It first flowered in cultivation in England in 1831. Its short joints have a fancied resemblance to species of 11 *Mesembryanthemum*.

A dried specimen of Haworth's plant is still preserved in London and through the kindness of N. E. Brown we have a photograph of it.

Rhipsalis echinata was published as a synonym by Pfeiffer (Enum. Cact. 136. 1837).

Illustrations: Cact. Journ. 1: 180; Curtis's Bot. Mag. 58: pl. 3078; Schumann, Gesamtb. Kakteen 633. f. 98, G; Monatsschr. Kakteenk. 2: 9; 4: 59; Arch. Jard. Bot. Rio de Janeiro 1: pl. 11; Goebel, Pflanz. Schild. 1: pl. 4, f. 7; Loddiges, Bot. Cab. 20: pl. 1920; Thomas, Zimmerkultur Kakteen 58.

Plate XXIV, figure 9, shows a fruiting plant obtained by Dr. Rose in Rio de Janeiro in 1915 (No. 20246); figure 10 shows a flowering plant sent by Alwin Berger in 1908.

2. *Rhipsalis cereuscula* Haworth, Phil. Mag. 7: 112. 1830.*Hariota saglionis* Lemaire, Cact. Aliq. 39. 1838.*Rhipsalis saglionis* Otto in Walpers, Repert. Bot. 2: 936. 1843.*Rhipsalis brachiata* Hooker in Curtis's Bot. Mag. 69: pl. 4039. 1843.*Hariota cereuscula* Kuntze, Rev. Gen. Pl. 1: 262. 1891.*Rhipsalis saglionis rubrodiscus* Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 80. 1915.

Stems and branches terete; stem slender, usually elongate, often erect, sometimes 6 dm. high, crowned by a cluster of short branches; upper branches short, 2 to 6 times as long as thick, somewhat angled, the areoles bearing 2 to 4 short bristles; flowers terminal or near the ends of the branches, 16 mm. broad; petals about 12, spreading, pinkish to white with yellowish midrib; stigma-lobes 3 or 4; berries white.

Type locality: Brazil.

Distribution: Uruguay to central Brazil.

Illustrations: Curtis's Bot. Mag. 69: pl. 4039; Loudon, Encycl. Pl. ed. 3. 1380. f. 19408, as *Rhipsalis brachiata*; Cycl. Amer. Hort. Bailey 4: f. 2101; Stand. Cycl. Hort. Bailey 5: f. 3377; Cact. Journ. 1: 180; Monatsschr. Kakteenk. 4: 75, as *R. saglionis*.

Plate XXVII, figure 3, is of a plant which flowered in the New York Botanical Garden in March 1912. Figure 221 is from a photograph of a flowering plant from Misiones, obtained by Dr. Rose in 1915 from Dr. Spegazzini.

3. *Rhipsalis prismatica* Rümpler in Förster, Handb. Cact. ed. 2. 884. 1885.*Hariota prismatica* Lemaire, Illustr. Hort. 10: Misc. 84. 1863.*Rhipsalis suareziana* Weber, Rev. Hort. 64: 425. 1892.*Rhipsalis tetragona* Weber, Rev. Hort. 64: 428. 1892.

Very much branched, prostrate; lower branches elongated and terete; upper branches short and somewhat angled; areoles more or less setose; flowers white; petals usually 5, obtuse; fruit small, pinkish to white, globose.

* Haworth spelled this *R. mesembryanthoides* and also *R. salicornioides*.

† The Index Kewensis gives the place of publication erroneously as Lemaire, Cact. Aliq. Nov. 39. 1838.

‡ According to the Index Kewensis, *Rhipsalis suarensis* Weber (Dict. Hort. Bois 1046. 1898) is the same.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Fruiting branch of *Rhipsalis heteroclada*.
2. Fruiting branch of same.

3. Fruiting branch of *Rhipsalis capilliformis*.
4. Fruiting branch of *Rhipsalis virgata*.

Type locality: Not cited, but Förster and Weber state that it came from Brazil.

Distribution: Brazil and northern Madagascar, but range not known.

Weber thought that *Rhipsalis tetragona* was the same as *R. prismatica* Rümpler, but because he was not certain he described it as new.

Illustration: Gartenwelt 16: 634, as *Rhipsalis suareziana*; Monatsschr. Kakteenk. 18: 74, as *R. tetragona*.

Plate xxxii, figure 3, shows a plant from Berlin which flowered in the New York Botanical Garden on November 23, 1915.



FIG. 221.—*Rhipsalis cereuscula*.

4. *Rhipsalis simmleri* Beauverd, Bull. Herb. Boiss. II. 7: 136. 1907.

Stems pendent, cylindric, 2 to 3 mm. in diameter, very much branched, the branches dichotomous or 3 or 4-verticillate, upper short and somewhat angled, quite unlike lower ones; flowers solitary, subterminal; petals white with pink tips, oblong, 6 to 8 mm. long; filaments 5 to 8 mm. long, white, filiform; style exserted, 9 mm. long; stigma-lobes ovate, reflexed, white; ovary obconic, 3 to 3.5 mm. in diameter; fruit white.

Type locality: Costa Rica.

Distribution: Costa Rica.

This species is named for Paul Simmler, chief gardener of the Boissier Collections at Geneva, Switzerland. The plant was introduced in a collection of orchids from Costa Rica and flowered in cultivation. Dr. Rose saw it when in Geneva in 1912 and obtained a small fragment, but he did not see it in flower.

Illustration: Bull, Herb. Boiss. II. 7: 137.

5. *Rhipsalis clavata* Weber, Rev. Hort. 64: 429. 1892.

Rhipsalis clavata delicatula Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 45. 1918.

Erect when young but soon hanging, often a meter long or more, much branched; joints all similar, narrowly clavate, sometimes 4-angled when young, short, 1 to 3 cm. long, deep green,

becoming brown, produced in terminal whorls of 2 to 7; areoles few, sometimes bearing 1 to 5 white hairs; flowers near end of branches, white, 1.5 cm. long; petals hardly spreading; fruit spherical, 6 mm. in diameter, white or yellowish; seeds 1.5 cm. long.

Type locality: Petropolis, in the state of Rio de Janeiro, Brazil.

Distribution: State of Rio de Janeiro.

This species is much like *Hatiora* and it was really referred to *Hariota* at one time by Weber, himself. Schumann gives only one locality for it, but Dr. Rose found it on Corcobado in Rio de Janeiro, altitude 465 meters, growing on branches of trees, and on this plant the description has been partly based. Weber's manuscript name, *Hariota clavata*, has appeared only as a synonym of this species (Monatsschr. Kakteenk. 5: 172. 1895).

Illustrations: Arch. Jard. Bot. Rio de Janeiro 2: pl. 17, as *Rhipsalis clavata delicatula*; Arch. Jard. Bot. Rio de Janeiro 1: pl. 13; Möllers Deutsche Gärt. Zeit. 25: f. 11, No. 16.

6. *Rhipsalis campos-portoana* Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 35. 1918.

Stem slender, terete, usually pendent, usually dichotomous; primary branches elongated; terminal branches in 2's or 4's, somewhat clavate, 3 to 4 cm. long; areoles few, naked; flowers terminal or usually so, white; petals about 8, slightly spreading, obtuse, up to 9 mm. long; fruit globose, 6 mm. in diameter, red.

Type locality: Serra de Itatiaya, Brazil

Distribution: Known only from the type locality.

This plant was collected by Dr. Rose and Campos Porto in July 1915 (No. 20612) and flowered in the Jardim Botânico do Rio de Janeiro in September of that year, and from this the description was drawn. Dr. Rose brought home living specimens but these have not yet flowered.

Illustration: Arch. Jard. Bot. Rio de Janeiro 2: pl. 7.

7. *Rhipsalis heteroclada* nom. nov.

Stems stiff, dark green, but purple about areoles and tips of branches, often erect in cultivation, much branched toward top of plant; branches often in verticillate clusters, much more slender than the main stem, 1 to 2 mm. in diameter; areoles small, often bearing a single bristle; flowers small, white or greenish; petals 5, obtuse, spreading or recurved; filaments about 20, white, erect; style white, sunken at base into a little cup; stigma-lobes 3, white; ovary green, about 2 mm. long; fruit globose, to 6 mm. in diameter, white.

This plant is very common in Brazilian collections, where it is planted on fruit trees. Dr. Rose found some beautiful examples in the Horto Bolanco Paulista, near São Paulo, and on Ilha Grande (Rose 20371, type).

Plate XXIII, figure 2, shows a fruiting branch obtained by Dr. Rose in Rio de Janeiro; plate XXV, figures 1 and 2, shows fruiting plants collected by Dr. Rose in Rio de Janeiro; plate XXXII, figure 1, shows a fruiting plant obtained by Dr. Rose in Rio de Janeiro.

8. *Rhipsalis capilliformis* Weber, Rev. Hort. 64: 425. 1892.

Rhipsalis gracilis N. E. Brown, Gard. Chron. III. 33: 18. 1903.

Stems and branches very slender and weak, the main branches often much elongated, the branchlets short, spreading or drooping; flowers numerous, scattered along sides of branches, cream-colored, rotate, 5 to 6 mm. broad; petals few, sometimes only 5, short and obtuse; fruit globose, naked, white or pinkish, 4 to 5 mm. in diameter; seeds very numerous.

Type locality: Not cited.

Distribution: Eastern Brazil, but not known to us in the wild state.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Rhipsalis cribrata*.
2. Fruiting branch of *Rhipsalis capilliformis*.
3. Flowering branch of same.
4. Flowering branch of same.
5. Flowering branch of *Rhipsalis teres*.

This is a very attractive little plant, often forming a dense mass of delicate branches. It is a rather shy bloomer, but grows well in damp greenhouses.

Illustration: Gartenwelt 13: 117.

Plate xxvi, figure 4, is from a plant obtained in the Botanical Garden at Brussels by Dr. Rose in 1912, which flowered and fruited in Washington in 1919; figure 3 shows a plant sent from Paris, France, which flowered in the New York Botanical Garden in 1911 (No. 14795); figure 2 is from a plant sent by R. Lamb, from Manchester, England; plate xxv, figure 3, shows a fruiting plant sent from Paris in 1901.

9. *Rhipsalis burchellii* nom. nov.

Rhipsalis cribrata Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 81. pl. 10. 1915. Not Rümpler, 1885.

Much branched, very weak, with long slender hanging branches, the branching usually dichotomous; ultimate branches usually 4 to 10 cm. long; flowers subterminal, campanulate, 10 to 12 mm. long, white; fruit turbinate, rose-colored.

This plant is very common in the forests about São Paulo. Dr. Rose collected it in the forest of Jabaquara, August 15, 1915 (No. 20857, type), and also in the Botanical Garden of Museu Paulista on August 14, 1915 (No. 20849).

This species is named for William John Burchell (1781-1863), who went to Brazil in 1825, where he made large and valuable collections.

Plate xxvii, figure 2, shows a fruiting branch taken from Dr. Rose's plant No. 20857.

10. *Rhipsalis cribrata* (Lemaire) Rümpler in Förster, Handb. Cact. ed. 2. 889. 1885.

Hariota cribrata Lemaire, Illustr. Hort. 4: Misc. 12. 1857.

Rhipsalis pendula Vöchting, Jahrb. Wiss. Bot. Leipzig 9: 371. 1873. Not Pfeiffer, 1837.

Rhipsalis penduliflora N. E. Brown, Gard. Chron. II. 7: 716. 1877.

Hariota penduliflora Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Rhipsalis cribrata filiformis Engelhardt in Möllers, Deutsche Gärt. Zeit. 18: 583. 1903.

Woody at base, much branched; branches of two forms; stems terete, elongated, at first erect, then hanging, without aerial roots; terminal branches very short, 2 to 3 cm. long, usually in whorls of 2 to 20; areoles small, often with 1 or 2 small setae; flowers generally terminal, pendulous, white or cream-colored, 8 to 10 mm. long; petals usually 5 to 7, obtuse, drying yellow; filaments erect, numerous, white, salmon-colored at base; style white; stigma-lobes 3 or 4, spreading, white; ovary naked; fruit small, globose, 2 to 3 mm. in diameter, pinkish, terminated by the old perianth.

Type locality: Brazil.

Distribution: States of Minas Geraes, Rio de Janeiro, and São Paulo, Brazil.

This species was introduced into Europe in 1856 from Brazil, as some of the other species have been, through sendings of orchids, where it was discovered by Lemaire, and when it flowered the following year it was named and described by him.

Hariota penduliflora (Monatsschr. Kakteenk. 1: 69. 1891) is listed but not described.

Rhipsalis penduliflora laxa, referred to by Schumann (Martius, Fl. Bras. 4: 276. 1890), comes from the gardens at Kew.

Illustrations: Möllers Deutsche Gärt. Zeit. 18: 585, as *Rhipsalis cribrata filiformis*; Blühende Kakteen 1: pl. 27, A; Arch. Jard. Bot. Rio Janeiro 1: pl. 9, as *R. penduliflora*.

Plate xxiii, figure 3, shows a fruiting branch collected by Dr. Rose in Rio de Janeiro in 1915; plate xxvi, figure 1, shows a flowering branch obtained by Dr. Rose in Rio de Janeiro.

11. *Rhipsalis cassutha* * Gaertner, Fruct. Sem. 1: 137. 1788.


Cassutha filiformis Miller, Gard. Dict. ed. 8. 1768. Not Linnaeus, 1753.

Cactus parasiticus Lamarck, Encycl. 1: 541. 1783. Not Linnaeus, 1768.

Cactus pendulus Swartz, Prodr. 77. 1788.

*The original spelling given by Gaertner is as above. The usual spelling, however, is *R. cassutha*.

- Rhipsalis parasitica* Haworth, Syn. Fl. Succ. 187. 1812.
*Cactus caripensis** Humboldt, Bonpland, and Kunth, Nov. Gen. at Sp. 6: 66. 1823.
Cereus caripensis De Candolle, Prodr. 3: 467. 1828.
Rhipsalis cassytha dichotoma De Candolle, Prodr. 3: 476. 1828.
Rhipsalis cassytha mauritiana † De Candolle, Prodr. 3: 476. 1828.
Rhipsalis cassytha mociniana De Candolle, Prodr. 3: 476. 1828.
Rhipsalis cassytha hookeriana De Candolle, Prodr. 3: 476. 1828.
Rhipsalis cassytha swartziana De Candolle, Mém. Mus. Hist. Nat. Paris 17: 80. 1828.
Rhipsalis dichotoma G. Don, Hist. Dichl. Pl. 3: 176. 1834.
Rhipsalis hookeriana G. Don, Hist. Dichl. Pl. 3: 176. 1834.
Rhipsalis cassythoides G. Don, Hist. Dichl. Pl. 3: 176. 1834.
Rhipsalis cassutha pendula Salm-Dyck in Pfeiffer, Enum. Cact. 134. 1837.
Rhipsalis undulata Pfeiffer, Enum. Cact. 136. 1837.
Hariota cassytha Lemaire, Cact. Gen. Nov. Sp. 75. 1839.
Cereus parasiticus Haworth in Steudel, Nom. ed. 2. 1: 335. 1840.
Rhipsalis aethiopica Welwitsch, Journ. Linn. Soc. Bot. 3: 152. 1859.
Rhipsalis minutiflora Schumann in Martius, Fl. Bras. 4²: 271. 1890.
Hariota parasitica Kuntze, Rev. Gen. Pl. 1: 262. 1891.
Rhipsalis comorensis Weber, Rev. Hort. 64: 424. 1892.
Rhipsalis zanzibarica ‡ Weber, Rev. Hort. 64: 425. 1892.

FIG. 222.—*Rhipsalis*.

Epiphytic or saxicolous, usually growing on trunk or branches of large trees, hanging in large clusters, 1 to 9 meters long, the branches weak and pendent; branches when young bearing 5 to 9 white bristles at the areoles, when old naked, terete, sometimes producing aerial roots, often only 3 mm. in diameter, light green, usually growing from tips of other branches, generally in pairs but sometimes in clusters of 6 or 8; flowers lateral, solitary, small, greenish in bud, sometimes subtended by a single bristle; petals 2 mm. long, cream-colored; stamens borne on disk; ovary exserted; fruit naked, white or pink, maturing a few days after flowering, globose, 5 mm. in diameter.

Type locality: Not cited.

Distribution: Florida, Mexico, Central America, West Indies, Panama to Dutch Guiana, eastern and southern Brazil, Colombia, Ecuador, Bolivia, and Peru, also in Ceylon and tropical Africa.

The fruit of *Rhipsalis cassutha*, while usually white, is sometimes described as red or pinkish. Hooker, in his Exotic Flora, figured and described the fruit as flesh-colored. Weber, who received a red-fruited form from Costa Rica, has named it variety *rhodocarpa* (Dict. Hort. Bois 1046. 1898). In the West Indies the plants inhabit moist districts and are most abundant in forests, but in the vicinity of Matanzas, Cuba, occur on cliffs.

*This name was written *Cactus garipensis* by Kunth (Syn. Pl. Aeq. 3: 370. 1824) and is so listed in the Index Kewensis.

† De Candolle gives *Cactus pendulinus* Sieber (Fl. Maur. 2. n. 259) as a synonym of this variety.

‡ Schumann (Gesamtb. Kakteen 623) spells the name *Rhipsalis sansibarica*.



M. E. Eaton del.

1. Flowering and fruiting branch of *Rhipsalis cassutha*.
2. Fruiting branch of *Rhipsalis burchelli*.
3. Flowering branch of *Rhipsalis cereuscula*.

A. Hoen & Co. Baltimore

Hitherto unknown wild within the continental United States, the plant was found on August 5, 1923, by C. A. Mosier on trees in Wallenstein's Hammock, Dade County, Florida.

Cactus cassythoides Mociño and Sessé was given by De Candolle (Prodr. 3: 476. 1828) as a synonym of *R. cassytha mociniana*.

Löfgren (Arch. Jard. Bot. Rio de Janeiro 2: 40. pl. 11. 1918) has figured and described as new a plant under the name of *Rhipsalis cassythoides* which may belong here. The name had already been used by Don and we have referred it as a synonym of *R. cassutha*.

Cactus epidendrum Linnaeus (Amen. Acad. 8: 257. 1785) is without description and has been referred to *Rhipsalis undulata*. It was from Surinam.

Cereus bacciferus (Hemsley, Biol. Centr. Amer. Bot. 1: 548. 1880) appears only as a synonym of *Rhipsalis cassutha*.

Cassytha baccifera Miller (De Candolle, Prodr. 3: 476. 1828) and *C. polysperma* Aiton (Gaertner, Fruct. Sem. Pl. 1: 137. 1788) are known in synonymy only.

Rhipsalis pendula Hortus (Pfeiffer, Enum. Cact. 133. 1837) occurs only as a synonym.

Rhipsalis caripensis Weber is listed as one of the synonyms of this species by Schumann (Gesamtb. Kakteen 622. 1898).

Rhipsalis cassytha vars. *major* and *pilosiuscula* (Salm-Dyck, Hort. Dyck. 228. 1834) and var. *tenuior* (Schumann, Monatsschr. Kakteenk. 1: 78. 1891) are only names. The first has been referred to *R. floccosa*, while the second is sometimes referred to *R. pulvinigera*.

Illustrations: De Tussac, Fl. Antill. 3: pl. 22, as *Cactus pendulus*; Plunkenet, Phyt. Pl. 172, f. 2. 1692, as *Cuscuta baccifera*, etc.; De Candolle, Mém. Mus. Hist. Nat. Paris 17: pl. 21; Förster, Handb. Cact. ed. 2. 888. f. 121. as *Rhipsalis cassytha mociniana*; Ann. Inst. Roy. Hort. Fromont 2: pl. 3, as *R. parasitica*; Arch. Jard. Bot. Rio de Janeiro 2: pl. 11, as *R. cassythoides*; Gartenwelt 13: 117, as *R. minutiflora*; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 18, as *R. sansibarica*; (Hortus malabaricus pl. 7, *fide* Miller); Gaertner, Fruct. Sem. Pl. 1: pl. 28, f. 1; Torrey 9: 154. f. 1; Ann. Rep. Smiths. Inst. 1908: 537. f. 1; Journ. N. Y. Bot. Gard. 11: f. 23; Loudon, Encycl. Pl. 413. f. 6907; Loddiges, Bot. Cab. 9: pl. 865; Goebel, Pflanz. Schild. 1: pl. 4, f. 2; Karsten, Deutsche Fl. 887. f. 501, No. 5; ed. 2, 2: 456. f. 605, No. 5; Stand. Cycl. Hort. Bailey 2: f. 712; Nov. Act. Nat. Cur. 1: pl. 16. f. 1; Hooker, Exot. Fl. 1: pl. 2; Curtis's Bot. Mag. 58: pl. 3080; Gartenwelt 16: 633.

Plate xxvii, figure 1, shows a plant received from the Hope Botanical Garden in Jamaica. Figure 222 is from a photograph showing branches of a plant sent us from R. Lamb's collection at Manchester, England.

12. *Rhipsalis virgata* Weber, Rev. Hort. 64: 425. 1892.

Main stem or branches meter long or more, terete, about 5 mm. thick, erect or ascending but in time often pendent, often bearing aerial roots; upper branches short, 1 to 6 cm. long, terete; areoles small, a little hairy, often with a white or pinkish bristle, subtended by a minute bract; flowers borne along sides of the 2 and 3-year old branches, solitary at areoles, rotate, 8 to 10 mm. broad, open throughout day; outer perianth-segments few, ovate, greenish yellow, sometimes tinged with red; inner perianth-segments 4 to 6, oblong, cream-colored, obtuse; filaments erect, white; style white, about as long as stamens; stigma-lobes, white; ovary broader than high, crowned by a circle of scales and bearing one on the side.

Type locality: Described from a garden plant supposed to have come from Brazil.

Distribution: Eastern Brazil.

Illustration: Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 12.

Plate xxv, figure 4, shows a plant, received from M. Simon of St. Ouen, Paris, in 1901, which flowered and fruited in the New York Botanical Garden in 1916.

13. *Rhipsalis teres* (Vellozo) Steudel, Nom. ed. 2. 2: 449. 1841.

Cactus teres Vellozo, Fl. Flum. 207. 1825.

Rhipsalis conferta Salm-Dyck, Cact. Hort. Dyck. 1849. 229. 1850.

Hariota conferta Kuntze, Rev. Gen. Pl. 1: 262. 1891.

Hariota teres Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Stems erect or spreading, woody at base, 10 to 12 mm. in diameter, much branched, especially above, with 5 to 12 short ultimate branches at top of main ones; old branches terete, green or blotched with red; flowers usually several at top of short terminal branches and scattered all along the primary ones, 10 to 12 mm. broad, pale yellow; petals widely spreading; filaments and style white, erect.

Type locality: Brazil.

Distribution: States of Minas Geraes, Rio de Janeiro, and São Paulo, Brazil.

Rhipsalis floribunda Schott was given by Schumann (Martius, Fl. Bras. 4²: 274. 1890) as a synonym of this species.

Illustrations: Vellozo, Fl. Flum. 5: pl. 30, as *Cactus teres*; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 22; Garten-Zeitung 4: 182. f. 42, No. 7, as *Rhipsalis conferta*.

Plate XXVI, figure 5, shows a plant received from Kew in 1902 which flowered in the New York Botanical Garden in 1917.




FIG. 223.—*Rhipsalis shaferi*.

14. *Rhipsalis lindbergiana* Schumann in Martius, Fl. Bras. 4²: 271. 1890.

Rhipsalis erythrocarpa Schumann in Engler, Pflanzenw. Ost. Afrikas 282. 1895.

Hariota lindbergiana Kuntze, Rev. Gen. Pl. 32: 107. 1898.

Rhipsalis densiareolata Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 41. 1918.

Very much branched, hanging from tree-trunks in great festoons, 1 to 2 meters long; joints elongated, 3 to 5 mm. in diameter; areoles filled with hairs and 2 bristles; flowers numerous, lateral, pinkish; ovary naked or nearly so; fruit light red, globose, 2 to 3 mm. in diameter, 16 to 20-seeded.


Type locality: Near the city of Rio de Janeiro.

Distribution: Mountainous regions in the state of Rio de Janeiro, Brazil, and. Mount Kilman-Djaro, Africa.

Rhipsalis erythrocarpa Schumann was described from herbarium specimens, collected on Mount Kilman-Djaro, in tropical Africa. We refer it to *R. lindbergiana* Schumann in deference to the opinion of Mr. Roland-Gosselin* but we have not had specimens for study.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 2: pl. 12, as *Rhipsalis densiareolata*. Rev. Hort. 85: f. 152, in part, as *R. erythrolepis*; Martius, Fl. Bras. 4²: pl. 53; Arch. Jard. Bot. Rio de Janeiro 1: pl. 4.

*Bull. Soc. Bot. France 59: 100.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Rhipsalis neves-armondii*.
2. Flowering branch of *Rhipsalis paradoxa*.
3. Fruiting branch of *Rhipsalis pulvinigera*.

4. Flowering branch of *Rhipsalis tucumanensis*.
5. Flowering branch of same.

Plate XXI, figure 4, shows a branch from the plant collected by Dr. Rose on Tijuca in 1915 (No. 21 174); figure 2 shows another plant collected by him in Rio de Janeiro, Brazil (No. 20309).

15. *Rhipsalis shaferi* sp. nov.

Stems at first stiff, erect or ascending, afterwards spreading or procumbent, 4 to 5 mm. thick, terete, green or more or less purplish at tips; juvenile and lower branches often bearing several bristles at areoles; upper branches without bristles or with a single appressed one; scales subtending the areoles small but broad; flowers numerous, scattered all along side of branch, solitary (rarely in pairs) at areoles, small, rotate, greenish white, 8 to 10 mm. broad; petals 5 or 6, short-oblong, obtuse; filaments greenish, erect; stigma-lobes, white; ovary not sunken in branch; fruit small, globose, 2 to 3 mm. in diameter, white or sometimes tinged with pink.

Collected by John A. Shafer on trees at Asunción, Paraguay, March 18, 1917 (No. 139), on trees at Trinidad, Paraguay, March 17, 1917 (No. 134, type), again in Paraguay (Nos. 145 and 147), and on trees at Posados, Misiones, Argentina (No. 131).

Plate XXIV, figure 7, shows a branch in flower; plate XXI, figure 3, shows a branch in fruit of the type which flowered in the New York Botanical Garden in 1921. Figure 223 is from a photograph of Shafer's No. 131, which flowered and fruited in Washington in 1921.

16. *Rhipsalis fasciculata* (Willdenow) Haworth, Suppl. Pl. Succ. 83. 1819.

Cactus fasciculatus Willdenow, Enum. Pl. Suppl. 33. 1813.

Rhipsalis horrida Baker, Journ. Linn. Soc. 21: 347. 1884.

Rhipsalis madagascarensis Weber, Ind. Sem. Hort. Paris 1889; Rev. Hort. 64: 424. 1892.

Hariota fasciculata Kuntze, Rev. Gen. Pl. 1: 262. 1891.

Hariota horrida Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Stems woody, terete, much branched; branchlets clavate to cylindric, faintly ribbed when old, 4 mm. in diameter, with numerous areoles, each with a cluster of fragile hairs 3 to 4 mm. long; flowers lateral but not described; ovary not sunken in the branch; fruit globose, small, bearing a few areoles, these pubescent and setose.

Type locality: Not cited.

Distribution: Brazil and Madagascar.

We have studied Madagascan specimens of this plant sent from Kew and one sent from Bahia, Brazil, to Dr. Rose by L. Zehntner in 1920. De Candolle (Plantes Grasses 1: pl. 59) states that it occurred in Santo Domingo. Roland-Gosselin * says that it inhabits American Islands; our very extensive explorations in the West Indies have failed to discover it. The Brazilian plant differs only from the Madagascan by having fewer hairs at the areoles.

Rhipsalis pilosa Weber is listed by Schumann (Martius, Fl. Bras. 4²: 300. 1890) with the statement that it occurs in P. Rebut's Catalogue without description; A. Berger in a letter (dated March 7, 1920) states that this name is said to be a synonym of *R. madagascarensis*. It is illustrated (Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 20). *Rhipsalis madagascarensis dasycerca* Weber is listed by R. Lamb (Collection of Cacti 73. 1908.)

Illustrations: De Candolle, Pl. Succ. 1: pl. 59, as *Cactus parasiticus*; Curtis's Bot. Mag. 58: pl. 3079; Gartenwelt 13: 117, Ann. Inst. Roy. Hort. Fromont 2: pl. 1, f. G, as *R. fasciculata*; Loudon, Encycl. Pl. 413. f. 6908. as *R. parasitica*.

17. *Rhipsalis pulchra* Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 75. 1915.

Stems much branched, often pendent; branches often in whorls of 3's or 4's, 3 to 4 mm. in diameter, bright green; areoles minute, reddish; flowers few, usually from near the tips of terminal branches, purplish red, large, 12 to 14 mm. long; petals oblong, obtuse; stigma-lobes white; ovary purplish red.

Type locality: Serra da Mantiqueira, Brazil.

Distribution: State of Rio de Janeiro.

* Bull. Soc. Bot. France 59: 99.

Our living specimens came from the Organ Mountains, Rio de Janeiro, Brazil, obtained by J. N. Rose through Ph. Luetzelburg, September 21, 1915 (No. 21157).

Dr. Rose examined the type collected by A. O. Darby in 1915 in the Museu Paulista and obtained a fragment of it through the kindness of the Director.

Rhipsalis pulcherrima Löfgren (Monatsschr. Kakteenk. 9: 136. 1899) seems to have been the name first given to this plant.

Illustration: Arch. Jard. Bot. Rio de Janeiro 1: pl. 5.

Plate xxxi, figure 2, shows a flowering branch of the plant obtained by Dr. Rose in 1915 which flowered in the New York Botanical Garden in 1918 (No. 21151).

18. *Rhipsalis lumbricoides* Lemaire, Illustr. Hort. 6: Misc. 68. 1859.

Cereus lumbricoides Lemaire, Cact. Gen. Nov. Sp. 60. 1839.

Rhipsalis sarmentacea Otto and Dietrich, Allg. Gartenz. 9: 98. 1841.

Lepismium sarmentaceum Vöchting, Jahrb. Wiss. Bot. Leipzig 9: 399. 1873.

Hariota lumbricalis Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Hariota sarmentacea Kuntze, Rev. Gen. 3: 107. 1898.



FIG. 224.—*Rhipsalis lumbricoides*.

Stems terete when glowing, but angled when dormant, 3 to 4 meters long, about 6 mm. thick rooting freely, much branched; young growth with 5 to 10 white bristles from each areole, usually spreading, but old branches naked; flowers white to cream-colored, sometimes tinged with green; petals few, often only 5, lanceolate, acute, 10 to 12 mm. long, acuminate; style slender, greenish, longer than the stamens; stigma-lobes 4, spreading, greenish; ovary naked; fruit white.

Type locality: Montevideo, Uruguay.

Distribution: Uruguay and Paraguay, also probably southern Brazil. Hooker says that it is a native of Buenos Aires, but this is doubtless an error.

This plant flowered in Washington on March 16, 1915. Schumann's drawing of the flower is not very good.

Rhipsalis sarmentosa (Monatsschr. Kakteenk. 4: 46. 1894) and *R. larmentacea* (Illustr. Hort. 6: 88. 1859) are misspellings for *R. sarmentacea*.

According to Lemaire (Cact. Gen. Nov. Sp. 60. 1839) *Cereus flagelliformis minor* Salm-Dyck (Hort. Dyck. 64. 1834) belongs here. Grisebach (Symb. Fl. Argem. 139) referred *Cereus donkelaarii* here.

Illustrations: Martius, Fl. Bras. 4: pl. 59; Curtis's Bot. Mag. 85: pl. 5136; Dict. Gard. Nicholson 4: 598. f. 60; Suppl. 635. f. 646; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 69, D, E; Gard. Chron. III. 2: 465. f. 95; Watson, Cact. Cult. 232. f. 90; ed. . f. 66, as *Rhipsalis sarmentacea*; Schumann, Gesamtb. Kakteen 633. f. 98, F; Arch. Jard. Bot. Rio de Janeiro 1: pl. 3; Gartenwelt 13: 117.



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Rhipsalis floccosa*.
2. Flowering branch of same.
3. Fruiting branch of *Rhipsalis puniceo-discus*.

4. Flowering branch of *Rhipsalis gibberula*.
5. Branch of *Rhipsalis dissimilis*.
6. Flowering and fruiting branch of same.

Figure 224 is from a photograph taken by H. Buch which was given to Dr. Rose when he was in La Plata, Argentina, in 1915.

19. *Rhipsalis aculeata* Weber, Rev. Hort. **64**: 428. 1892.

Stems terete, 3 to 4 mm. in diameter, somewhat angled and roughened in dried specimens; areoles close together, bearing wool and 8 to 10 appressed white bristles or spines; fruit not immersed, globose, 7 to 8 mm. in diameter, dark purple to nearly black, either naked or with 3 or 4 hairy areoles.

Type locality: Catamarca, Argentina.

Distribution: Northern Argentina, in the provinces of Catamarca and Tucuman.

A round-stemmed species collected by Otto Kuntze on the Sierra de Santa Cruz, Bolivia, and labeled *Hariota sarmentacea* may belong here.

This species is described by Schumann as 8 to 10-ribbed, but no ribs are shown in growing plants; in drying the branches are somewhat angled but one could hardly describe them as ribbed. Dr. Shafer made a single collection of this plant at Tucuman in 1917 (No. 92); part of this material is living in the New York Botanical Garden. Dr. Rose also obtained a specimen through one of his Argentina correspondents from Catamarca.

Plate xxiv, figure 8, is from Dr. Shafer's plant mentioned above.

20. *Rhipsalis grandiflora* Haworth, Suppl. Fl. Succ. **83**. 1819.

Cactus funalis Sprengel, Syst. **2**: 479. 1825.

Cactus cylindricus Vellozo, Fl. Flum. **207**. 1825. Not Lamarck, 1783. Not Ortega, 1800.

Rhipsalis funalis Salm-Dyck in De Candolle, Prodr. **3**: 476. 1828.

Hariota funalis Lemaire, Cact. Gen. Nov. Sp. **74**. 1839.

Rhipsalis cylindrica Steudel, Nom. ed. **2**: 448. 1841.

Hariota cylindrica Kuntze, Rev. Gen. Pl. **1**: 262. 1891.

Hariota grandiflora Kuntze, Rev. Gen. Pl. **1**: 262. 1891.

Rhipsalis robusta Lindberg, Monatsschr. Kakteenk. **6**: 53. 1896. Not Lemaire, 1860.

Rhipsalis hadrosoma Lindberg, Monatsschr. Kakteenk. **6**: 96. 1896.

Branches divaricate, often reddish, especially about the areoles, stout, 8 to 10 mm. in diameter; flowers numerous, scattered all along branches, 12 mm. long, 2 cm. broad, light rose or cream-colored; sepals reddish; petals few, oblong, obtuse, widely spreading; anthers and style white; stigma-lobes 4, white; fruit naked, purplish, 6 to 7 mm. in diameter.

Type locality: Not cited.

Distribution: State of Rio de Janeiro, Brazil.

We have not seen the type specimen of this species, but through the kindness of Mr. N. E. Brown of Kew we have seen a photograph of Haworth's specimens, which are the same as the species here described. Haworth's plant was received from Brazil in 1816, sent by Messrs. Bowie and Cunningham.

Rhipsalis calamiformis (Pfeiffer, Enum. Cact. **135**. 1837) was published as a synonym of *R. funalis*.

Walpers gives *Rhipsalis funalis gracilior* Pfeiffer (Repert. Bot. **2**: 279. 1843) as a synonym.

Illustrations: Gartenwelt **13**: 117; Watson, Cact. Cult. **228**. f. 89; ed. **3**. f. 65; Amer. Gard. **11**: 465; Dict. Gard. Nicholson **3**: 289. f. 365; Gartenflora **42**: 234. f. 48; Link and Otto, Icon. Pl. Rar. pl. **38**, as *Rhipsalis funalis*; Vellozo, Fl. Flum. **5**: pl. 31, as *Cactus cylindricus*; Monatsschr. Kakteenk. **6**: 55, as *R. robusta*; Blühende Kakteen **3**: pl. 141; Monatsschr. Kakteenk. **7**: 151. f. 1 to 8; Arch. Jard. Bot. Rio de Janeiro **1**: pl. 7, as *R. hadrosoma*; Curtis's Bot. Mag. **54**: pl. 2740; Schumann, Gesamtb. Kakteen **633**. f. 98, A; Martius, Fl. Bras. **4**²: pl. 54; Monatsschr. Kakteenk. **7**: 151. f. 9 to 11; Arch. Jard. Bot. Rio de Janeiro **1**: pl. 6.

Plate xxxi, figure 3, shows a plant collected by Dr. Rose near Rio de Janeiro in 1915 (No. 20746) which flowered in the New York Botanical Garden in 1918; figure 1 is of a plant which also flowered in the New York Botanical Garden, April 3, 1912; plate xxi, figures 1 and 6, shows the flowers and fruit of specimens sent by Alwin Berger in 1908.

21. *Rhipsalis megalantha* Löfgren, Monatsschr. Kakteenk. **9**: 134. 1899.*Rhipsalis novaesii* Gürke, Monatsschr. Kakteenk. **19**: 12. 1909.

Plants stout, up to 1 cm. thick, at first erect but in time spreading or with pendent branches, dull green, often spotted with purple; areoles rather prominent, especially after flowering; flowers large, 4 cm. broad; petals 8 to 12, oblong, often shortly acuminate or obtuse, white; filaments erect, orange at base, rose-colored above; style thick, longer than the stamens; stigma-lobes 6 to 8; fruit surrounded with white hairs, rather small, 6 mm. in diameter, white or tinged with red; seeds nearly black.

Type locality: Island of São Sebastião, Brazil.

Distribution: Known only from the type locality, an island off the coast of Brazil, belonging to the state of São Paulo.

This plant is known wild only from the collection of Dr. Löfgren, but is now widely found in cultivation, sometimes under the names *Rhipsalis grandiflora* or *R. novaesii*. It has the largest flower of any species of *Rhipsalis*.

Illustrations: Blühende Kakteen **2**: pl. 116; Monatsschr. Kakteenk. **19**: 13, as *Rhipsalis novaesii*; Monatsschr. Kakteenk. **9**: 137; Schumann, Gesamtb. Kakteen Nachtr. 147. f. 35; Arch. Jard. Bot. Rio de Janeiro **1**: pl. 8.

Plate xxiv, figure 3, shows a fruiting branch obtained by Dr. Rose in Rio de Janeiro, Brazil, in 1915 (No. 20400).

22. *Rhipsalis leucorhaphis* Schumann, Monatsschr. Kakteenk. **10**: 125. 1900.

Epiphytic, much branched, about dm. long, rooting abundantly along the branches, jointed, 5 to 8 mm. in diameter, terete or showing 4 or 5 ribs in herbarium specimens; bristles 1 to 5, appressed, early deciduous; areoles subtended by an ovate papery bract; flowers white, nodding, large, 1.5 cm. long; petals only slightly spreading; filaments purplish or white with orange-colored base; stigma-lobes 3 or 4, greenish, spreading; ovary not sunken in the branch; fruit globose, bright red, 6 to 8 mm. in diameter; seeds numerous, brown.

Type locality: Estancia Tagatiya, Paraguay.

Distribution: Paraguay and northern Argentina.

We did not know this species until it was brought back by Dr. Shafer in 1917 from Paraguay, where he obtained good specimens; he also found it abundant in northern Argentina. Like many of the other species it grows in various situations, sometimes sprawling over rocks or growing on forest trees. One of his living plants fruited in the New York Botanical Garden and from this we have drawn part of our description.

Plate xxiv, figure 1, shows the plant in flower, and figure 2 shows it in fruit, collected by Dr. Shafer at Trinidad, Paraguay (No. 143).

23. *Rhipsalis loefgrenii* nom. nov.*Rhipsalis novaesii* Löfgren, Arch. Jard. Bot. Rio de Janeiro **1**: 69. 1915. Not Gürke, 1909.

Stems long and slender, rooting freely all along stem, pale green to purple, terete, 3 mm. in diameter; areoles small, subtended by a large scarious bract with appressed hairs in axils when young; flowers very numerous, 12 to 15 mm. long, white, campanulate; filaments purplish at base; fruit purplish, 5 to 8 mm. in diameter.

Type locality: Near Campinas, Brazil.

Distribution: Brazil.

Dr. Rose saw the Löfgren type in the Botanical Garden at Rio de Janeiro and obtained living and herbarium specimens of the plant. Dr. Shafer also obtained living specimens from Löfgren in 1917.

Unfortunately, Löfgren's name was given to another plant by Gürke and for this reason we have renamed it in honor of Dr. Alberto Löfgren (1854-1918), who long studied this genus and published an excellent monograph of it in 1915.

Illustration: Arch. Jard. Bot. Rio de Janeiro **1**: pl. 2, as *Rhipsalis novaesii*.



M. E. Eaton del.

A. Hoehn & Co. Baltimore

1. Flowering branch of *Rhipsalis gonocarpa*.
2. Flowering branches of *Rhipsalis warmingiana*.
3. Fruiting branch of *Rhipsalis tonduzii*.

4. Fruiting branch of *Rhipsalis trigona*.
5. Flowering branch of *Rhipsalis pentaptera*.
6. Fruiting branch of same.

Figure 225*a* shows two branches with a single fruit 1.33 times natural size; figure 225*b* shows a branch twice natural size; figure 225*c* shows one of the bracts which subtend the areoles, 4 times natural size, all drawn from plants obtained by Dr. Shafer from Dr. Löfgren in 1917 and since grown in the New York Botanical Garden.

24. *Rhipsalis nevesarmondii*
Schumann in Martius, Fl.
Bras. 4²: 284. 1890.

? *Rhipsalis rigida* Löfgren,
Arch. Jard. Bot. Rio de
Janeiro 1: 93. 1915.

Stems elongated, much branched, and hanging from trees in large clusters; branches arranged in whorls of 3 to 10, 4 to 5 mm. thick, terete, elongated, deep green; flowers widely spreading, 2 cm. broad, white to cream-colored; petals about 12, acute; style erect, white; stigma-lobes 5, white; ovary sunken in the branch; fruit globose, red, 10 mm. in diameter; seeds brown.




FIG. 225.—*Rhipsalis loefgrenii*. *a*, fruiting branch; *b*, tip of branch; *c*, bract.

Type locality: Mount Tijuca, Rio de Janeiro, Brazil.

Distribution: Rio de Janeiro, Brazil.

There has long been much uncertainty regarding this species and Dr. Rose, during his trip to South America, in 1915, endeavored to solve the problem. He first visited one of the three localities mentioned in the original description, namely Tijuca, a mountain near Rio de Janeiro. Here he found two species which belonged to the same group, *Rhipsalis grandiflora* and *R. pulvinigera*. He then visited the herbarium of the Museo Nacional, where he found specimens of *R. neves-armondii*. Unfortunately, they did not bear an original label but one doubtless written after the appearance of the description in the Flora Brasiliensis for the three localities mentioned therein. After studying this material carefully, he visited the mountain region just above Tijuca, namely Pica Popagaya, where he feels certain he has collected the true form, although the joints are more terete and the flowers are pure white instead of yellow; it is a singular *Rhipsalis* and a very shy bloomer. A second visit was then made to Tijuca, but lower down on the mountain, and here he again found this species.

Illustrations: Martius, Fl. Bras. 4²: pl. 56; Arch. Jard. Bot. Rio de Janeiro 1: pl. 19; Blühende Kakteen 2: pl. 80, A.

Plate xxviii, figure 1, shows a flowering plant collected by Dr. Rose at the type locality in 1915 (No. 20673), which flowered in the New York Botanical Garden in 1916; Plate xxiv, figures 4 and , show branches from the same plant, in fruit.

25. *Rhipsalis pittieri* sp. nov.

Epiphytic, resembling in habit *Rhipsalis cassutha*; branches 5 to 6 mm. in diameter, dull green, terete; petals greenish yellow, 5 to 6 mm. long; ovary sunken in the stem, surrounded by white hairs; fruit maturing very slowly, white; seeds black.

Collected by H. Pittier near Hacienda Koster, Borburata, near Puerto Cabello, Venezuela, in 1913 (No. 6467), and flowered first in Washington in the fall of 1914 (October 16), the fruit maturing March 16, 1915. The plant has repeatedly flowered since. This species

is perhaps nearest *Rhipsalis floccosa*, from Brazil, and is the most northern representative of the Series *Floccosae*.

Plate xxiv, figure 6, is of a fruiting specimen of the type plant.

26. *Rhipsalis pulvinigera* Lindberg, *Gartenflora* **38**: 186. 1889.

Rhipsalis funalis minor Pfeiffer, *Enum. Cact.* 135. 1837.

Plant epiphytic, rather stout, at first erect but in time hanging, and then sometimes 3 to 5 meters long, the branches dull green with purple about the areoles, 5 to 7 mm. in diameter; terminal branches often in whorls of 3 to 5; flowers at first white, in age yellowish, 2 cm. broad; ovary sunken in the branch; fruit globose, red, 8 mm. in diameter.

Type locality: Brazil.

Distribution: In the coastal mountains of central Brazil.

Schumann gives *Rhipsalis grandiflora minor* (Gesamtb. Kakteen 644. 1898) as a synonym of this species, but he evidently meant *R. funalis minor*.

Rhipsalis cassytha pilosiuscula Salm-Dyck (*Hort. Dyck.* 228. 1834), although never described, probably is to be referred here.

Illustrations: *Gartenflora* 42 f. 48, as *Rhipsalis funalis*; *Gartenflora* **38**: f. 33, 34; Rümpler, *Sukkulenten* 210. f. 119; 211. F. 120; *Rev. Hort.* **85**: f. 152, in part.

Plate xxviii, figure 3, is from a plant collected by Dr. Rose near Rio de Janeiro in 1915, which fruited in the New York Botanical Garden in 1915 (No. 43060).

27. *Rhipsalis floccosa* Salm-Dyck in Pfeiffer, *Enum. Cact.* 134. 1837.

Hariota floccosa Lemaire, *Cact. Gen. Nov. Sp.* 75. 1839.

Rhipsalis rugulosa Lemaire, *Illustr. Hort.* 8: after pl. 293. 1861.

Hariota rugosa Kuntze, *Rev. Gen. Pl.* 1: 263. 1891.

Stems slender, 5 to 8 mm. in diameter, much branched, at first erect, becoming pendent; branches alternate; flowers lateral, 2 cm. broad, white, tinged with yellow, surrounded by a tuft of wool; ovary sunken in the branch; fruit globose, 5 mm. in diameter, rose-colored or nearly white.

Type locality: Not cited.

Distribution: Brazil.

Rhipsalis cassytha major Salm-Dyck (Pfeiffer, *Enum. Cact.* 134. 1837), a synonym only, is referred here by Pfeiffer.

Hariota floccosa Cels was used as a synonym by Förster (*Handb. Cact.* 458. 1846), but was not technically published until 1891.

Illustration: *Gartenflora* **38**: 185. f. 35.

Plate xxix, figure 1, shows a flowering branch from a specimen sent by Mr. Lamb from Manchester, England, in 1914, and figure 2 shows a flowering branch collected by Dr. Rose in Brazil in 1915 which flowered in the New York Botanical Garden on February 24, 1922.

28. *Rhipsalis tucumanensis* Weber, *Rev. Hort.* **64**: 426. 1892.

Hariota tucumanensis Kuntze, *Rev. Gen. Pl.* 3: 107. 1898.

Epiphytic on forest trees, when young setose, but soon naked, much branched; branches often pendent, sometimes in whorls of 4, 4 to 10 mm. in diameter, when young nearly terete, bright green with a red spot at the areoles, when old angled, yellowish green; flowers one from an areole, 15 to 18 mm. in diameter, rosy white to cream-colored; sepals, white but rose-colored on the back; petals 8, ovate-lanceolate; stamens numerous, white, spreading, much shorter than petals; style white; stigma-lobes 4 or 5; ovary sunken in the branch, surrounded by a tuft of wool; fruit described as white tinged with red, but often red or pinkish, 8 to 10 mm. broad.

Type locality: Tucuman, Argentina.

Distribution: Tucuman to Catamarca, Argentina, and perhaps Bolivia and Paraguay.



M. E. Eaton del.

1. Flowering branch of *Rhipsalis grandiflora*.
2. Flowering branch of *Rhipsalis pulchra*.
3. Flowering branch of *Rhipsalis grandiflora*.

A. Hoen & Co. Baltimore

This species is common in northern Argentina, where it was repeatedly collected by Dr. Shafer in 1917.

Of this relationship, but perhaps specifically distinct, is the plant sent by M. Bang (No. 2323) from Coripati, Yungas, Bolivia, distributed as *Rhipsalis salicornioides*. Here we have tentatively referred K. Fiebrig's plant (No. 5801) from the Upper Paraná, Paraguay.

Of plate xxviii, figures 4 and 5 show flowering and fruiting branches from Dr. Shafer's collection from Calilegua, Argentina (Nos. 55 and 68), painted at the New York Botanical Garden, May 24, 1922.




FIG. 226.—*Rhipsalis stilcata*. Reduced.

FIG. 227.—*Rhipsalis gibberula*. $\times 0.5$.

29. *Rhipsalis gibberula* Weber, Rev. Hort. 64: 426. 1892.

Stems 3 to 6 mm. thick, yellowish green, with dichotomous or trichotomous branches or sometimes with terminal whorls of 6; areole small; buds obtuse, pinkish, hairy when in flower; flowers scattered along branches toward tip, white to pale pink, 8 to 9 mm. long, 12 to 15 mm. broad; petals not widely spreading (at least in our specimen); stigma-lobes 3 to 6, white; fruit white, somewhat depressed, 8 to 10 mm. in diameter, 7 to 8 mm. high, the base sunken in the branch.

Type locality: Brazil.

Distribution: Organ Mountains, Brazil.

The species was described from plants brought to Paris from Brazil in 1887, their habitat not recorded, but Dr. Rose traced it to the Organ Mountains in 1915 and his plant flowered in the New York Botanical Garden in February 1921 (No. 21161). In 1902 a specimen was sent from Paris to the New York Botanical Garden and one specimen was obtained from R. Lamb, Superintendent of Parks at Manchester, England, in 1914, but neither has done well in cultivation.

Plate xxix, figure 4, is from a plant collected by Dr. Rose in the Organ Mountains in 1915, which flowered in the New York Botanical Garden, February 17, 1921. Figure 227 shows the plant received from Paris in 1902 which flowered in the New York Botanical Garden on March 6, 1917.

30. *Rhipsalis puniceo-discus* G. A. Lindberg, Gartenflora 42: 233. 1890.

Rhipsalis foveolata Weber, Dict. Hort. Bois 1047. 1898. According to Roland-Gosselin.

Rhipsalis chrysocarpa Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 94. 1915.

? *Rhipsalis chrysantha* Löfgren, Arch. Jard. Bot. Rio de Janeiro 1: 99. 1915.

Branches slender, almost filiform, hanging, pale green when young, freely rooting; branches in terminal whorls, often as many as 6; flowers large, 1.5 cm. long, white; perianth-segments widely spreading; stamens orange-colored, at least at base; fruit at first dark red but in age golden yellow.

Type locality: Not cited.

Distribution: Brazil.

This plant first passed in living collections as *R. funalis gracilis* (Gartenflora 42: 233. 1893.)

Dr. Löfgren gave Dr. Rose a cutting of the original plant of *Rhipsalis chrysantha*.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 1: pl. 20, as *Rhipsalis chrysocarpa*; Rev. Hort. 79: 106. f. 33, as *R. foveolata*; Gartenflora 42: 235. f. 49; Arch. Jard. Bot. Rio de Janeiro 1: pl. 21.

Plate XXIX, figure 3, shows a plant also brought by Dr. Rose from Brazil (No. 20662) which flowered and fruited in the New York Botanical Garden, March 7, 1921.

31. *Rhipsalis dissimilis* (G. A. Lindberg) Schumann in Martius, Fl. Bras. 4: 286. 1890.

Lepismium dissimile G. A. Lindberg, Gartenflora 39: 148. 1890.

Rhipsalis dissimilis setulosa Weber, Rev. Hort. 64: 428. 1892.

Rhipsalis pacheco-leonii Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 38. 1918.

In clumps on large limbs of trees and freely rooting; branches very diverse, some with numerous bristly hairs from the areoles, others naked, erect, prostrate or even hanging; hairy branches with 9 very low ribs, the areoles close together, each with about 15 long white bristles; glabrous branches, 5-angled, with the areoles alternating as in *Rhipsalis paradoxa*; flower-buds red; flowers solitary, about 6 mm. broad; petals few, oblong, obtuse, widely spreading, sometimes turned back, pinkish; stamens erect, numerous, white; ovary sunken in the branch; style pinkish, erect; stigma-lobes 3 or 4, white.

Type locality: São Paulo, Brazil.

Distribution: States of São Paulo and Rio de Janeiro, Brazil.

We have referred *Rhipsalis pacheco-leonii* here after studying living specimens of *R. dissimilis* and specimens from the type collection obtained by Dr. Rose in 1915 (No. 20707).

Rhipsalis setulosa Weber (Hort. Bois Paris) was published as a synonym of *R. dissimilis* var. *setulosa*.

Illustrations: Gartenflora 39: 148. f. 36, 37, as *Lepismium dissimile*; Arch. Jard. Bot. Rio de Janeiro 2: pl. 10, as *Rhipsalis pacheco-leonii*; Curtis's Bot. Mag. 131: pl. 8013, as *R. dissimilis setulosa*; Blühende Kakteen 2: pl. 80, B; Gartenflora 40: f. 121.

Plate XXIX, figures 4 and 6, shows the two diverse forms which this plant takes, as does also plate XXXII, figures 6 and 7. The specimens were collected by Dr. Rose in the state of Rio de Janeiro in 1915 and are a part of the type material of *R. pacheco-leonii*.

32. *Rhipsalis pentaptera* Pfeiffer in Dietrich, Allg. Gartenz. 4: 105. 1836.

Hariota pentaptera Lemaire, Cact. Gen. Nov. Sp. 75. 1839.

Branches stiff, bright green, 6 to 15 mm. in diameter, strongly 6- or 6-ribbed, the ribs indented at areoles; areoles often 2 cm. apart, small, subtended by broad bracts, usually bearing 2 white bristles; flowers usually scattered along whole length of branches, opening in daytime, 1 to 4 from an areole; scales 4 or at base of corolla, broad and obtuse; petals 5, reddish on back, cream-colored on face, 4 mm. long, obtuse; stamens numerous, about 25, free from petals, white, about as long as style; style and stigma-lobes white; ovary truncate, naked; fruit 3 to 4 mm. in diameter, white, naked, or with an occasional small scale.

Type locality: Not cited. Otto says, in a note, probably Brazil.

Distribution: Southern Brazil and Uruguay.

A very common species in cultivation, flowering freely in March and April.

Hariota pentaptera Lemaire and *Rhipsalis pentagona* are given as synonyms of this species by Förster (Handb. Cact. 453. 1846).

Illustrations: Pfeiffer and Otto, Abbild. Besch. Cact. 1: pl. 17, f. 1; Goebel, Pflanz. Schild. 1: pl. 4, f. 4; Gartenwelt 13: 117; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 21; Rev. Hort. 85: f. 152, in part.

Plate XXX, figures 5 and 6, shows a plant which flowered and fruited in the New York Botanical Garden in 1912 and 1915, obtained from Paris, France, in 1902.

33. *Rhipsalis sulcata* Weber, Dict. Hort. Bois 1046. 1898.

Stems woody, sometimes 10 to 15 mm. in diameter, often long and pendent; branches elongated, the joints 2 to 3 dm. long, 5-angled, light green; areoles remote (2.5 to 5 cm. apart), usually near the



M. E. Eaton del.

1. Fruiting branch of *Rhipsalis heteroclada*.
2. Fruiting branches of *Disocactus biformis*.
3. Flowering and fruiting branch of *Rhipsalis prismatica*.
4. Flowering branch of *Rhipsalis coriacea*

5. Fruiting branch of same.
6. Fruiting branch of *Rhipsalis dissimilis*.
7. Branch of same.

A. Hoen & Co. Baltimore

center of a purple blotch; flowers solitary at the areoles, rather large, rotate, white to pinkish; ovary naked.

Type locality: Not cited.

Distribution: Not known in the wild state.

Weber found this plant in cultivation under the name of *Rhipsalis micrantha*, but it is very different from the true *R. micrantha* which comes from Ecuador.

Figure 226 shows a plant received from Paris in 1902 which flowered in the New York Botanical Garden on March 21, 1912.

34. *Rhipsalis trigona* Pfeiffer, Enum. Cact. 133. 1837.

Hariota trigona Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Stems stout, very much branched, 1.5 cm. in diameter, strongly 3-angled, the angles or ribs alternating with those of adjoining joints; flowers solitary, white to pinkish, widely spreading, sometimes 2 cm. broad; sepals usually 3, short, obtuse; petals generally 7, oblong, obtuse; filaments numerous, white; style white; stigma-lobes 4, white; ovary sunken in the branch; fruit globose, 8 to 10 cm. in diameter, red.

Type locality: Brazil.

Distribution: Brazil.

Wildeman states that the species is probably from the state of Rio de Janeiro.

Illustrations: Wildeman, Icon. Select. 5: pl. 193; Arch. Jard. Bot. Rio de Janeiro 1: pl. 23; Gartenflora 40: 38. f. 15, 16; Gartenwelt 13: 117.

Plate xxx, figure 4, shows a plant sent to Dr. Rose by R. Lamb of Manchester, England, in 1914, which flowered and fruited in the New York Botanical Garden in 1919.

35. *Rhipsalis paradoxa* Salm-Dyck, Cact. Hort. Dyck. 1844. 39. 1845.

Lepismium paradoxum Salm-Dyck in Pfeiffer, Enum. Cact. 140. 1837.

Hariota alternata Lemaire, Hort. Univ. 2: 39. 1841.

Rhipsalis alternata Lemaire, Cactées 80. 1868.

Hariota paradoxa Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Plants freely giving off aerial roots, branched, hanging in large clusters 1 meter long or more; branches in zigzag links, terminal, in pairs or in whorls of to 8, more or less spreading, 3-winged, pale; flowering areoles very woolly, setose when young, borne at upper ends of ribs; flowers sub-terminal, large, 2 cm. long, white; ovary sunken in stem; fruit not seen.

Type locality: Brazil.

Distribution: Brazil, especially near the city of São Paulo, Brazil.

The young growth is glossy green, the areoles subtended by broad round bracts. Seedling plants are very different from the adult plant; they are strongly 4-angled, with each angle bearing closely-set areoles, filled with slender bristles and showing no resemblance to the typical form; gradually as the plants grow older their mature joints take on the normal form. This plant is a prolific bloomer and in the garden of the Museo Paulista it remains in flower for three weeks.

Pfeiffer (Enum. Cact. 140. 1837) gives *Cereus pterocaulis* Hortus as a synonym of *Lepismium paradoxum* while Förster (Handb. Cact. 453. 1846) gives *Rhipsalis pterocaulis* as a synonym of *R. paradoxa*.

Lepismium alternatum Hortus (Loudon, Hort. Brit. Suppl. 3: 576. 1850) appeared as a questionable synonym of *Lepismium paradoxum*.

Illustrations: Herb. Génér. Amat. II. 2: pl. 38; Hort. Univ. 2: pl. 50, as *Hariota alternata*; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 69, A, B; Schumann, Gesamtb. Kakteen 633. f. 98, B; Karsten, Deutsche Fl. 887. f. 501, No. 4; ed. 2. 2: 456. f. 605, No. 4; Martius, Fl. Bras. 4: pl. 55, f. 1; Arch. Jard. Bot. Rio de Janeiro 1: pl. 22; Goebel, Pflanz. Schild. 1: pl. 1, f. 5; Rev. Hort. 85: f. 152, in part; Karsten and Schenck, Vegetationsbilder 1: pl. 6, f. c.

Plate xxviii, figure 2, is from a plant received from La Mortola in 1908 which flowered in the New York Botanical Garden in 1916.

36. *Rhipsalis houlletiana* Lemaire, Illustr. Hort. 5: Misc. 64. 1858.*Rhipsalis houlletii* Lemaire in Curtis's Bot. Mag. 100: pl. 6089. 1874.*Rhipsalis regnellii* Lindberg, Gartenflora 39: 119. 1889.*Hariota houlletiana* Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Stems 1 to 2 meters long, slender, terete below but flat and broad above; branches flat and thin, 1 to 5 cm. broad, tapering into a petiole-like base; margin serrate; flowers numerous, bell-shaped with a red eye; petals cream-colored, turning pale yellow, lanceolate, acute; stamens numerous; ovary not sunken in the branch, strongly 4 to 5-angled; fruit not angled, globose, red, 5 to 6 mm. in diameter.

Type locality: Not cited.

Distribution: Brazil, in the states of Minas Geraes, Rio de Janeiro, and São Paulo.

This species grows on trees in the mountains at an altitude of 1,000 meters.

Rhipsalis regnelliana appears in the general index for the Monatsschrift für Kakteenkunde (volumes 1–20) in place of *R. regnellii*.

Illustrations: Blühende Kakteen 1: pl. 56; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 69, C; Gartenflora 39: f. 29, 31 to 33; Schumann, Gesamtb. Kakteen f. 98, E; Martius, Fl. Bras. 4²: pl. 8; Möllers Deutsche Cart. Zeit. 25: 477. f. 11, No. 14, as *Rhipsalis regnellii*; Curtis's Bot. Mag. 100: pl. 6089; Gartenflora 39: f. 30; Rümpler, Sukkulente 212. f. 121, as *R. houlletii*; Rev. Hort. Belge 40 after 186, as *R. kegnelli* (in error for *R. regnellii*); Blühende Kakteen 2: pl. 111; Arch. Jard. Bot. Rio de Janeiro 1: pl. 17.

Plate xxxiii, figure 1, shows a flowering plant collected by Dr. Rose in Rio de Janeiro, Brazil, in 1915 (No. 20307), which flowered in the New York Botanical Garden in 1918; figure 2 shows a plant obtained from M. Simon of St. Ouen, Paris, in 1901, as *Rhipsalis regnellii*, which flowered in the New York Botanical Garden December 16, 1916; figure 3 shows a dissected flower and figure 4 a fruiting branch; plate xxxiv, figure 1, shows a plant with flowers obtained in Paris in 1901; figure 2 shows a flower cut through the center.

37. *Rhipsalis warmingiana* Schumann in Martius, Fl. Bras. 4²: 291. 1890.

At first erect, then spreading or hanging; branches elongated, jointed, 10 mm. wide or less, either flat or sharply 3 or 4-angled, more or less blotched or colored throughout with purple or red; flowers one at an areole, 20 mm. long, white, directed forward, the perianth-segments spreading, acute; stamens 25 to 30, white; ovary strongly angled; fruit globose, 5 to 6 mm. in diameter, dark purple to nearly black, capped by the withered flower.

Type locality: Near Lagoa Santa, Minas Geraes; two localities were cited when first described, this being the first.

Distribution: State of Minas Geraes, Brazil.

The plant has long been in cultivation, where it does well and blooms freely. Dr. Rose brought back a fresh supply from Brazil in 1915. According to Robert Lamb, the flowers have a perfume resembling that of a hyacinth.

Illustrations: Monatsschr. Kakteenk. 9: 151; Arch. Jard. Bot. Rio de Janeiro 1: pl. 18; Gartenflora 41: f. 5, 6, 7.

Plate xxx, figure 2, shows a plant from M. Simon which flowered in the New York Botanical Garden in 1912; plate xxxiv, figures 3 and 4, shows two fruiting branches received from the Berlin Botanical Garden in 1902.

38. *Rhipsalis gonocarpa* Weber, Rev. Hort. 64: 427. 1892.

Very much branched; joints narrowly lanceolate to linear, crenate, 3-angled or flattened, becoming purplish; flowers lateral, white, 15 mm. long; petals 7 or 8, lanceolate; stamens 20 to 30, white; ovary strongly 3-angled; stigma-lobes 3 or 4; fruit terete, dark purple to black, globular to short-oblong, 10 to 12 mm. long.

Type locality: São Paulo, Brazil.

Distribution: State of São Paulo, Brazil.



M. E. Eaton del.

1. Fruiting branch of *Rhipsalis houlettiana*.
2. Fruiting branch of same.

3. Flower of same.
4. Fruiting branch of same.

A. Hoen & Co. Baltimore

Schumann (Gesamtb. Kakteen 641. 1898) refers here as a synonym *Rhipsalis pterocarpa* Weber, a name which he had previously listed in the Flora Brasiliensis (4²: 300. 1890).

Plate xxx, figure 1, is from a plant sent to Dr. Rose in 1914 by R. Lamb of Manchester, England, which flowered in the New York Botanical Garden in 1920.

39. *Rhipsalis linearis* Schumann in Martius, Fl. Bras. 4²: 296. 1890.

Stems at first erect but afterwards spreading or prostrate, 6 to 8 dm. long, much branched; branches vary narrow, serrate, narrowed at base and woody; flowers white, 16 to 18 mm. long; fruit white, 5 mm. in diameter.

Type locality: Southern Brazil, but no definite locality cited. Localities in Paraguay and Argentina also cited in the original place of publication.

Distribution: Southern Brazil, Uruguay, Paraguay, and northern Argentina.

We know this species only from description.

40. *Rhipsalis micrantha* (HBK.) De Candolle, Prodr. 3: 476. 1828.

Cactus micranthus Humboldt, Bonpland, and Kunth, Nov. Gen. et Sp. 6: 65. 1823.

Hariota micrantha Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Either epiphytic and pendulous or clambering over rocks; branches 3 or 4-angled or flattened, to 8 mm. broad; areoles small, remote, bearing often 1 to bristles; flowers white, lateral, 7 mm. long including the ovary; petals cream-colored, spreading, obtuse; filaments, style, and stigmalobes white; fruit 8 to 10 mm. long, naked, white to reddish, globose; seeds black.




FIG. 228.—*Rhipsalis micrantha*. $\times 0.5$.

Type locality: Near Olleros, formerly in Ecuador, now in northern Peru.

Distribution: Ecuador and northern Peru.

Schumann describes this species as having 5 angles and cites only Humboldt's plant. The original description says 3 or 4-angled or compressed. The plant which he actually described is doubtless *Rhipsalis sulcata*, which has long passed in collections as *R. micrantha*.

Dr. Rose found this species quite common in southern Ecuador and brought back living specimens of it. The specimen in the New York Botanical Garden which came from Berlin agrees with Schumann's description.

Figure 228 shows a branch from the plant brought by Dr. Rose from southern Ecuador in 1918 (No. 23248).

41. *Rhipsalis tonduzii* Weber, Dict. Hort. Bois 1046. 1898.

Stems giving off aerial roots freely, at first erect but branches hanging, 1 cm. in diameter or less, normally 4 or 5-angled, sometimes 7-angled, but terminal branches often 3-angled or occasionally flattened and 2-angled; branches about 10 cm. long, usually terminal but always in clusters of 2 to 6, pale green; areoles close together, forming notches in the branch; flowers small, 22 mm. long, white; ovary exserted (Schumann says immersed); fruit globose, short-oblong, white, 7 to 10 mm. long, usually on upper half of terminal branches, resembling fruit of *Rhipsalis cassutha* but much longer, sometimes abortive and covered with hairs, thus resembling a small chestnut-bur, perhaps the result of insect stings; seeds oblong, numerous, black.

Type locality: Costa Rica.

Distribution: Costa Rica but range unknown.

This species flowered in Washington in March 1912, in June 1919, and again in April 1920; fruit was obtained July 31, 1919, and in April 1920.

Plate xxx, figure 3, shows a branch of a plant brought back from Costa Rica by Dr. Maxon in 1906.

Of this relationship is the following:

RHIPSALIS WERCKLEI Berger, Monatsschr. Kakteenk. **16**: 64. 1906.

Epiphytic, much branched, hanging, 3 to 6 dm. long; branches 2 to 4-angled, mostly 3, 8 to 10 cm. long, 10 mm. broad or less, without aerial roots; flowers borne singly along the whole branch, small; sepals 2; petals 4, creamy white; ovary not sunken in the branch; fruit globose, naked or with an occasional small scale, white, 5 mm. long; seeds numerous, brownish.

Type locality: Navarro, Costa Rica.

Distribution: Costa Rica.

The above description with regard to flowers and fruit has been copied. Our living specimens suggest that it may be different from *Rhipsalis tonduzii*, but whether specifically distinct will require further study to determine.

42. *Rhipsalis boliviana* (Britton) Lauterbach in Buchtien, Contr. Fl. Bolivia **1**: 145. 1910.

Hariota boliviana Britton in Rusby, Mem. Torr. Club **3**: 40. 1893.

Stems somewhat 4-angled and narrowly winged at base, setose at the areoles, the setae 5 to 10, yellowish white, about 2 mm. long; branches 5.5 to 30 cm. long, flattened and thin, 1 to 2 cm. broad, broadly crenate, the crenations 1.5 to 3 cm. long; flowers usually solitary but sometimes 2 or 3 at an areole, about 15 mm. long, one-half to two-thirds as broad, "yellow"; fruit globose, nearly 1 cm. in diameter, truncate at apex.

Type locality: Yungas, Bolivia.

Distribution: Wet forests of Bolivia.

43. *Rhipsalis lorentziana* Grisebach, Abh. Ges. Wiss. Göttingen **24**: 139. 1879.

Epiphytic on forest trees or clambering over rocks, freely rooting along stems; lower part of stem often terete; branches thin, flattened or sometimes 3-angled, usually elongated and narrow, sometimes more or less constricted near middle, 3 cm. broad or less, coarsely serrate, usually cuneate at base; flowers white, about 4 cm. long; ovary oblong, strongly angled, naked except a few scales at the top; fruit globose, purplish, 3 mm. in diameter.

Type locality: Oran, Argentina.

Distribution: Northwestern Argentina and to be expected in southern Bolivia.

Dr. Kurtz gave to Dr. Rose when he was in Córdoba, Argentina, in 1915, a part of the plant collected by Lorentz and Hieronymus in 1893 (No. 454), which proves to be the type.

44. *Rhipsalis ramulosa* (Salm-Dyck) Pfeiffer, Enum. Cact. **130**. 1837.

Cereus ramulosus Salm-Dyck, Hort. Dyck. 340. 1834.

Hariota ramulosa Lemaire, Cact. Gen. Nov. Sp. 75. 1839.*

Stems woody, 3 dm. or more high, erect, terete; branches 7 to 12 cm. long, 1.2 to 2.5 cm. broad, pale green, with distant low crenations 12 to 20 mm. apart, when young often ciliate at areoles but in age naked; flowers solitary at the areoles, small, rotate, greenish white; sepals and petals 6 or 7, ovate-lanceolate, adhering to the base of the ovary, persistent; stamens 12 to 18; style filiform; stigma lobes inconspicuous; fruit glabrous, 5 to 6 mm. in diameter, white and subpellucid with 2 to 3 minute scales; seeds small, black.

Type locality: Not cited.

Distribution: Western Brazil and the adjacent borders of Bolivia and Peru (according to Vaupel).

Collected by R. S. Williams at Isapuri, Bolivia, altitude 5,550 feet, October 1, 1901 (No. 734). We have also referred here H. H. Rusby's No. 749 from trunk of trees near the cataracts of Bopi River, Bolivia, altitude 2,500 feet, September 8, 1921.

We know this plant from herbarium specimens; it is similar to *Rhipsalis lorentziana* but bearing scales on the ovary.

* Lemaire, in 1839 (Cact. Gen. Nov. Sp. 74, 75), combines *Rhipsalis* with *Hariota*, and 8 of the 10 species which he lists had not heretofore been referred to *Hariota*. They are, therefore, to be credited to Lemaire rather than to Otto Kuntze (Rev. Gen. Pl. **1**: 262. 1891), as has been done in the Index Kewensis.



M. E. Eaton del.

A. Hoern. & Co. Baltimore

1. Flowering branch of *Rhipsalis houlettiana*.

2. Flower of same.
3. Fruiting branch of *Rhipsalis warmingiana*.

4. Fruiting branch of same.

Rhipsalis ramulosa has long been a doubtful species. Its origin was unknown at the time of its first publication, but Schumann in 1890 attributed it to Costa Rica, but this was evidently a mistake.

Vaupel has recently published an article (Zeitschrift für Sukkulantenkunde 1: 19. 1923) in which he states that the type was cultivated in the Botanical Garden of Berlin in 1833 and that specimens are now preserved in the herbarium there. He states that these are the same as the plant collected by Ule at Seringal, San Francisco, in the Upper Acre region of Brazil, about 10° south latitude, towards the border of Bolivia and Peru. He would also refer here a plant collected by Tafalla in 1790 at Pozugo in eastern Peru. *Cactus dentatus* Ruiz (Martius, Fl. Bras. 4²: 288. 1890), given as a synonym of *Rhipsalis alata* by Schumann, is based on Tafalla's plant and according to Vaupel should not have been credited to Ruiz.

Epiphyllum ramulosum, *E. ciliare*, and *E. ciliatum* were all given by Pfeiffer (Enum. Cact. 130. 1837) as synonyms of *Rhipsalis ramulosa*.

Figure 229 shows a drawing made from Mr. Williams's specimen.




FIG. 229.—Top of fruiting branch of *Rhipsalis ramulosa*. $\times 0.75$.

45. *Rhipsalis purpusii* Weingart, Monatsschr. Kakteenk. 28: 78. 1918.

Plant epiphytic; stems 8 mm. in diameter, woody, terete, brown; branches weak, elongated, terete below, flattened above, thin, remotely crenate; flowers small, white, solitary.

Type locality: Cerro de Boqueron, Chiapas, Mexico.

Distribution: Known only from the type locality.

This must be related to the Costa Rican plant, *Rhipsalis coriacea*, and perhaps conspecific.

Illustrations: Monatsschr. Kakteenk. 28: 79; Möllers Deutsche Gärt. Zeit. 35: 117.

46. *Rhipsalis coriacea* Polakowsky, Linnaea 41: 562. 1877.

Hariota coriacea Kuntze, Rev. Gen. Pl. 1: 262. 1891.

Rhipsalis angustissima Weber Bull. Mus. Hist. Nat. Paris 8: 465. 1902.

Rhipsalis leiophloea Vaupel, Zeitschrift Sukkulentk. 1: 20. 1923.

Stems 2 to 10 cm. high, woody and terete at base, with many lateral branches; branches often hanging, 1 to 3.5 cm. broad, thin, serrate, the teeth 1.5 to 2.5 cm. apart, bearing the small areoles; young branches purple, terete at first, but finally broad and flattened above; areoles at base of branch and sometimes but rarely on flattened part, bearing 2 to 7 long, hairy bristles; flowers rather narrow, including ovary 12 mm. long, each subtended by a shallow scale; sepals and petals erect below; sepals usually 3, cream-colored, tinged with red; petals greenish white to pinkish, usually 5 to 10, obtuse, 7 to 8 mm. long; stamens numerous, white; style white; stigma-lobes short, white; fruit white, 7 mm. in diameter, bearing several broad, rounded scales; seeds black.

Type locality: Near Cartago, Costa Rica.

Distribution: Widely distributed in Costa Rica.

This species flowers in March.

Rhipsalis coriacea, which originally came from Costa Rica, Schumann referred to *R. alata* of Jamaica, a plant of similar habit but yet very distinct.

Illustration: Bull. Mus. Hist. Nat. Paris **8**: 466, as *Rhipsalis angustissima*.

Of plate xxxii, figure 4 shows a flowering specimen and figure 5 a fruiting specimen from a plant collected by Wm. R. Maxon at Tunialba, Costa Rica, in April 1900, painted at the New York Botanical Garden on April 12, 1912.

47. *Rhipsalis jamaicensis* Britton and Harris, *Torreyia* **9**: 159. 1909.

Pendent from trees, 3 to 10 dm. long, the main axis angular; joints 1 to 4 dm. long, 1 to 2.5 cm. broad, thin, dull green, bluntish at apex, narrowed into a short or elongated stipe at base, the margins low-crenate; flowers yellowish green, about 6 mm. long; perianth-segments about , oblong to oblanceolate, only a little spreading, obtusish; ovary oblong, bearing a few small scales stamens 20 to 30; style longer than stamens; stigma-lobes 3; fruit globose, 6 to 8 mm. in diameter, white, the scales 3 mm. broad.

Type locality: Troy, Cockpit Country, Jamaica.

Distribution: Forests of Jamaica.

Illustration: *Torreyia* **9**: 158. f. 3.

Plate xxii, figure 4, shows a plant with flowers and young fruit from Jamaica.

48. *Rhipsalis platycarpa* (Zuccarini) Pfeiffer, *Enum. Cact.* **131**. 1837.

Epiphyllum Platycarpum Zuccarini, *Cat. Cact. Monac.* 1836.

Cereus Platycarpus Zuccarini, *Abh. Bayer. Akad. Wiss. München* **2**: 736. 1837.

Hariota Platycarpa Kuntze, *Rev. Gen. Pl.* **1**: 263. 1891.

Branches broad and flat, 1 to 2 dm. long, 3 cm. broad or more, dull green becoming red when grown in sunlight, with broad deep crenations; flowers borne toward apex of branch, 1 to 3 from an areole, 16 to 18 mm. long, greenish yellow or dull white; petals 8 mm. long, ovate; stamens white; stigma-lobes , white; fruit (doubtless immature) naked, green, somewhat compressed, angled, truncate.

Type locality: Brazil.

Distribution: Organ Mountains, Brazil.

We have obtained plants of this species from Mr. Lamb at Manchester in 1904 and Dr. Rose found it wild in the Organ Mountains of Brazil in 1915 (No. 21159). It grows well in cultivation but it has never flowered with us.

Illustrations: Pfeiffer and Otto, *Abbild. Beschr. Cact.* **1** pl. 17, f. 2; *Blühende Kakteen* **2**: pl. 90.

Figure 230 shows a branch of the plant obtained by Dr. Rose in the Organ Mountains.





FIG. 230.—*Rhipsalis platycarpa*. $\times 0.4$

49. *Rhipsalis russellii* sp. nov.

Hanging in great clusters from the horizontal branches of trees; branches strongly flattened, 15 cm. long, , to 6 cm. broad, cuneate at base, strongly crenate, dark green or purplish along margins; flowers often 9 at an areole, minute; sepals few, obtuse, reddish at tips; petals usually 5, cream-colored, obtuse, 2 mm. long; fruit usually 1 at an areole, small, globose, to 6 mm. in diameter, purple.

Collected by J. N. Rose and Paul G. Russell near Toca da Onca, Bahia, Brazil, June 27 to 29, 1915 (No. 20106). This species suggests *Rhipsalis elliptica*, but has very different flowers and fruit.



M. E. Eaton del.

1. Fruiting branch of *Rhipsalis oblonga*.
2. Fruiting branch of *Rhipsalis elliptica*.
3. Flowering branch of *Rhipsalis crispata*.

A. Hoen & Co. Baltimore

Of plate xxxvii, figures 1 to 4 are from the type specimen which has repeatedly flowered and fruited in the New York Botanical Garden; figure 1 shows the tip of a flowering branch; figure 2 shows a cluster of six flowers; figure 3 shows a flower enlarged four diameters; figure 4 shows a fruiting branch.

50. *Rhipsalis elliptica* Lindberg in Martius, Fl. Bras. 4²: 293. 1890.

Rhipsalis chloroptera Weber, Dict. Hort. Bois 1045. 1898.

Rhipsalis elliptica helicoidea Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 44. 1918.

Plants growing in clumps, at first ascending, often hanging from trees; joints flat and broad, oblong to elliptic, 3 to 20 cm. long, 2 to 7 cm. broad, the margin faintly to strongly crenate; flowers generally 1, sometimes 2 or 3 at an areole, 12 mm. broad; petals usually 5, yellowish, widely spreading, oblong, obtuse; filaments numerous, nearly erect, white; style white; stigma-lobes white, 5; ovary not sunken in the branch; fruit reddish, a little longer than broad, 6 to 7 mm. long.

Type locality: Near Sorocaba, south of Santos, São Paulo, Brazil.

Distribution: States of São Paulo and Santa Catharina, Brazil.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 2: pl. 16, as *Rhipsalis elliptica helicoidea*; Blühende Kakteen 2: pl. 104, as *Rhipsalis chloroptera*; Arch. Jard. Bot. Rio de Janeiro 1: pl. 15.

Plate xxxv, figure 2, is from a plant collected by Dr. Rose at Jabaquara, near Rio de Janeiro, Brazil, in 1915, which flowered in the New York Botanical Garden in 1916.

51. *Rhipsalis pachyptera* Pfeiffer, Enum. Cact. 132. 1837.

Cactus alatus Willdenow, Enum. Pl. Suppl. 35. 1813. Not Swartz, 1788.

Epiphyllum alatum Haworth, Suppl. Pl. Succ. 84. 1819.

Cactus triquetra Vellozo, Fl. Flum. 206. 1825. Not Willdenow, 1813. Not Haworth, 1803.

Cereus alatus Link and Otto, Icon. Pl. Rar. 77. 1830.

Lepismium fluminense Miquel, Bull. Neerl. 48. 1838.

Rhipsalis robusta Lemaire, Rev. Hort. IV. 9: 502. 1860.

Rhipsalis pachyptera purpurea Corderoy, Gard. Chron. III. 2: 468. 1887.

Hariota triquetra Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Hariota pachyptera Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Hariota robusta Kuntze, Rev. Gen. Pl. 1: 263. 1891.

Rhipsalis crassa Schumann, Keys 54. 1903.

Stems much jointed, pendent; joints often 3 to 6 dm. long, 5 to 7 cm. broad, thickish, stiff, sometimes nearly orbicular, often purple, deeply crenate; flowers numerous, but solitary, rarely 2 to 4 from the areole, large; petals widely spreading, yellowish; stamens numerous; stigma-lobes 4 or 5, slender; fruit globose, white.

Type locality: Originally given as the West Indies,* but this is doubtless a mistake.

Distribution: States of Rio de Janeiro, Minas-Geraes, Santa Catherina, and São Paulo, Brazil.

The species grows in the high mountains on trunks of trees, altitude 1,000 meters, down to nearly sea-level.

A variety, *crassior* Salm-Dyck (Pfeiffer, Enum. Cact. 132. 1837), with thick green orbicular joints, has been described.

Steudel's name of *Rhipsalis alata* (Nom. ed. 2. 1: 333. 1840), given as a synonym of *Cereus alatus* De Candolle, is referred here by Schumann, but probably relates to *Pseudorhipsalis*.

This species was for a long time confused with *Rhipsalis alata*, a very distinct species from Jamaica, now referred by us to the genus *Pseudorhipsalis*.

Illustrations: Curtis's Bot. Mag. 55: pl. 2820,* as *Cactus alatus*; Vellozo, Fl. Flum. 5: pl. 25, as *Cactus triquetra*; pl. 33, as to flower only; Link and Otto, Icon. Pl. Rar. pl. 39, as *Cereus alatus*; Monatsschr. Kakteenk. 6: 55; 7: 151, in part, as *Rhipsalis robusta*;

* The plant, however, which Hooker described and figured (Curtis's Bot. Mag. 55: pl. 2820) as *Cactus alatus* and which Pfeiffer cited in his original description, came from the Organ Mountains near Rio de Janeiro, Brazil.

Blühende Kakteen 1: pl. 34; Martius, Fl. Bras. 4²: pl. 57; Paxton's Fl. Gard. 1: pl. 99; Arch. Jard. Bot. Rio de Janeiro 1: pl. 14; Möllers Deutsche Gärt. Zeit. 25: 77. f. 11, No. 15; Gartenwelt 13: 117; 16: 633, 635; Karsten and Schenck, Vegetationsbilder 1: pl. 5, f. d.

Plate xxxvii, figure 6, shows a fruiting branch from the plant obtained by Dr. Rose in Brazil in 1915 (No. 20346); plate xxxvi, figure 1, shows a flowering branch from a plant obtained from M. Simon, of Paris, in 1901.

52. *Rhipsalis rhombea* (Salm-Dyck) Pfeiffer, Enum. Cact. 130. 1837.

Cereus rhombeus Salm-Dyck, Hort. Dyck. 341. 1834.

Hariota rhombea Lemaire, Cact. Gen. Nov. Sp. 75. 1839.

Stems terete or angled; branches usually flat and thin, but sometimes 3-angled; joints oblong, 1 to 3 cm. broad, cuneate at base, strongly crenate, dark green or purple; flowers usually solitary at areoles but sometimes in 2's, small, cream-colored, with a red spot at base of stamens; sepals reddish; petals obtuse; fruit dark red.




FIG. 231.—*Rhipsalis crispimarginata*.

Type locality: Not cited.

Distribution: Brazil, but range unknown.

Cereus crispatus crenulatus, *Epiphyllum crenulatum*, and *E. rhombeum* were referred by Pfeiffer (Enum. Cact. 130. 1837) as synonyms of this species.

Here perhaps also belongs *Cereus crispatus latior* (Salm-Dyck, Hort. Dyck. 66. 1834), which is without description.

Illustrations: Gartenwelt 16: 635; Karsten and Schenck, Vegetationsbilder 1: pl. 6, f. e; Möllers Deutsche Gärt. Zeit. 25: 477. f. 11, No. 13; Wildeman, Icon. Select. 2: pl. 67; Arch. Jard. Bot. Rio de Janeiro 1: pl. 16.

Plate xxxvi, figure 2, shows a flowering plant received from the Royal Botanic Garden at Kew in 1902 which flowered in the New York Botanical Garden in January 1912.

*According to the Index Kewensis this is *Cactus speciosus* Hooker, said to be equal to *Rhipsalis swartziana*.



M. E. Eaton del.

1. Flowering branch of *Rhipsalis pachyptera*.
2. Flowering branch of *Rhipsalis rhombea*.

A. Hoen & Co. Baltimore

53. *Rhipsalis crispimarginata* Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 37. 1918.

Plants pendulous, the main stem terete below, often 3-winged above; terminal branches in clusters, oblong, flat, obtuse, narrowed at base, shining green or tinged with purple, 4 to 6 cm. long; flowers usually solitary but sometimes 2 or 3 at an areole, white, drying pale yellow; sepals ovate-obtuse, reflexed; petals white, widely spreading, numerous; stigma-lobes white; fruit globose, white.

Type locality: Ilha Grande, near the city of Rio de Janeiro.

Distribution: State of Rio de Janeiro, Brazil.

Illustration: Arch. Jard. Bot. Rio de Janeiro 2: pl. 9.

Plate xxxvii, figure 5, shows a fruiting branch of the type collection obtained by Dr. Rose on Ilha Grande near Rio de Janeiro in 1915 (No. 20401). Figure 231 is from a photograph of Miss Eaton's painting of a plant given to Dr. Shafer by Dr. Löfgren in 1917 at Rio de Janeiro which flowered in the New York Botanical Garden in May 1922.

54. *Rhipsalis crispata* (Haworth) Pfeiffer, Enum. Cact. 130. 1837.

Epiphyllum crispatum Haworth, Phil. Mag. 7: 111. 1830.

Rhipsalis crispata latior Salm-Dyck in Pfeiffer, Enum. Cact. 130. 1837.

Hariota crispata Lemaire, Cact. Gen. Nov. Sp. 75. 1839.

Rhipsalis rhombea crispata Schumann, Gesamtb. Kakteen 638. 1898.




FIG. 232.—*Rhipsalis crispata*.

Branches divided into short flat joints 6 to 10 cm. long, broad both at base and apex, green, more or less crenate; flowers solitary or sometimes 2 to 4 at an areole, 10 to 12 mm. broad, cream-colored to light yellow; filaments numerous; fruit white, 7 mm. in diameter.

Type locality: Brazil.

Distribution: Brazil, but range unknown.

The synonyms *R. crispa* and its var. *major* (Förster, Handb. Cact. 450. 1846) and *R. crispa latior* Salm-Dyck (Walpers, Repert. Bot. 2: 279. 1843); *Hariota crispata latior* Lemaire (Cact. Gen. Nov. Sp. 75. 1839) doubtless belong here. Schumann, in Nachtrag, page 144, refers here *Rhipsalis swartziana* Pfeiffer.

Pfeiffer publishes as a synonym of the above *Cereus crispatus* (Enum. Cact. 130. 1837), as does also Förster. Schumann (Gesamtb. Kakteen 638. 1898) makes this species a variety of *Rhipsalis rhombea*, but later (Gesamtb. Kakteen Nachtr. 145. 1903) recognizes two species.

Cactus torquatus (Walpers, Repert. Bot. 2: 342. 1843), referred to *Rhipsalis rhombea* by Walpers, was only a garden name.

Illustrations: Arch. Jard. Bot. Rio de Janeiro 1: pl. 16, as *R. rhombea*; Gartenwelt 13: 117; Garten-Zeitung 1: 459. f. 109; Rev. Hort. 85: f. 152, in part.

Plate xxxv, figure 3, shows a flowering plant received from A. Berger in 1908. Figure 232 is from a photograph of Miss Eaton's painting of the plant obtained by Dr. Rose in Brazil in 1915 (No. 20708) which flowered and fruited in the New York Botanical Garden in 1922.

55. *Rhipsalis oblonga* Löfgren, Arch. Jard. Bot. Rio de Janeiro 2: 36. 1918.

In cultivation bushy; main branches terete below, more or less flattened above; ultimate branches narrowly oblong, 5 to 15 cm. long, 1 to 2 cm. broad, shining green even in sunlight; flowers borne along the sides of the branches, solitary at the areoles; fruit globular to short-oblong, 3 to 4 mm. long, nearly white, naked, crowned by the withered perianth.

Type locality: On Ilha Grande, Brazil.

Distribution: Known only from the type locality.

Illustration: Arch. Jard. Bot. Rio de Janeiro 2: pl. 8, as *Rhipsalis oblonga*.

Plate xxxv, figure 1, shows the plant grown by Dr. Löfgren at Rio de Janeiro and given to Dr. Shafer in 1917, which flowered and fruited in the New York Botanical Garden in May 1922.

56. *Rhipsalis cuneata* sp. nov.

Epiphytic on trees; joints oblong to spatulate, 8 to 12 cm. long, thin, obtuse, cuneate at base, strongly crenate, naked at the areoles or with a bristle or two; flowers so far as known solitary; fruit globose, 4 mm. in diameter, naked.




FIG. 233.—*Rhipsalis cuneata*.

Collected by R. S. Williams above San Juan, Bolivia, altitude 5,500 feet, April 2, 1902 (No. 2458). This species is known to us only from herbarium specimens.

Figure 233 is from a photograph of the specimen in the U. S. National Herbarium.

57. *Rhipsalis roseana* Berger, Zeitschrift für Sukkulantenkunde 1: 22. 1923.

Lower joints flat, 15 to 20 mm. broad, distinctly alternately notched; areoles small, with a little tuft of white wool and a single short brown hair, 15 to 20 mm. apart, the upper ones more closely set; upper joints narrower and more linear or linear-lanceolate, 10 to 15 mm. broad and 6 to 12 cm. long or more, equally notched, smooth, bright green; some of the uppermost joints often narrower, 8 to 10 mm. broad and only shallowly notched, others triangular with prominent notched angles and excavated sides, others 1 cm. wide, with 3 or 4 prominent wing-like distinctly but remotely notched ribs and areoles about 4 cm. apart; flowers small, whitish yellow.

This species was described from cultivated plants of unknown origin. We believe that it may be from Colombia and we would refer here the following specimens: Wilson Popenoe's No. 518 from near San Miguel, Perdona, Tolima, 1921, and Ellsworth P. Killip's No. 8203 from mountains west of Popayán, 1922.

Mr. Berger writes: "This new *Rhipsalis* is decidedly distinct from *R. wercklei*; its branches are shorter, broader, more deeply notched and of a firmer nature. Its growth too is far less quick and it does not form so promptly long and pendent shoots as *R. wercklei*."



M. E. Eaton del.

A. Hoen & Co. Baltimore

1. Flowering branch of *Rhipsalis russelli*.
2. Cluster of flowers of same.
3. Flower of same.

4. Fruiting branch of same.
5. Fruiting branch of *Rhipsalis crispimarginata*.
6. Fruiting branch of *Rhipsalis pachyptera*.

UNPUBLISHED OR INCOMPLETELY DESCRIBED SPECIES.

RHIPSALIS CHRYSANTHA Löfgren, Arch. Jard. Bat. Rio de Janeiro 1: 99. 1915.

We know this species only from description. Löfgren places it in his subgenus *Lepismium* near *Rhipsalis dissimilis*, but his descriptions suggest *R. rosea* (our *Rhipsalidopsis rosea*). Both names are based on Dr. P. Dusen's collections from Paramá, Brazil. It seems near *R. puniceo-discus*.

RHIPSALIS FRONDOSA Wercklé, Subregion Fitogeografica Costa Ricense 42. 1909.

The above name is given without description.

Weingart (Monatsschr. Kakteenk. 20: 185. 1910) refers to this plant as a new species represented in a sending from Costa Rica by Wercklé. Nothing further is known about it.

RHIPSALIS RIEDELIANA Regel, Ind. Sem. Hort. Petrop. 1860: 49. 1860.

Hariota riedeliana Kuntze, Rev. Gen. Pl. 1: 263. 1891.

This plant was sent from Brazil by Riedel, but we do not know it. Schumann did not know it.

Rhipsalis bucheni Béhagnon (Rev. Hort. 85: 436. f. 152. 1913) we know only from the illustration of a poor potted plant and an incomplete description.

Rhipsalis carnosa and *R. lagenaria* are names mentioned by Vöchting (Jahrb. Wiss. Bot. Leipzig 9: 368, 372. 1873).

Rhipsalis erythrolepis Bénagnon (Rev. Hort. 85: f. 152, part) is known only from a potted plant of some species with broad, flat joints.

Rhipsalis filiformis seems to be only a garden name (Monatsschr. Kakteenk. 6: 47. 1896). It may be the same as *R. cribrata filiformis* Engelhardt (Möllers Deutsche Gärt. Zeit. 18: 585. 1903).

Rhipsalis itatiaiae Weber appears in Robert Lamb's Collection of Cacti, page 72, 1908, without description. In 1914 Mr. Lamb sent Dr. Rose a specimen under this name, but it has not bloomed. A part of this plant from Mount Itatiaia, Brazil, is now growing in the New York Botanical Garden (Rose, No. 888).

Rhipsalis macabensis Glaziou (Bull. Soc. Bot. France Mem. III. 326. 1909) is only a name. According to Glaziou he collected it on rocks and trees at Alto Macahé, Rio de Janeiro (No. 18262).

Rhipsalis microcarpa Steudel, is a name found only in Schumann's Index (Gesamtb. Kakteen 832. 1898).

Rhipsalis miquelii Lemaire (Cactées 80. 1868) is not described but it is grouped with *R. pachyptera*, *R. rhombea*, and other flat-jointed species. Lemaire also lists *R. turpinii* on the same page, associating it with *R. micrantha* and *R. trigona*.


Rhipsalis oligosperma Lindberg (Monatsschr. Kakteenk. 7: 21. 1897) is a name only.

Rhipsalis spathulata Otto (Sweet, Hort. Brit. ed. 3. 1839) Schumann thought might be a mistake for *Pereskia spathulata*. Kuntze takes it up, however, as *Hariota spathulata* (Rev. Gen. Pl. 1: 263. 1891).

The name *Rhipsalis taglionis* occurs in the Index Kewensis Supplement 1, by error for *R. saglionsis*.

Rhipsalis wettsteinii Schumann (Monatsschr. Kakteenk. 17: 48. 1907) is a name only.

The illustration (fig. 233a) given on next page is a reproduction of Plumier's plate 92 of Burmann Plantarum Americanum published in 1755 and now referred to *Neoabbottia paniculata*, discussed on page 280 of this volume.



CACTUS brachiatus & articulatus .

APPENDIX.

During the progress of our investigations much information has been received from numerous sources which could not be included in publication at the logical places. Some of this was taken up in the appendix to the first volume (Cactaceae **1**: 2 16-225) and some in the appendix to the second (Cactaceae **2**: 223-226); what remains is included in this appendix to the whole work.

Dr. David Griffiths, who studied the species of *Opuntia*, especially with relation to their economic possibilities, and grew many of them at experimental stations of the United States Department of Agriculture at Brownsville, Texas, and at Chico, California, has published and described many species as new. We have included these in our studies of the genus and have grouped them with the species known to us as accurately as has been possible from his published descriptions and illustrations and after examination of as many of his type specimens as we have been permitted to see; however, conditions were such that we have not been able to study a number of them. They have not been arranged for ready reference by students.

The preface to Volume I gives a list of volunteers who have made valuable contributions of specimens and data to this investigation. Many of these have continued to aid us.

Dr. Britton, in continuing his West Indian studies, investigated the cacti of Grenada and of Trinidad in 1920 and 1921 and published an account of the Trinidad species.

Dr. John K. Small has continued his investigation of the southeastern United States and of Florida in particular, in cooperation with Mr. Charles Deering, and has greatly increased our knowledge of the cacti existing there, including the discovery of many undescribed species of *Opuntia*.

Dr. Francis W. Pennell, Curator of Botany at the Academy of Natural Sciences of Philadelphia, and Mr. E. P. Killip, of the United States National Museum, made extensive botanical collections in Colombia in 1922, including some specimens of cacti, which we have studied.

Dr. Henry H. Rusby led the Mulford Biological Exploring Expedition to Bolivia in 1921-1922 and with the assistance of Dr. O. E. White obtained for us specimens of several little-known cacti.

Dr. Philip A. Munz has sent us cacti from the deserts of southern California.

Mr. C. Z. Nelson obtained cacti from southern Mexico, including a beautiful new species of *Selenicereus*.

Mr. Francis J. Dyer, while connected with the Consular Service in Honduras and at Nogales, Mexico, sent us many specimens from those stations.

Professor Harvey M. Hall, while making extensive explorations in the western United States in connection with his own work, has forwarded interesting cactus plants.

Dr. and Mrs. Charles D. Walcott have sent specimens from Alberta, Canada, some of them coming from near the most northern range of the family.

Dr. W. L. Abbott and Mr. E. C. Leonard made extensive collections in Haiti in 1920 and obtained a number of rare and little-known plants, including one which had been collected by Charles Plumier about 1698 and which proved to be a new genus; this was named by us for Dr. Abbott. More recently Dr. Abbott has sent us specimens from Santo Domingo.

Dr. George F. Gaumer, the veteran collector in Yucatan, has sent very important collections from his region, including a number of new species.

Mr. Ivan M. Johnston, who accompanied the scientific expedition sent out by the California Academy of Sciences in 1921 to explore the islands of the Gulf of California, 249

collected many cacti, especially species of *Neomammillaria*, duplicating many of the important discoveries made by Dr. Rose on the same islands in 1911. He has also sent us cacti from Colorado.

Professor Fortunato L. Herrera has sent some very interesting plants from eastern Peru, especially from about his home at Cuzco.

Mr. Robert Runyon has collected extensively in southern Texas and northern Mexico and has supplemented his specimens with some very beautiful photographs.

Dr. L. H. Bailey and his daughter, Miss Ethel Zoe Bailey, obtained valuable cacti from Venezuela, especially from the region about Ciudad Bolívar, on the Orinoco, in 1921.

Mr. W. B. Alexander was sent to Argentina by the Australian Government in 1920 and 1921 in search of enemies of the weed prickly pears and there made many important observations, especially on the genus *Opuntia*. He sent us two undescribed species.

Dr. B. P. Reko, a very diligent collector, has sent many cacti from Mexico, especially from Oaxaca, including several new to science.

Señor Octavio Solís, in charge of the cactus garden belonging to the Mexican Government in the City of Mexico, has sent many living plants from his country, especially of the genus *Neomammillaria*. To him we have dedicated the genus *Solisia*.

Señor J. G. Ortega has collected extensively on the west coast of Mexico, especially in the state of Sinaloa, and for him we have named *Neomammillaria ortegae*.

Mr. J. Francis Macbride and Mr. William Featherstone, who were in charge of the botanical expedition of the Field Museum to Peru in 1922 and 1923, made large and valuable collections of cacti in central and eastern Peru.

Mr. E. C. Rost has collected and photographed many interesting cacti for us in southern California and Lower California.

Dr. W. S. W. Kew explored extensively in Lower California in 1921 and sent not only many specimens but numerous habit photographs.

Mr. James H. Ferriss, while making various excursions through the western United States, has sent in many specimens. Among his interesting discoveries was the finding of *Neomammillaria pottsii* in southern Texas.

Mrs. S. L. Pattison, an enthusiastic collector in western Texas, has sent many valuable specimens, including new species collected by herself or for her by local collectors.

Mrs. Ruth C. Ross spent considerable time in eastern Arizona in 1921 and collected cacti along the route traversed by Emory in 1847, re-collecting certain species which he had discovered at that time.

Mr. Harry Johnson was located for about a year in Guatemala, during which time he sent a number of very interesting cacti, especially species of *Epiphyllum*. Some of these were accompanied by full notes and drawings.

Señor P. Campos-Porto has sent a number of interesting specimens from Brazil belonging to the genus *Rhipsalis*.

The following persons have contributed valuable specimens, usually from about their homes or while engaged in other work: G. W. Goldsmith, Colorado; B. C. Tharp, Texas; Charles O. Chambers, Oklahoma; James S. Holmes, Washington, D. C.; Joseph A. Holmes, Wyoming; William Hertrich, California; William Tell, Texas; Albert Ruth, Texas; D. C. Parman, Texas; Karl Reiche, Mexico; Gerold Stahel, Surinam; Rev. Louis Mille, Ecuador; H. M. Pilkington, Haiti; Percy L. Ports, Washington, D. C.; W. E. Broadway, Trinidad; A. F. Moeller, Mexico; W. E. Meyer, Bolivia; Stephen E. Aguirre, Mexico; Mrs. Elsie McElroy Slater, Texas; Paul C. Standley, Central America; R. D. Camp, Brownsville, Texas; George L. Fisher, Texas; A. V. Frič, Mexico; and Dana Lee, Wyoming.

As treated in this monograph the Cactus family is composed of 3 tribes. The first and second tribes are taken as units, but the third is composed of 8 subtribes. The number of genera recognized is 124 and the number of species is 1,235.

CORRECTIONS AND ADDITIONS TO VOLUME I.

On page 11, vol. 1, under *Pereskia pereskia*, add to illustrations: Garten-Zeitung 4: 182. f. 42, No. 5; Gard. Chron. III. 20: f. 108; Stand. Cycl. Hort. Bailey 2: f. 714, as *Pereskia aculeata*; Loudon, Encycl. Pl. ed. 3. 413, as *Cactus pereskia*; Möllers Deutsche Gärt. Zeit. 23: 256. 1. 15, as *Pereskia godseffiana*.

Also insert: *Pereskia longispina rubescens* Pfeiffer and *P. longispina rotundifolia* Salm-Dyck were given by Walpers (Repert. Bot. 2: 283. 1843) as synonyms of *P. aculeata*, but they were not described.

On page 12, vol. 1, under *Pereskia autumnalis*, add to distribution: Common in Salvador where it is much planted for hedges.

Also add to illustrations: Monatsschr. Kakteenk. 25: 35, as *Pereskopsis autumnalis*; Engler and Drude, Veg. Erde 13: f. 10, as *Pereskia guatemalensis*.

On page 14, vol. 1, under *Pereskia sacharosa*, add the synonym: *Pereskia amapola argentina* Weber in Weingart, Monatsschr. Kakteenk. 14: 87. 1894.

On page 17, vol. 1, under *Pereskia guamacho*, insert: Illustration: Carnegie Inst. Wash. 269: pl. 11, f. 92, 93.

On page 20, vol. 1, under *Pereskia grandifolia*, add to illustrations: Rümpler, Sukkulanten f. 128; Engler and Prantl, Pflanzenfam. 3^{6a}: f. 57, J; Blühende Kakteen 3: pl. 137; Watson, Cact. Cult. f. 6, in part; 222. f. 87; ed. 3. f. 63; Karsten, Deutsche Fl. ed. 2. 2: 456. f. 605, No. 9; Loudon, Encycl. Pl. ed. 3. 1202. f. 17371; Van Géel, Sert. Bot. 4: pl. 1, as *Pereskia bleo*; Dict. Gard. Nicholson 3: 75. f. 81; Monatsschr. Kakteenk. 15: 81.

Also add synonym: *Cactus grandiflorus* Link, Enum. Hort. Berol. 2: 25. 1822.

On page 21, vol. 1, *Pereskia zinniaeflora*, add to illustrations: Watson, Cact. Cult. ed. 1 and 2. 223. f. 88; ed. 3. f. 64; Dict. Gard. Nicholson 4: 586. f. 55.

On page 21, vol. 1, under *Pereskia horrida*, substitute for this name:

Pereskia humboldtii nom. nov.

Cactus horridus Humboldt, Bonpland, and Kunth, Nov. Gen. et Sp. 6: 70. 1823. Not Salisbury, 1796.
Pereskia horrida De Candolle, Prodr. 3: 475. 1828.

On page 24, vol. 1, at end of *Pereskia*, add: *Pereskia recurvifolia* and *P. galeottiana* are two names marked with an asterisk by Lemaire (Cactées 95. 1868), indicating that they are new. So far as we know they were never described.

On page 24, vol. 1, at end of *Pereskia* insert:

Pereskia pflanzii Vaupel, Zeitschrift Sukkulantenk. 1: 56. 1923.

Tree about 15 meters high, with verticillate branches, not very spiny; leaves thick, ovoid, narrowed at base, 4 cm. long by 2 cm. broad; flowers solitary at apex of leafy branches; corolla 3 cm. long, rose-colored.

Type locality: Vicinity of Laguna Santa Isabel, Bolivia.

Distribution: Bolivia, but known only from type locality.

Pereskia verticillata Vaupel, Zeitschrift Sukkulantenk. 1: 55. 1923.

Erect shrub, 2 meters high, very spiny, with verticillate branches; leaves thick, lanceolate, 5 cm. long by 1.5 cm. broad; flowers borne at apex of leafy branches; corolla 1.5 cm. long, rose-colored.

Type locality: Vicinity of Laguna Santa Isabel, Bolivia.

Distribution: Bolivia, but known only from type locality.

On page 27, vol. 1, under *Pereskopsis chapistla*, add to illustration: Smiths. Misc. Coll. 50: pl. 43.

On page 28, vol. 1, under *Pereskopsis porteri*, add the synonym: *Opuntia rotundifolia* Brandegees, Zoe 2: 21. 1891. Not *Pereskia rotundifolia* De Candolle, 1828.

On page 29, vol. 1, under *Pereskopsis spathulata*, insert: Illustration: Möllers Deutsche Gärt. Zeit. 25: 488. f. 22, No. 1, as *Pereskia spathulata*.

On page 29, vol. 1, under *Pereskiaopsis pititache*, add to illustrations: Deutsche Gärt. Zeit. 8: 33, as *Pereskia calandriniaefolia*.

On page 30, vol. 1, insert the following:

11. *Pereskiaopsis scandens* sp. nov.

Slender, climbing or clambering over walls, up to 10 meters long; branches terete, grayish, smooth; areoles circular, white-woolly when young, gray in age, with a short spine (mm. long) and a bunch of brown glochids in the upper edge; leaves ovate, 1.5 to 2 cm. long, glabrous, acute; flowers yellow, from the areoles on old branches, appearing in June; fruit maturing slowly (perhaps requiring 2 to 3 years to ripen), very narrow, 5 to 7 cm. long, somewhat tubercled, with a deep umbilicus; seeds few.

Living specimens of *P. scandens* were sent by Dr. George F. Gaumer from Izamal, Yucatan, Mexico, in July 1921 (type). It was also collected by A. Schott at Mérida in 1865 (No. 409).

Withdraw the name *Pereskia zehntneri* from page 14, vol. 1, and substitute the following at the end of *Pereskiaopsis* on page 30:

1a. *QUIABENTIA* gen. nov.

A low, leafy, much branched shrub with numerous horizontal branches, usually in whorls; leaves fleshy but flattened, stiff, borne at right angles to the branches; areoles large, white-felted, often with numerous spines, these acicular and white, the upper part of areole bearing glochids; flowers terminal, very large, bright red; ovary leafy, very narrow; stamens numerous, a little shorter than the style, much shorter than the petals; style short and stiff; stigma-lobes very short, obtuse; seeds white, a little flattened, covered with a hard bony aril as in *Opuntia*.

A monotypic genus, native of the semiarid region of Bahia, Brazil. The generic name is from quiabento, the native name of the plant.

1. *Quiabentia zehntneri* Britton and Rose.

Pereskia zehntneri Britton and Rose, Cactaceae 1: 14. 1919.

Flowers at ends of branches, large, 7 to 8 cm. broad, 3 to 4 cm. long, bright red, appearing in November; petals broad, retuse; ovary borne in the upper end of the branch, very narrow, 3 to 4 cm. long, bearing the usual leaves, areoles, and spines of the branches; fruit oblong to clavate, 6 to 7 cm. long, 1.5 cm. in diameter at the top, slightly angled by the low elongated tubercles running downward from the small scattered areoles, and finally without leaves, spines, or bristles, sterile below, with thick fleshy walls and with a small narrow seed-cavity; umbilicus broad, slightly depressed; seeds thick with flattened sides rounded on the back, 5 mm. in diameter.

In its large, red, rotate flowers this plant at once suggests a *Pereskia*. Its red flowers are so similar to those of *P. bahiensis* of the same region that at first we considered the two species congeneric. Now that we have studied the fruit and seed it is evident that *P. zehntneri* belongs to a very different genus. Then, too, the old areoles develop deciduous spines or bristles which are doubtless glochids; these occur on the upper part of the areoles but do not form the definite brush of the *Opuntiae*. These glochids would exclude it from the *Pereskiae*. It must therefore be referred to the *Opuntieae* and next to *Pereskiaopsis*. In its broad, thick leaves it resembles that genus, but its flowers are terminal, very large, and rotate; its fruit is much elongated and the seeds are glabrous.

We are indebted to Dr. Leo Zehntner, a very keen observer, for many fine specimens and much information regarding it. He has found it only on a small calcareous mountain near the city of Born Jesus da Lapa, Brazil, but it has been transplanted to the Horto Florestal of Joazeiro where it is well established and where it flowered three years after being replanted. In 1915 Dr. Rose brought living specimens to the New York Botanical Garden from this stock (No. 19722).

On page 32, vol. 1, under *Pterocactus tuberosus*, add the synonym: *Opuntia tuberosa albispina* Salm-Dyck in Förster, Handb. Cact. ed. 2. 911. 1885.

Also add to illustrations: Haage and Schmidt, Cat. Gen. 230. 1908; De Laet, Cat. Gén. f. 74, as *Pterocactus kuntzei*.

On page 34, vol. 1, under *Nopalea cochenillifera*, add the synonyms: *Cactus nopal* Thierry, Dict. Sci. Nat. 6: 103. 1817; *Cactus splendidus* Thierry, Dict. Sci. Nat. 6: 103. 1817; *Cactus campechianus* Thierry, Dict. Sci. Nat. 6: 103. 1817; *Nopalea coccifera* Lemaire, Cactées 89. 1868.

Also add to illustrations: Loudon, Encycl. Pl. ed. 1 and 3. 412. f. 6888, as *Cactus cochenillifer*; Contr. U. S. Nat. Herb. 8: pl. 48, as spineless opuntia; Knorr, Thesaurus pl. 0, 1.

On page 37, vol. 1, under *Nopalea auberi*, insert:

Opuntia auberi was described as from Cuba, but as no *Nopalea* is known from Cuba we have been unable to account for this reference. The following incidents may explain it:

L. Pfeiffer described the plant in 1840 just after his return from Cuba, where he had gone with Otto in 1838. At Havana they visited the Botanical Garden, then in charge of Pedro Auber, for whom this plant was doubtless named. It is also stated that, although Pfeiffer made this trip especially to gather cacti, he saw only one species, *Opuntia horrida*. The probabilities, therefore, are that this plant was obtained from the Botanical Garden at Havana, perhaps with a statement from Auber that it was Cuban.

On page 37, vol. 1, under *Nopalea dejecta*, add the synonym: *Nopalea angustifrons** Lindberg, Act. Soc. Sc. Fenn. 10: 123. 1871.

Add to illustrations: Act. Soc. Sc. Fenn. 10: pl. 2, as *Nopalea angustifrons*.

On page 41, vol. 1, under *Maihuenia poeppigii*, add to illustrations: Gartenflora 30: 412, as *Pereskia poeppigii*.

On page 42, vol. 1, under *Maihuenia brachydelphys*, insert the synonym: *Opuntia brachydelphis* Schumann in Just, Bot. Jahresb. 26: 343. 1898.

Insert: *Mammillaria brachydelphis* is a clerical error for *Opuntia brachydelphis*.

On page 42, vol. 1, under *Opuntia*, add the synonym: *Cactus* Lemaire,† Cactées 86. 1868. Not Linnaeus, 1753.

On page 46, vol. 1, under *Opuntia ramosissima*, insert: *Opuntia tessellata denudata*, according to C. R. Orcutt, is only a form—spiny joints frequently occurring on the same plant with the spineless form; it is common in the Mojave Desert, California. It was mentioned by Alverson (Cact. Cat. 6) while *O. ramosissima denudata* is listed by Weinberg (Cacti 22). *O. ramosissima cristata* is mentioned by Schelle (Handb. Kakteenk. 41. 1907).

Also add to illustrations: Cact. Journ. 1: pl. for February; Monatsschr. Kakteenk. 8: 71, as *Opuntia tessellata cristata*; Stand. Cycl. Hort. Bailey 4: f. 2596, 2610.

On page 47, vol. 1, under *Opuntia leptocaulis*, add the synonym: *Opuntia californica* Engelm. in Emory, Mil. Reconn. 158. 1848.

Also insert: *Opuntia stipata* (Schumann, Index Gesamtb. Kakteen 830. 1898) refers to *O. leptocaulis stipata*.

Also add to illustrations: Emory, Mil. Reconn. 158. No. 11, as *Opuntia californica*; Gartenwelt 11: 75, as *O. vaginata*; Carnegie Inst. Wash. 269: pl. 10, f. 89; pl. 11, f. 96; Stand. Cycl. Hort. Bailey 2: f. 717; Schelle, Handb. Kakteenk. 41. f. 2; Möllers Deutsche Gärt. Zeit. 25: 475. f. 9, No. 21.

On page 49, vol. 1, under *Opuntia caribaea*, insert: Dr. Britton endeavored to find this plant in Trinidad in 1920 and 1921 but failed and he could not learn anything about it. It appears probable that the drawing sent by Mr. Lockhart to Kew in 1825 was made from a Venezuelan plant.

On page 54, vol. 1, under *Opuntia clavellina*, add to illustration: Karsten and Schenck, Vegetationsbilder 13: pl. 18, in part.

On page 56, vol. 1 under *Opuntia whipplei* in last line of description read cm as mm.

*The Index Kewensis refers this name to *Opuntia leucacantha*, but the illustration shows that it belongs to *Nopalea*.

† Lemaire in his Les Cactées, published in 1868, takes up the name *Cactus* for certain of the low, depressed, much branched or cespitose species of *Opuntia*. He lists a number of these on pages 8 and 88, but as they are not connected through published species their identification is made only by inference.

Add to illustration: Bull. Agr. Exper. Sta. N. Mex. **78**: pl. 11, 12; North Amer. Fauna **7**: pl. 9; Pac. R. Rep. **4**: pl. 17, f. 1 to 4; Stand. Cycl. Hort. Bailey **4**: f. 2609.

On page 57, vol. 1, under *Opuntia acanthocarpa*, add to illustration: Stand. Cycl. Hort. Bailey **4**: f. 2606; Gartenwelt **11**: 75.

On page 57, vol. 1, under *Opuntia echinocarpa* and *O. parryi*, respectively, add the synonyms: *Cactus echinocarpus* and *C. parryi* Lemaire, Cactées 88. 1868.

On page 58, vol. 1, under *Opuntia bigelovii*, add to illustrations: MacDougal, Bot. N. Amer. Des. pl. **47**: Shreve, Veg. Des. Mt. Range pl. 4; Contr. U. S. Nat. Herb. **16**: pl. 10; Stand. Cycl. Hort. Bailey **4**: f. 2607; Karsten and Schenck, Vegetationsbilder **4**: pl. 40, B.

On page 61, vol. 1, under *Opuntia cholla*, insert: *Opuntia chella* (Index Kew. Suppl. **1**: 302) is a typographical error for *O. cholla*.

On page 62, vol. 1, under *Opuntia versicolor*, add to illustrations: Carnegie Inst. Wash. **269**: pl. 8, f. 81; pl. 9 MacDougal, Bot. N. Amer. Des. pl. 58; Plant World **9**¹²: f. 50.

On page 63, vol. 1, under *Opuntia imbricata*, add the synonym: *Cactus imbricatus* Lemaire, Cactées 88. 1868. Also add to distribution: Oklahoma.

Insert: Rydberg (Fl. Rocky Mountains 576. 1917) reports this species from Utah under the name of *Opuntia arborescens*; we have seen no specimens of it from Utah.

Insert: *Cactus subquadriflorus* Mociño and Sessé (De Candolle, Prodr. **3**: 471. 1828), given as a synonym of *Opuntia rosea*, doubtless belongs here. Schumann's reference, *C. quadriflorus*, is incorrect. *C. subquadrifolius* (Cactaceae **3**: 65) is a clerical error.

Add to illustrations: Dict. Gard. Nicholson Suppl. 179. f. 195, as *Opuntia decipiens*; Dict. Gard. Nicholson **4**: 581. f. 52, as *O. rosea*; Stand. Cycl. Hort. Bailey **4**: f. 2608; Engler and Drude, Veg. Erde **13**: f. 28, in part; Gartenwelt **4**: 159, as *O. arborescens*; Bot. Jahrb. Engler **58**: Beibl. **129**: 33. f. 10.

On page 66, vol. 1, under *Opuntia tunicata*, add to illustrations: Garden **13**: 107,* as *Opuntia exuviata*; Möllers Deutsche Gärt. Zeit. **25**: 476. f. 9, No. 7; Goebel, Pflanz. Schild. **1**: f. 36, as *O. stapeliae*; Contr. U. S. Nat. Herb. **10**: pl. 17, f. A.

On page 68, vol. 1, under *Opuntia fulgida*, add to illustrations: MacDougal, Bot. N. Amer. Des. pl. 57, as *Opuntia mamillata*; MacDougal, Bot. N. Amer. Des. pl. 87.

On page 68, vol. 1, under *Opuntia spinosior*, insert: This plant is sometimes found in the trade as *Opuntia arborescens spinosior* (see Grässner).

Add to illustrations: Emory, Mil. Recon. App. 2. f. 10, as *Opuntia arborescens*; Shreve, Veg. Des. Mt. Range pl. 2, A.

On page 71, vol. 1, under *Opuntia vestita*, insert: Illustration: Möllers Deutsche Gärt. Zeit. **25**: 476. f. 9, No. 8.

On page 73, vol. 1, under *Opuntia clavarioides*, add to illustrations: Garden **13**: 107, as *Opuntia clavarioides cristata*; Rother, Praktischer Leitfaden Kakteen 106; Möllers Deutsche Gärt. Zeit. **15**: 67; **25**: 476. f. 9, No. 19; Thomas, Zimmerkultur Kakteen Wiener Ill. Gärt. Zeit. **28**: f. 18; Monatsschr. Kakteenk. **32**: 131.

On page 73, vol. 1, under *Opuntia salmiana*, insert: Extend range to central Argentina and habit to rocky hillsides (according to W. B. Alexander).

On page 75, vol. 1, under *Opuntia subulata*, add to illustrations: Deutsche Gärt. Zeit. **8**: 32, as *Pereskia subulata*; Haage and Schmidt, Haupt-Verz. Kakteen **1919**: 169; Goebel, Pflanz. Schild. **1**: f. 5; Möllers Deutsche Gärt. Zeit. **25**: 476. f. 9, No. 15.

On page 78, vol. 1, under *Opuntia cylindrica*, add to illustrations: Möllers Deutsche Gärt. Zeit. **25**: 476. f. 9, No. 12; Gartenwelt **15**: 539; Rother, Praktischer Leitfaden Kakteen 107; Cact. Journ. **1**: 100; Schelle, Handb. Kakteenk. **42**. f. 4, as *Opuntia cylindrica cristata*; Wiener Illustr. Gartenz. **29**: f. 22, No. 10; De Laet, Cat. Gén. f. 88; Monatsschr. Kakteenk. **13**: 71; Schelle, Handb. Kakteenk. **42**. f. 3.

*This illustration is very poor and is only tentatively referred here. If native to California, as one might infer from the account which accompanies the illustration, it may refer to a form of *Opuntia prolifera* or *O. echinocarpa*.

On page 80, vol. 1, under *Opuntia stanlyi*, add the synonym: *Cactus emoryi* Lemaire, Cactées 88. 1868.

Also add to illustrations: Schelle, Handb. Kakteenk. 38. f. 1, as *Opuntia emoryi*; Nat. Geogr. Mag. 21: pl. on p. 716, as *O. kunzei*.

On page 80, vol. 1, under *Opuntia schottii*, insert: *Opuntia greggii* occurs only in Schumann's Index (Gesamtb. Kakteen 829) with page reference to *O. schottii greggii*.

On page 81, vol. 1, under *Opuntia clavata*, insert the synonym: *Cactus clavatus* Lemaire, Cactées 88. 1868.

Add to illustrations: Stand. Cycl. Hort. Bailey 4: f. 2605.

On page 82, vol. 1, under *Opuntia pulchella*, add to illustration: MacDougal, Bot. N. Amer. Des. pl. 26, as *O. pusilla*.

On page 83, vol. 1, under *Opuntia bulbispina*, insert: *Cactus bulbispinus* Lemaire. (Cactées 88. 1868) was intended as a synonym of this species.

On page 89, vol. 1, under *Opuntia glomerata*, insert: Extend range to central and northern Argentina.

Insert: *Tephrocactus polyacanthus* (Index Kewensis Suppl. 1: 421) was intended for *T. platyacanthus* Lemaire (Förster, Handb. Cact. ed. 2. 915. 1885).

Add to illustrations: Watson, Cact. Cult. ed. 1 and 2. 257. f. 97; ed. 3. f. 60, as *Opuntia papyracantha*; Dict. Gard. Nicholson 2: 503. f. 755; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 1, as *O. platyacantha*; Schelle, Handb. Kakteenk. 45. f. 7, as *O. andicola*; De Laet, Cat. Gén. f. 60; Rev. Hort. Belg. 40: after 186; Schelle, Handb. Kakteenk. 44. f. 6; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 2, as *O. diademata*.

On page 92, vol. 1, under *Opuntia aoracantha*, add to illustrations: Schelle, Handb. Kakteenk. 44. f. 5.

On page 93, vol. 1, under *Opuntia hickenii*, insert: Mr. W. B. Alexander suggests that *Opuntia platyacantha* Spegazzini (not Salm-Dyck) is probably a synonym of this species.

On page 94, vol. 8, insert:

64a. *Opuntia wetmorei* sp. nov.

Forming low mounds of considerable extent with hundreds of branches; joints 4 to 10 cm. long, terete, turgid, 2. cm. in diameter or less, slightly tapering towards each end, dull green, but




Fig. 234.—*Opuntia wetmorei*, fruit, stem, and seeds.

dull purple around and especially below the areoles; leaves subtending the minute areoles, 1 to 2 mm. long, caducous; areoles circular, bearing tawny or white wool when young; glochids short, yellowish; spines numerous, very unequal, scarcely pungent, white to straw-colored or brownish, 3 or 4 of lower ones almost hair-like, reflexed or appressed to joints, 3 or 4 of uppermost erect or ascending, flattened, 2 to 3.5 cm. long; flowers not known; immature fruit glabrous at first, dull green, becoming reddish purple especially about the areoles, 3 cm. long, bearing long white bristly spines, especially from upper areoles, deeply umbilicate.

Collected by W. B. Alexander in the barranca of the Tunuyán River near Tunuyán, Mendoza, Argentina, March 22 and 23, 1921.

This species is perhaps nearest *Opuntia darwinii*. We are under great obligation to W. B. Alexander for sending us very fine living plants by Alexander Wetmore, who brought them to us directly from Argentina. Mr. Wetmore was with Mr. Alexander when the plant was collected and he has given us a word picture of the plant; we take pleasure in naming the species for him, not only in recognition of this service but also for obtaining other valuable specimens of cacti.

Figure 234 is from a photograph of the type plant, one-half natural size.

On page 95, vol. 1, under *Opuntia corrugata*, insert: *Tephrocactus rectrospinus* (Index Kewensis Suppl. 1: 421) is a misspelling for *T. rectrospinosus* Lemaire.

Also insert: *Illustrations*: Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 11; 488. f. 22, No. 8.

On page 95, vol. 1, under *Opuntia ovata*, add:

Opuntia pusilla Salm-Dyck (Observ. Bot. 3: 10. 1822. Not Haworth, 1812) was referred by Schumann to *O. corrugata*. We have seen a photograph of Haworth's specimen (bearing the date November 8, 1824) which seems to answer to Salm-Dyck's plant which we would refer here.

On page 96, vol. 1, under *Opuntia sphaerica*, add the synonym: *Opuntia ovata leonina* Schelle, Handb. Kakteenk. 46. 1907.

Also add to illustrations: Deutsche Gärt. Zeit. 7: 313, as *Opuntia leonina*; Schelle, Handb. Kakteenk. 46. f. 8, as *O. grata leonina*.

On page 97, vol. 1, under *Opuntia pentlandii*, add the synonym: *Cactus bolivianus* Lemaire, Cactées 88. 1868.

Also add to illustrations: Watson, Cact. Cult. ed. 3. 106. f. 54; Deutsche Gärt. Zeit. 7: 312; Schelle, Handb. Kakteenk. 58. f. 16, as *Opuntia boliviana*; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 14.

On page 99, vol. 1, insert:

76a. *Opuntia alexanderi* sp. nov.

Low, depressed, forming a small clump; joints readily detached, grayish green, strongly tubercled, globose, 2 to 3 cm. in diameter, nearly hidden by the numerous spines; areoles small, close together, circular; spines 4 to 12, up to 4 cm. long, flexible, white below, dark above or with black tips, scurfy-pubescent even in age; flowers not known; fruit red, dry, obovoid, 2 cm. long, lower areoles not spiny, but upper ones bearing 2 to 8 long, white, erect, weak spines overtopping the fruit; umbilicus of fruit depressed; seeds white, 5 to 6 mm. broad.

Collected by W. B. Alexander, between Chilecito and Famatina, province of La Rioja, Argentina, February 19, 1921. Mr. Alexander studied this species in the field but could not identify it and sent it to us for study. It belongs to the subgenus *Tephrocactus*, but is not near any of the known species. We take great pleasure in naming it for Mr. Alexander, who has extensively studied the cacti in Argentina.

On page 100, vol. 1, under *Pumilae*, add to distribution: Venezuela.

On page 100, vol. 1, under *Opuntia pumila*, insert: Illustration: Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 5.

On page 101, vol. 1, under *Opuntia pubescens*, add the synonym: *Cactus pubescens* Lemaire, Cactées 87. 1868.

On page 102, vol. 1, under *Opuntia curassavica*, add to illustrations: Dillenius, Hort. Elth. 2: pl. 295, as tuna; Loudon, Encycl. Pl. 413. f. 6897, as *Cactus curassavicus*; Knorr, Thesaurus pl. o.2.

On page 102, vol. 1, insert:

80a. *Opuntia abjecta* Small, sp. nov.

Prostrate, often growing in large irregular patches on almost bare limestone or where some sand and humus has accumulated, irregularly branched; joints suborbicular, sometimes nearly subglobose, oval, or broadly obovate, mostly 4 to 8 cm. long, very thick, frequently turgid, light green, loosely attached to each other; leaves ovoid to conic-ovoid, 2 to 3 mm. long, ascending and slightly curved upward, green or purplish; glochids yellowish; spines setaceous-acicular, mostly solitary, brown, or reddish purple, mottled light and dark, becoming chalky gray when dry; the larger ones 2 to 6 cm. long; flowers usually solitary on a joint; berry urceolate, 1 to 1.5 cm. long, somewhat tuberculate, red or purple-red, rounded at base; umbilicus relatively broad, concave; seeds few, flattish, about 4 mm. wide.

On edge of hammock, southern end of Big Pine Key, Florida. Type collected in May 1921 by J. K. Small, preserved in the herbarium of the New York Botanical Garden.

Similar to *Opuntia drummondii* but with shorter joints, longer and more slender spines, and different fruit.

On page 105, vol. 1, under *Opuntia drummondii*, add to illustration: Journ. Elisha Mitchell Sci. Soc. 34: pl. 13, 14.

On page 105, vol. 1, under *Opuntia tracyi*, insert:

Type Locality: Biloxi, Mississippi. *Distribution:* Southern Mississippi, southeastern Georgia to northern Florida.

On page 105, vol. 1, insert:

86a. *Opuntia impedata* Small, sp. nov.

Prostrate, ultimately copiously branched, the joints often piled several layers deep and forming viciously armed mats, elliptic or oblong, mostly 7 to 15 cm. long, rather thick, pale green;



FIG. 235.—*Opuntia impedata*.

leaves stout-subulate, 4 to 6 mm. long, erect or ascending, slightly curved upward, dark green; glochids brownish; spines subulate, usually numerous, solitary or 2 together, light gray, except the brown tip, salmon-colored when dry, and faintly banded when wet; flowers often several on a joint; ovary obconic, nearly terete; sepals green, outer lanceolate to ovate, 4 to 8 mm. long, acuminate, the inner much larger, with shoulders of very broad body narrowed into stoutish tip; corolla bright yellow, 4.5 to 5.5 cm. wide; petals about 12, 2.5 to 3 cm. long, broadly obovate to cuneate-obovate, broadly rounded at apex, mucronate; anthers nearly 2 mm. long; berry clavate, about 3 cm. long, narrowed at base; umbilicus rather small, somewhat concave; seeds rather few, 4 to 4.5 mm. in diameter.

Sand dunes, northeastern Florida. Type in the herbarium of the New York Botanical Garden; collected on dunes at Atlantic Beach, Florida, in April 1921, by J. K. Small.

Dr. Small notes that the stiff spines may penetrate leather shoes and that the plant is very prolific, both vegetatively and through its fruit.

Figure 235 is from a photograph taken by Dr. Small of the type plant.

On page 110, vol. 1, insert:

Series 3a. PISCIFORMES.

Plants in dense colonies with turgid, very spiny, narrow, deep green joints, the spines conspicuously long and slender, salmon-colored in the first year, gray in the second; flowers numerous, bright yellow; berry turbinate-obovoid, 4 cm. long or less. The only species inhabits Florida.

96a. *Opuntia pisciformis* Small, sp. nov.

Prostrate, copiously branched, forming dense mats often 1 to 3 meters in diameter, with joints piled several layers deep, roots fibrous; joints narrowly elliptic, linear-elliptic, or spatulate, mostly 1 to



FIG. 236.—*Opuntia pisciformis*.

3 dm. long, very thick, deep green, readily detached; leaves stout-subulate, 2 to 4 mm. long, incurved; areoles rather prominent, mostly armed; spines solitary or 2 or 3 together, cream-colored, becoming salmon-colored and gray with a dark tip when dry, salmon when wet, the longer ones 5 to 6 cm. long; flowers numerous; ovary turbinate, angular and tuberculate; sepals green, the outer lanceolate to triangular-lanceolate, 9 to 12 mm. long, acuminate, the inner much larger, the broad ovate or suborbicular base broadly tapering into the very stout tip; corolla bright yellow, 6 to 7 cm. wide; petals about 12, 3 to 4 cm. long, broadly cuneate, mostly truncate or emarginate at apex, mucronate; anthers nearly 2 mm. long; berry broadly turbinate-obovoid, 3.5 to 4 cm. long, purple, narrowed at base, the umbilicus deeply concave; seeds rather numerous, 5 to 5.5 mm. in diameter.

Sand dunes, estuary of the Saint Johns River, Florida. Type in the herbarium of the New York Botanical Garden; collected on dunes at Atlantic Beach, Florida, in April 1921, by J. K. Small,

Figure 236 is from a photograph by Dr. Small of the type plant.

On page 113, vol. 1, under *Opuntia tuna*, in first line read 1769 as 1768.

Add the synonyms: *Cactus horridus* Salisbury, Prodr. 348. 1796; *Opuntia tuna humilior* Salm-Dyck, Cact. Hort. Dyck. 1849. 66. 1850.

Insert: *Opuntia maidenii* Griffiths (Bull. Torr. Bot. Club 46: 201. 1919) described from a cultivated plant sent from Australia and grown at Chico, California, seems referable to this species.

Add to illustrations: Loudon, Encycl. Pl. 411. f. 6880, as *Cactus polyanthos*; Monatsschr. Kakteenk. 6: 25, as *Opuntia polyantha*; Deutsche Gärt. Zeit. 7: 447, as *O. humilis*; Watson, Cact. Cult. ed. 3. f. 62; Cact. Journ. 2: 169; Useful Wild Plants U. S. Canada, opp. 18, 108, 174; Stand. Cycl. Hort. Bailey 4: f. 2599; Schelle, Handb. Kakteenk. 51. f. 13; Remark, Kakteenfreund 24.

On page 115, vol. 1, under *Opuntia antillana*, insert: *Opuntia domingensis* appears without description in Urban's Symbolae (8: 466. 1920). It was a manuscript name for which *O. antillana* was substituted.

On page 117, vol. 1, under *Opuntia decumbens*, add to illustrations: Bull. U. S. Dept. Agr. 31: pl. 7, f. 1, as *Opuntia puberula*; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 3.

On page 119, vol. 1, under *Opuntia basilaris*, insert: *Opuntia dorffii* is advertised by Haage and Schmidt (Monatsschr. Kakteenk. 29: September). We have had a cutting which we would refer to one of the forms of *O. basilaris*.

Also add to illustrations: Cact. Journ. 2: 163, as *Opuntia basilaris albiflora*; Cact. Journ. 11: pl. for October; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 13, as *O. basilaris cordata*; Möllers Deutsche Gärt. Zeit. 25: f. 9, No. 9, as *O. basilaris minima*; Watson, Cact. Cult. ed. 3. f. 53; Deutsche Gärt. Zeit. 7: 312; Remark, Kakteenfreund 23; Monatsschr. Kakteenk. 7: 125; Stand. Cycl. Hort. Bailey 4: f. 2597; Gartenflora 31: 280; Schelle, Handb. Kakteenk. 47. f. 10.

On page 121, vol. 1, under *Opuntia microdasys*, add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 488. f. 2, No. 4, as *Opuntia microdasys monstrosa*; Garden 13: 107,* as *O. pubescens*; Schelle, Handb. Kakteenk. 47. f. 9; Möllers Deutsche Gart. Zeit. 25: 476. f. 9, No. 16; Karsten and Schenck, Vegetationsbilder 2: pl. 22, B.

On page 123, vol. 1, under *Opuntia pycnantha*, insert: *Opuntia pycnacantha* (Just's Jahresb. 24²: 380. 18) seems to have been a misspelling for *O. pycnantha*.

On page 127, vol. 1, under *Opuntia opuntia*, add the synonym: *Opuntia compressa* Macbride, Contr. Gray Herb. 11. 65: 41. 1922.

Also add to illustrations: Contr. U. S. Nat. Herb. 21: pl. 23, B; Bailey, Sand Dunes Indiana 94; Ann. Inst. Roy. Hort. Fromont 2: pl. 1, f. F; Deutsches Mag. Gart. Blumen. 1869: pl. 17. opp. 257; Kraemer, Appl. Econ. Bot. f. 341, as *Opuntia vulgaris*; Watson, Cact. Cult. 212. f. 84; Ann. Rep. Bur. Amer. Ethn. 33: pl. 20, A; Clements and Clements, Rocky Mt. Fl. pl. 32, f. 6; Clements, Fl. Mount. Plain pl. 32, f. 7, as *O. humifusa*; Wiener, Ill. Cart. Zeit. 2: 40. f. 10, as *O. rafinesquiiana*; Deutsche Gärt. Zeit. 7: 447; Wiener Ill. Gärt. Zeit. 2: f. 112, as *O. rafinesquiiana arkansana*; Watson, Cact. Cult. ed. 3. f. 61;

*This illustration is very poor and the identification is based largely upon the description.

Schelle, Handb. Kakteenk. 50. f. 12; Belg. Hort. 26: pl. 8; Illustr. Hort. 15: pl. Opp. 51; Deutsches Mag. Gart. Blumen. 1869: pl. 17, Opp. 257, as *O. rafinesquei*; Kraemer, Appl. Econ. Bot. f. 341.

On page 130, vol. 1, insert:

121a. *Opuntia eburnispina* Small, sp. nov.

Prostrate, widely branched and forming mats on dune sands, with tuberous roots; joints oval or suborbicular, varying to broadest above middle, thickish, 6 to 13 cm. long, pale green, somewhat shining, especially when young; leaves ovoid-subulate, 4 to 5 mm. long, pale green, recurved-spreading; spines relatively stout, 2 to 4 at an areole or sometimes solitary, 1 to 2 cm. long, ivory-white with yellowish tips when young, becoming dark gray, not spirally twisted, greenish when wet; flowers few; ovary obconic; sepals triangular, green, 5 to 7 mm. long; corolla clear yellow, 4 to 5 cm. wide; petals few, narrowly cuneate, often minutely pointed; berries obovoid, 2 cm. long or less.

Coastal sands, Cape Romano, Florida. Type specimens in the herbarium of the New York Botanical Garden; collected in May 1922, by J. K. Small.

Figure 237 is from a photograph by Dr. Small of the type plant.




FIG. 237.—*Opuntia eburnispina*.

On page 131, vol. 1, under *Opuntia macrorhiza*, add to illustrations: Watson, Cact. Cult. ed. 3. f. 59 Dict. Gard. Nicholson 4: 580. f. 50, 51.

On page 131, vol. 1, under *Opuntia tortispina*, add the synonym: *Opuntia cymochila montana* Engelm., Proc. Amer. Acad. 3: 296. 1856.

Also add to distribution: Southeastern Colorado. Established and slowly spreading east of Cincinnati, Ohio (E. T. Wherry).

Also add to illustrations: Watson, Cact. Cult. ed. 3. pl. 102; Meehans' Monthly 11: 57, as *Opuntia mesacantha*; Meehans' monthly 5: 172, as *O. oplocarpa*.

On page 134, vol. 1, *Opuntia sulphurea*, insert: Mr. W. B. Alexander writes as follows concerning this species:

"This is by far the commonest species of *Opuntia* in the Argentine, where it is commonly known as 'penca,' i. e. the spiny plant, sometimes being distinguished from other larger species by the name 'penquilla' or 'penca chica.' The writer met with it in the provinces of Buenos Aires, Córdoba, San Luis, Mendoza, San Juan, La Rioja, Catamarca and Santiago del Estero."

Add to illustrations: Wiener Ill. Gärt. Zeit. 28: f. 17, as *Opuntia maculacantha*; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 18.

On page 134 vol. 1, under *Opuntia soehrensii*, add the synonyms: *Cactus ayrampo* Azara, Voy. 2: 526. 1809; *Opuntia haenquiiana* Herrera, Rev. Univ. Cuzco 8: 60. 1919.

Also insert: Azara's original description is interesting and a translation of it is given:

"A species of tunilla (cactus) which is found in the temperate gorges near the Cordillera produces the seed in question. The plant is found in arid and sterile soil where ordinarily this family of plants grows and thrives by creeping on the ground in such a way as to stifle all the others. From the seed confined within the round and spiny fruit is derived a color of a clear violet, brilliant and extremely agreeable to the eye but very superficial and very light, although it acquires a little stability and durability by the means of alum and some other chemicals."

On page 135, vol. 1, insert:

129a. *Opuntia macbridei* sp. nov.

A low bush, 6 dm. high, forming broad impenetrable thickets on gravelly river flats; joints obovate, 6 to 8 cm. broad, 8 to 15 cm. long, glabrous, at first light green, in age dark green; leaves minute, 1 to 2 mm. long, caducous; areoles on young joints hemispheric, brown-felted and



FIG. 238.—*Opuntia macbridei*.

with brown glochids, on old joints 2 to 3 cm. apart; spines 2 to 4, in age gray to horn-colored, with yellowish tips, very unequal, the longest up to 5 cm. long, stout-subulate; flowers very small, orange to orange-red; petals only 4 to 5 mm. long; ovary tuberculate, bearing many brown-felted tubercles but without spines, deeply umbilicate; fruit deeply umbilicate, red to purple.

Collected by Macbride and Featherstone at Huanuco, Peru, altitude 2,300 meters, August 28 to September 3, 1922 (No. 2365, type), and April 8, 1923 (No. 3250).

Mr. Macbride states that the seeds are brown. All the fruits we have seen were sterile; these sterile fruits on falling to the ground take root and form new plants.

This interesting plant, which proves to be undescribed, we have named for Mr. J. Francis Macbride, who led the Botanical Expedition of 1922 to South America, sent out by the Field Museum of Natural History, under the Captain Marshall Field fund.

Figure 238 is from a photograph showing the habit of this plant.

On page 135, vol. 1, under *Opuntia penicilligera*, insert: Mr. W. B. Alexander sends us the following account of this plant:

"This plant was met with close to the coast at Bahia Blanca, and near the foot of the Andes at Tunuyán. As remarked by Spegazzini, this species is very distinct from any other found in Argentina and there seems no reason for thinking that it may belong to the Series *Sulphureae* in

which it is tentatively placed in the Cactaceae. It should surely either be the type of a separate series or be placed in the Series *Basilares*, to the members of which, judging by illustrations, it shows great resemblance."

On page 138, vol. 1, under *Opuntia pottsii*, add to illustrations: Watson, Cact. Cult. ed. 3. 1. 58; Dict. Gard. Nicholson 4:580. f. 49, as *Opuntia filipendula*.

On page 145, vol. 1, under *Opuntia phaeacantha*, add to illustrations: Deutsche Gärt. Zeit. 7: 447, as *Opuntia camanchica*; Meehans' Monthly 11: 57, as *O. phaeacantha major*; Shreve, Veg. Des. Mt. Range pl. 5, A, as *O. toumeyii*; De Laet, Cat. Gén. f. 58.

On page 147, vol. 1, under *Opuntia engelmannii*, add the synonym: *Opuntia engelmannii discata* C. Z. Nelson, Trans. Ill. State Acad. Sci. 12: 124. 1919.

Also add to illustrations: Cact. Journ. 1: pl. for February; 2: 162, as *Opuntia engelmannii cristata*; Gard. Chron. 111. 39: 148. f. 58; Plant World 9¹²: f. 49; Shreve, Veg. Des. Mt. Range pl. 5, B; Stand. Cycl. Hort. Bailey 4: f. 2601; Scientific Month. 17: 70, 71, 72.

On page 149, vol. 1, under *Opuntia discata*, add to illustrations: Carnegie Inst. Wash. 269: pl. 10, f. 87.

On page 153, vol. 1, under *Opuntia bergeriana*, add to illustrations: Gartenwelt 11: 75.

On page 153, vol. 1, under *Opuntia elatior*, add to illustrations: Loudon, Encycl. Pl. ed. 3. 411. f. 6879, as *Cactus nigricans*.

On page 155, vol. 1, under *Opuntia boldinghii*, add to distribution: Chacachacare and Patos Islands, Trinidad.

On page 156, vol. 1, under *Opuntia vulgaris*, insert: *Opuntia gracilior* (Index Kewensis 3 357. 1894) is a mistake for *O. inonacantha gracilior* Lemaire.

Add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 20, as *Opuntia monacantha variegata*; Pl. Utiles Madagascar 124. f. 39; 125. f. 39.

On page 158, vol. 1, under *Opuntia arechavaletai*, add to illustration: Karsten and Schenck, Vegetationsbilder 11: pl. 1.

On page 158, vol. 1, under *Opuntia bonaerensis*, insert: Mr. W. B. Alexander writes of this species as follows:

This species was seen only on rocky slopes in the Sierra de la Ventana in the south of the province of Buenos Aires. It is known only from the few Sierras which rise from the pampas in the east of the province. There is little doubt that it is nearly related to *Opuntia vulgaris* Miller (*O. monacantha* Haworth) which was found by the writer at Rio de Janeiro and is familiar in Australia.

Add to illustration: Anal. Mus. Nac. Montevideo 5: pl. 33, as *Opuntia chakensis*.

On page 159, vol. 1, insert after *Opuntia scheeri*: *Opuntia diversispina* Griffiths (Bull. Torr. Club 46: 197. pl. 9. 1919) grown from seed of unknown origin at Brownsville, Texas, is described as similar to *O. scheeri* and in the accompanying illustration the joints resemble those of that species.

On page 160, vol. 1, *Opuntia chlorotica*, add to illustrations: Bull. N. Mex. Coll. Agr. No. 78. pl. [4]; Stand. Cycl. Hort. Bailey 4: f. 2600.

On page 161, vol. 1, under *Opuntia laevis*, add to illustrations: MacDougal, Bot. N. Amer. Des. pl. 56.

On page 163, vol. 1, under *Opuntia dillenii*, add to illustrations: Garden 13: 107,* as *Opuntia crassa*; Bull. Torr. Club 46: pl. 10, as *O. maritima*; Lindley, Veg. King. ed. 3. 746. f. 498, No. 1, 2 Knorr, Thesaurus pl. o; Watson, Cact. Cult. ed. 3. f. 6.

On page 163, vol. 1, insert:

174a. *Opuntia ochrocentra* Small, sp. nov.

Erect, 1 meter tall or less, much branched or sometimes diffuse, with fibrous roots; joints elliptic to oval varying to broadest above the middle x to 3 dm long thickish light green not

* This illustration is not very good for this species. It is, however, the same one that Nicholson used (f. 757) and that W. Watson used (f. 86) as *Opuntia tuna*, which we have referred here.

repand; leaves ovoid, 2 to 4 mm. long, often purplish; areoles rather prominent; glochids yellowish brown; spines 5 to 6 together or sometimes fewer on new joints, yellow, stiff, subulate, reflexed, becoming gray when dry, yellowish green when wet, straight, the longer ones 4.6 to 5 cm. long; flowers rather few; ovary turbinate, even; sepals often purple-tinged, deltoid to rhombic-orbicular or rhombic-reniform, acute; corolla bright lemon-yellow, 7 to 8.5 cm. wide; petals few, cuneate, somewhat crisped; berry obovoid, red, about 2 cm. long.

On edge of hammock, southeastern end of Big Pine Key, Florida. Type specimens collected in December 1921, by J. K. Small, in the herbarium of the New York Botanical Garden.

Related to *O. dillenii*, differing in shape of the joints, which are not repand, and the strongly reflexed, scarcely flattened spines.

On page 166, vol. 1, under *Opuntia lindheimeri*, add to illustrations: Journ. Hered. Washington, 6⁴: f. 19, as *Opuntia ellisiana*; Journ. Hered. Washington 6⁴: f. 15, 16, as *O. cacanapa*; Journ. Hered. Washington 6⁴: f. 17, 18; as *O. subarmata*; Journ. Hered. Washington 5: 223. f. 13; Schulz, 500 Wild Fl. San Antonio pl. 12.

Also insert: Dr. Small has found this plant established, after cultivation, in pine lands west of Halenville, Florida.

On page 167, vol. 1, under *Opuntia cantabrigiensis*, add to illustrations: Gartenwelt 10: 560; Gard. Chron. III. 33: 98. f. 42.

Also insert: Professor Duncan S. Johnson found this species naturalized on sand dunes at Beaufort, North Carolina, in 1899, and Doctor Small studied it there in 1922.

At Cambridge, England, it has passed through many winters out of doors.

On page 168, vol. 1, under *Opuntia beckeriana*, insert: *Opuntia prostrata spinosior* (Schumann, Gesamtb. Kakteen 723. 1898) seems to have been a garden name which Schumann would refer to *O. beckeriana*.

On page 173, vol. 1, under *Opuntia tomentosa*, add to illustrations: Blanc, Cacti 82. No. 2200, as *Opuntia lurida*; Reiche, Elem. Bot. f. 165; Gartenwelt 11: 75.

On page 175, vol. 1, under *Opuntia leucotricha*, add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 4, as *Opuntia leucacantha*; Cassell's Dict. Gard. 2: 138; Bull. U. S. Dept. Agr. 31: pl. 6, f. 2; pl. 7, f. 2; U. S. Dept. Agr. Bur. Pl. Ind. Bull. 262: pl. 4; pl. 5, f. 1.

Insert: Dr. John K. Small has found this plant naturalized in a hammock south of Fort Pierce, Florida, where it is reported as established during the Seminole wars.

On page 176, vol. 1, under *Opuntia orbiculata*, add to the illustrations: Schelle, Handb. Kakteenk. 48. f. 11, as *Opuntia crinifera*; Gartenwelt 11: 76, as *O. lanigera*.

In third line of description on page 177 read cm. as dm.

On page 178, vol. 1, under *Opuntia ficus-indica*, add to illustration: Engler and Prantl, Pflanzenfam. 3^{6a}: f. 57, H; Gard. Chron. III. 34: 89. f. 34; 92. f. 42; Karsten, Deutsche Fl. 887. f. 501. No. 10, 11; ed. 2. 2: 456. f. 605. No. 10, 11; Journ. Dept. Agr. S. Austr. 13: 764; Garten-Zeitung 4: 182. f. 42, No. 1; Stand. Cycl. Hort. Bailey 4: f. 2598; Watson, Cact. Cult. ed. 3. f. 57.

On page 180, vol. 1, under *Opuntia maxima*, add the synonym: *Cactus maximus* Colla, Mem. Accad. Sci. Torino 33: 140. 1826 (?).

Also insert: *Illustration*: Möllers Deutsche Gärt. Zeit. 25: 488. f. 22, No. 3, as *Opuntia labouretiana*.

On page 181, vol. 1, under *Opuntia hernandezii*, insert: *Opuntia hernandezii* first appeared in De Candolle's Prodrum (3: 474. 1828).

Also insert: *Nopal silvestre* Thierry (Förster, Handb. Cact. ed. 2. 929. 1885) is cited as a synonym of *Opuntia hernandezii*. This reference is given also in the Index Kewensis.

Also insert: *Illustration*: Förster, Handb. Cact. ed. 2. 930. f. 128.

On page 184, vol. 1, under *Opuntia streptacantha*, add to illustrations: Useful Wild Pl. U. S. Canada opp. 18, 108, 174, as *Opuntia tuna*.

On page 185, vol. 1, under *Opuntia megacantha*, insert: *Opuntia effulgia* Griffiths (Bull. Torr. Club 46: 195. 1919) was obtained from San Luis Potosí, Mexico, and grown at Chico, California; *O. hispanica* Griffiths (Bull. Torr. Club 46: 198. 1919) was described from a plant received from Spain and grown at Chico; *O. chata* Griffiths (Bull. Torr. Club 46: 199. 1919), from Aguascalientes, Mexico, was grown at Brownsville, Texas, and at Chico; *O. obovata* Griffiths (Bull. Torr. Club 46: 202. 1919) from Hepasote, Mexico, was also grown at Brownsville and at Chico; *O. amarilla* Griffiths (Bull. Torr. Club 46: 205. 1919) was obtained in cultivation at Cardenas, Mexico, and grown at Chico. These are known to us only from descriptions and appear to be races of *O. megacantha* or of some of the related tall, white-spined species.

Add to illustrations: Ann. Rep. Smiths. Inst. 1917: pl. 16, f. 2.

On page 191, vol. 1, under *Opuntia robusta*, insert: *Opuntia cyanea* Griffiths (Bull. Torr. Club 46: 196. 1919) judging from the original description may be related to *O. robusta*.

Add to illustrations: Engler and Prantl, Pflanzenfam. 3^{6a}: f. 56, g, as *Opuntia albicans*.

On page 194, vol. 1, *Opuntia fragilis*, add to illustrations: Watson, Cact. Cult. ed. 3. f. 5; Deutsche Gärt. Zeit. 7: 313; Remark, Kakteenfreund 22, as *Opuntia brachyarthra*; Schelle, Handb. Kakteenk. 6. f. 15, as *O. fragilis brachyarthra*; Meehans' Monthly 11: 57.

On page 195, vol. 1, under *Opuntia arenaria*, add to illustration: Meehans' Monthly 11: 57.

On page 195, vol. 1, *Opuntia erinacea*, add the synonym: *Opuntia ursus horribilis* Walton, Cact. Journ. 2: 152. 1899.

Also add to illustrations: Cact. Journ. 1: 93, as *Opuntia*; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 10; Cycl. Amer. Hort. Bailey 3: 1149. f. 1548; Stand. Cycl. Hort. Bailey 4: 2363. f. 2603, as *O. ursina*; Meehans' Monthly 4: 9; Monatsschr. Kakteenk. 14: 10; N. Amer. Fauna 7: pl. 11, as *O. rutila*.

On page 198, vol. 1, under *Opuntia rhodantha*, add to illustrations: Monatsschr. Kakteenk. 30: 153, as *Opuntia xanthostemma*.

On page 199, vol. 1, under *Opuntia polyacantha*, add the synonym: *Opuntia missouriensis watsonii* Schumann, Gesamtb. Kakteen 735. 1898.

Also insert: Extend range to northwestern Oklahoma.

Add to illustrations: Rep. Mo. Bot. Gard. 13: pl. 13; Schelle, Handb. Kakteenk. 54. f. 14, as *Opuntia missouriensis*; Möllers Deutsche Gärt. Zeit. 25: 476. f. 9, No. 6, as *O. schwärziana*; Scientific American 124: 492; Meehans' Monthly 11: 57; Stand. Cycl. Hort. Bailey 4: f. 2604.

On page 201, vol. 1, under *Opuntia grandis*, add to illustration: The Garden 62: 425; Möllers Deutsche Gärt. 25: 476. f. 9, No. 17.

On page 203, vol. 1, under *Opuntia nashii*, insert: *Illustration*: Journ. N. Y. Bot. Gard. 6: f. 3.

On page 204, vol. 1, *Opuntia spinosissima*, insert at end: A species of this series, *Spinossissimae*, occurs on Navassa Island off the southeastern point of Haiti; specimens were sent us by Mr. F. P. Dillan, Superintendent of Light Houses, San Juan, Porto Rico, but they are not complete enough to be specifically referred.

On page 206, vol. 1, under *Opuntia moniliformis*, add the synonyms: *Cactus reticulatus* Index Kewensis 1: 369. 1893; * *Opuntia reticulata* Karsten, Deutsche Fl. ed. 2. 2: 457. 1895; *Opuntia picardae* Urban, Repert. Sp. Nov. Fedde 16: 35. 1919.

On page 208, vol. 1, under *Opuntia rubescens*, add to illustration: Carnegie Inst. Wash. 269: pl. 10, f. 90. 91, as *Opuntia catacantha*.

*The Index Kewensis refers *Cactus reticulatus* to Descourtilz (Fl. Med. Antill. 1: pl. 68), but the formal name was not used by him.

On page 209, vol. 1, under *Opuntia brasiliensis*, add to distribution: Peru.

Insert: *Opuntia brasiliensis gracilior* Salm-Dyck was given by Förster (Handb. Cact. 500. 1846) as a synonym of *O. brasiliensis minor*.

Also insert: Dr. Small has found this plant established after planting on shell mounds and waste places in southern Florida.

Add to illustrations: Goebel, Pflanz. Schild. 1: f. 37, 38.

On page 211, vol. 1, under *Ammophilae*, substitute for characters of the series:

Erect species, sometimes with a definite continuous trunk, often much branched, the joints broad and flat, spiny or unarmed, the spines (when present) subulate or subulate-acicular, whitish, gray or reddish, the large flowers yellow.

The series now appears to be most nearly related to the Series *Tortispinae* (vol. 1: 126) and may be placed to follow it as series a. *Opuntia austrina* Small, of southern Florida, may be transferred from the *Tortispinae* to the *Ammophilae*.

On page 211, vol. 1, under *Opuntia ammophila*, insert: More recent collections of this plant by Dr. Small, show that its range extends south to Cape Romano, Florida, and that the definite trunk, at first taken as characteristic of it, is not always developed; his living plants from different stations show slight individual differences which do not appear to be specific. This species has been erroneously referred by Dr. Griffiths (Bull. Torr. Club 46: 201) to *Opuntia bartramii* Rafinesque.

On page 213, vol. 1, insert:

239a. *Opuntia turgida* Small, sp. nov.

Plant erect, more or less diffusely branched, 0.5 meter tall or less, with fibrous roots; joints elliptic to elliptic-obovate, 5 to 12 cm. long, thickish, deep green, sometimes slightly glaucous when young; leaves subulate, 6 to 10 mm. long, spreading and more or less recurved, green, sometimes accompanied by fine bristles, but without spines; areoles scattered, often prominent and densely bristly on the older joints; spines (as far as known) wanting; flowers often several on a joint; ovary obovoid or obconic-obovoid, 2 to 2.5 cm. long, slightly tubercled; sepals green or purple-tinged, the outer subulate to lanceolate, 4 to 10 mm. long, acute, the inner rhombic-ovate, fully .5 cm. long, stout-pointed; corolla bright yellow, 5.5 to 6.5 cm. wide; petals 10 to 12, about 3 cm. long, broadly cuneate, abruptly narrowed, rounded or subtruncate at the apex, mucronate; anthers 2 mm. long; berry obovoid, 2 to 2.5 cm. long, greenish purple, even, broadly rounded at the base, the umbilicus flat or a little depressed at the middle; seeds rather numerous, about 4 mm. in diameter, somewhat turgid.

Hammocks near Yulee and on the mainland along the Halifax River south of Daytona, Florida. Type collected about five miles south of Daytona, in December 1919, by J. K. Small, preserved in the herbarium of the New York Botanical Garden.

This spineless, small-jointed species is tentatively referred to the Series *Ammophilae* on account of its fruit characters and erect habit. A plant sent from Kew to the New York Botanical Garden in 1902, under an unpublished name, very closely resembles this species.

On page 214, vol. 1, insert the following:

Opuntia napolea, offered for sale by Grässner (Monatsschr. Kakteenk. February 1920) we have not seen.

The name *Opuntia spirocentra* Engelmann and Bigelow (Haage, Verz. Cact. 30), found in the Index Kewensis, we have not been able to verify. As the name is credited to Engelmann and Bigelow and the habitat of the plant is said to be New Mexico it is doubtless an error and probably was intended for *O. macrocentra*.

Opuntia todari (Haage and Schmidt, Haupt-Verz. 230. 1912) is known only in the trade.

Cactus italicus referred by the Index Kewensis to Tenore (Steudel, Nom. ed. 2. 2: 246. 1840) occurs first in 1831 (Tenore, Syll. Pl. Neop. 241) where also occurs the name *Opuntia italica*. Both are unpublished but doubtless refer to some species of *Opuntia*.

CACTUS PARVIFOLIUS Ehrenberg in F. G. Dietrich, Vollst. Lex. Gaertn. 2: 416. 1802.

An upright, cylindrical, almost articulate stem; the upper part bedecked with small, cylindrical, fleshy, pointed leaves; on lower part of the stem, at the place where the leaves are attached, stiff bristles are formed which are surrounded at the base by a whitish-gray, woolly substance; in old age the stem requires a support on account of its slender growth; if the stem is cut through in the middle and the wound well dried, young sprouts make their appearance at this place which serve to propagate the plant. South America is its home.

The above paragraph is a free translation of the description.

We have not been able to identify this plant, but it is probably some species of *Opuntia* or possibly *Tacinga funalis*.

Cereus vulnerator Cortes (Fl. Colombia 69. 1897) and *C. guasabara* Cortes (Fl. Colombia 208. 1897) are different names for the same plant. From the brief descriptions it is difficult to identify this plant but it certainly is not a *Cereus*. It suggests some sheathed-spined *Opuntia* such as *O. tunicata* which has been introduced into South America and is common in northern Ecuador. It is known as curuntilla or guasabara in Colombia.

CORRECTIONS AND ADDITIONS TO VOLUME II.

On page 4, vol. II, under *Cereus hexagonus*, add the synonyms: *Cereus regalis* Haworth, Suppl. Pl. Slice. 75. 1819; *Cactus regalis* Sprengel, Syst. 2: 476. 1825; *Cereus childsi* Blanc, Cacti 39, No. 375.

Insert: *Cereus cyaneus* Hortus is listed by Berger (Hort. Mortola 69. 1912) as a South American plant grown at La Mortola. From drawings sent by Berger it is probably to be referred to *C. hexagonus*.

Add to illustrations: Andrews, Bot. Rep. 8: pl. 513; Reichenbach, Fl. Exot. pl. 322; Van Géel, Sert. Bot. 1: pl. 114, as *Cactus hexagonus*; Blanc, Cacti 39, No. 375, as *Cereus childsi*.

On page 8, vol. II, under *Cereus jamacaru* insert: *Cereus caracore* (Gosselin, Bull. Soc. Acclim. France 51: 58. 1905) belongs to the group containing *C. jamacaru*, that is, it is a true *Cereus*, according to Gosselin. He does not claim that it is a good species. No species of *Cereus*, however, are natives of Chile, from which this plant is said to have come. If indigenous to that country it is more likely to be *Trichocereus chiloensis*.

On page 9, vol. II, under *Cereus jamacaru*, add to illustrations: Monatsschr. Kakteenk. 26: 181; Karsten, Deutsche Fl. ed. 2. 2: 456. f. 605, No. 8.

On page 11, vol. II, under *Cereus peruvianus*, add the synonyms: *Piptanthocereus peruvianus* Riccobono, Boll. R. Ort. Bot. Palermo 8: 232. 1909; *Piptanthocereus peruvianus monstrosus* Riccobono, Boll. Ort. Bot. Palermo 8: 233. 1909.

Also add to illustrations: Saint-Hilaire, Exp. Fam. Nat. 2: pl. 95, in part as f. 1(?); De Candolle, Pl. Succ. Hist. 1: pl. 8, as *Cactus peruvianus*; Blanc, Cacti 36. No. 252; Rother, Praktischer Leitfaden Kakteen 15, as *Cereus peruvianus monstrosus*; Karsten and Schenck, Vegetationsbilder 1: pl. 41; 42, f. b; Gartenwelt 6: 133; Mem. Acad. Roy. Sci. pl. 4, 5; Haage and Schmidt, Haupt-Verz. 1919: 134. f. 10737; Goebel, Pflanz. Schild. 1: f. 5, 53.

On page 14, vol. II, under *Cereus pernambucensis*, add to illustration: Remark, Kakteenfreund 7, as *Cereus formosus monstrosus*.

On page 17, vol. II, under *Cereus aethiops*, add to illustrations: Förster, Handb. Cact. ed. 2. 207. f. 15, as *Cereus landbeckii*; Blanc, Cacti 26. No. 27; Gartenwelt 16: 537, as *Cereus coerulescens*.

On page 19, vol. II, insert:

25. *Cereus trigonodendron* Schumann, Bot. Jahrb. Engler **40**: 413. 1908.

Simple, or in age with a much branched top, 15 meters high; trunk 5 meters long, smooth, dm. in diameter or more; ribs to 6, 2 to 3 cm. high; areoles in young growth 2 to 3 cm. apart, producing abundant white wool, 1 cm. long or more; spines 4 to 7, at first brown, subulate, 2 to 5.5 cm. long; flowers as in typical species of *Cereus*, 10 to 15 cm. long; fruit smooth, edible.

Type locality: Department of Loreto, Peru.

Distribution: Valleys of eastern Peru and Bolivia.

This species is briefly described on page 19 of volume II of The Cactaceae, but at that time we knew little about it and were disposed to exclude it from the genus *Cereus*. We have since had a photograph of the type specimen from Berlin. In December 1922 F. L. Herrera sent us flowers from the Santa Ana Valley, province of Convención, Peru, and in February 1923 we received herbarium specimens of branches and flowers from W. E. Meyer, collected in 1922 at Cachuela-Esperanza, Boni, Bolivia. It is found only in the Atlantic drainage of Peru and Bolivia and is therefore geographically within the range of the genus *Cereus* as limited by us.

Illustration: Bot. Jahrb. Engler **40**: pl. 10.

Figure 239 is from a photograph sent by Dr. Vaupel from Berlin.




FIG. 239.—*Cereus trigonodendron*.

On page 20, vol. II, add at end of page: *Cereus amalonga* and its variety *cristata* are described in the Cactus journal (**2**: 93, 104, 119) and both are illustrated in the plate for August of that volume. They are said to have been imported from Mexico. We are unable to identify these plants either from the descriptions or illustrations.

On page 21, vol. II, under *Cereus lormata*, insert: We listed *Cereus lormata* among the species unknown to us but we have since seen an illustration (Wiener Ill. Gart. Zeit. **11**: pl. 3, in part) of a barren plant. It has about 10 vertical ribs with clusters of subulate spines, some of them greatly elongated. It is probably not a true *Cereus*.

On page 21, vol. II, at end of *Cereus* add: *Cereus perviridis* Weingart is advertised by Haage & Schmidt (Cat. 1914). We have seen a cutting but do not know its relationship.

Cereus pitahaya variabilis Weingart (Monatsschr. Kakteenk. **16**: 158. 1906) is only a form and is not described.

Cereus roezlii Haage jr. (Schumann, Gesamtb. Kakteen 64. f. 12. 1897) was described as columnar with 9 obtuse ribs, 9 to 12 radial spines, and one central spine much larger than the radials. Its flowers were unknown. It is said to come from the Andes of Peru or Ecuador. It is probably some species of *Lemaireocereus* or *Trichocereus*.

Cereus stolonifer Weber is listed by Schumann (Monatsschr. Kakteenk. 5: 43. 1895) as a plant grown in the Botanical Garden in Paris.

Cereus tripteris Salm-Dyck (De Candolle, Prodr. 3: 468. 1828) was described from barren plants of unknown origin and has never been identified.

Cereus uspenshi Haage jr. is mentioned in a report by Karl Hirscht (Monatsschr. Kakteenk. 8: 109. 1898).

Cereus auratus Labouret (Rev. Hort. iv. 4: 27. 1855) is a tall *Cereus*-like plant, originally reported as from Peru, but the Index Kewensis says it is from Mexico. The four following varieties: *genuinus*, *intermedius*, *mollissimus*, and *pilosus* are briefly described by Regel & Klein (Ind. Sem. Hort. Petrop. 1860: 4. 1860); *Pilocereus auratus* (Rümpel in Förster, Handb. Cact. ed. 2. 650. 1885) is doubtless the same.

On page 22, vol. II, under *Monvillea cavendishii*, add to illustrations: Blühende Kakteen 3: pl. 171, as *Cereus euchlorus*; Blühende Kakteen 3: pl. 172, as *C. rhodoleucanthus*; Blühende Kakteen 3: pl. 178, as *C. cavendishii*.

On page 23, vol. II, under *Monvillea spegazzinii*, add the synonyms: *Piptanthocereus spegazzinii* Riccobono, Boll. R. Ort. Bot. Palermo 8: 233. 1909; *Cereus spegazzinii hassleri* Weingart, Monatsschr. Kakteenk. 32: 163. 1922.

Add to illustrations: De Laet, Cat. Gén. f. 28, as *Cereus spegazzinii*.

On page 27, vol. II, under *Cephalocereus senilis*, add to illustrations: Journ. Intern. Gard. Club 3: 640, as *Cephalocereus* sp.; Gard. Chron. III. 32: 35; Journ. Hort. Home Farm. III. 59: 625; Amer. Garden 11: 479; West Amer. Sci. 13: 16, as *Cereus senilis*; West Amer. Sci. 13: 23, as *C. hoppenstedtii*; Möllers Deutsche Gart. Zeit. 25: 473. f. 5, No. 19; Remark, Kakteenfreund 20, as *Pilocereus hoppenstedtii*; Cact. Journ. 1: pl. 5; Gartenflora 27: 114; Deutsche Gärt. Zeit. 6: 64; Gard. Chron. 1873: f. 15; Garten-Zeitung 4: 182. f. 42, No. 6; Gartenwelt 2: 574; 116: 175; Watson, Cact. Cult. ed. 2. 260. f. 98; ed. 3. f. 34; West Amer. Sci. 9: 2; Journ. Intern. Gard. Club 3: 640; Blanc, Cacti 76. No. 1755; Weinberg, Cacti 26, as *Pilocereus senilis*; Palmer, Cult. Cact. 148, as *Pilocereus*; Engler and Drude, Veg. Erde 13: f. 30; Tribune Hort. 4: 283; Möllers Deutsche Gart. Zeit. 25: 473. f. 5, No. 3; Schelle, Handb. Kakteenk. 108. f. 44, 45; Floralia 42: 370; Balt. Cact. Journ. 1: 116.

On page 30, vol. II, under *Cephalocereus fluminensis*, add to illustrations: Goebel, Pflanz. Schild. 11: pl. 3, f. 1 to 3, as *Pilocereus*.

On page 31, vol. II, under *Cephalocereus macrocephalus*, add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 2.

On page 32, vol. II, under *Cephalocereus polylophus*, insert: *Pilocereus angulosus* Förster, according to Lemaire (Rev. Hort. 1862: 428. 1862) is little known; it is perhaps to be referred here.

Add to illustration: Gard. Chron. III. 50: 135. f. 64, c, as *Cereus polylophus*.

On page 42, vol. II, under *Cephalocereus arrabidaei*, insert: The following names relate to this species and other names associated with it: *Pilocereus sublanatus* Förster (Haage, Verz. Cact. 22) is referred to *Cereus sublanatus* by the Index Kewensis. *Pilocereus tilophorus* (Index Kewensis) is evidently a mistake for *Cereus tilophorus*. *Pilocereus oligogonus* Lemaire (Rev. Hort. 1862: 428. 1862) is said to come from Mexico; the two varieties, *houlettianus* and *sublanatus*, given at this same place as synonyms, may or may not belong with it; they should doubtless be referred to the species bearing the same names respectively.

Add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 9, as *Pilocereus exerens*.

On page 44, vol. II, under *Cephalocereus nobilis*, add the synonyms *Cereus polyptychus*

Lemaire, Cact. Gen. Nov. Sp. 56. 1839; *Pilocereus polyptychus* Rümpler in Förster, Handb. Cact. ed. 2. 680. 1885.

Insert: The plant upon which this name was based was a small, barren one of unknown origin.

Insert: *Pilocereus houlletianus niger* (Förster, Handb. Cact. ed. 2. 676. 1885) is only a name given as a synonym of *P. niger*, while *P. niger aureus* is briefly described on the same page.

Add to illustrations: Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 4, as *Pilocereus curtisi*.

On page 47, vol. II, under *Cephalocereus polygonus*, add the synonym: *Cephalocereus sch-lumbergeri* Urban, Symb. Antill. 8: 464. 1920.

On page 49, vol. II, under *Cephalocereus lanuginosus*, add the synonym: *Pilocereus lanuginosus virens* Salm-Dyck in Förster, Handb. Cact. ed. 2. 672. 1885.

Insert: Curran reports that this fruit is edible (Inventory No. 50. p. 50. U. S. Dept. Agr. Bur. Plant Industry).

On page 51, vol. II, under *Cephalocereus royenii*, add to illustrations: Journ. N. V. Bot. Gard. 15: pl. 133, 134.

On page 52, vol. II, under *Cephalocereus leucocephalus*, add to illustrations: Watson, Cact. Cult. 145. f. 6; Deutsche Gärt. Zeit. 7: 312, as *Pilocereus houlletianus*; Gard. Chron. III. 29: f. 79, as *P. houlletianus leucocephalus*; Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 14, as *P. cometes*; De Laet, Cat. Gén. No. 51, 52, 53 Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 7; Blühende Kakteen 2: pl. 79; West Amer. Sci. 13: 24; Schelle, Handb. Kakteenk. 101. f. 40. as *P. houlletii*; Gard. Chron. III. 32: 253, as *Cereus houlleti*.

On page 56, vol. II, under *Cephalocereus purpusii*, insert: Wilhelm Weingart, under date of June 18, 1921, wrote of this species as follows:

"*Cephalocereus purpusii* sp. nov. was collected by C. A. Purpus in 1902 near Mazatlán, was sent to me February 18, 1907, and bloomed in Darmstadt in 1918."


Figures 240 and 241 are reproduced from a drawing furnished by Wilhelm Weingart.

On page 56, vol. II, under *Cephalocereus cat-tingicola*, add to illustrations: Vegetationsbilder 6: pl. 14, as *Cereus cattingicola*; Engler, Bot. Jahrb. 40: Suppl. pl. 5, as *C. cattingae*.

On page 58, vol. II, insert:

49. *Cephalocereus collinsii* sp. nov.

About 3 meters high with few slender, elongated branches, these 3 to 4 cm. in diameter; ribs about 7, obtuse; tubercles about 1.5 cm. apart, circular, long-woolly as well as spiny; flowering areoles not much more woolly than the others; spines numerous, acicular, longer ones 3 to 4 cm. long; flowers borne near ends of branches, about 5 cm. long; fruit somewhat depressed, about 3 cm. broad; seed shining, black, 1.5 to 2 mm. broad.



FIGS. 240, 241.—*Cephalocereus purpusii*, flower.

Common in thickets near Tehuantepec, Oaxaca, Mexico. The type was collected by Dr. C. A. Purpus near Gerónimo in April 1923. It was reported by O. F. Cook and C. N. Collins from this region in 1902. The plant is named for Mr. Collins, who first brought it to our attention more than 20 years ago while carrying on field work in southern Mexico for the U. S. Department of Agriculture.

Figure 242 is from a photograph taken by Mr. Collins in 1902; it is three-fourths

natural size.

On page 58, vol. II, under *Cephalocereus hermentianus*, add: Illustration: Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 10, as *Pilocereus hermentianus*.

On page 58, vol. II, under *Pilocereus albisetosus*, add the synonyms: *Cactus albisetosus* Sprengel, Syst. 2: 496. 1825; *Cactus albisetus* Steudel, Nom. ed. 2. 1: 245. 1840.

On page 61, vol. II, under *Espostoa lanata*, add to illustrations: Schelle, Handb. Kakteenk. 105. f. 41, as *Pilocereus lanatus*; Schelle, Handb. Kakteenk. 105. f. 42, as *P. lanatus cristatus*; Wiener Ill. Gart. Zeit. 11: pl. 3, in part, as *P. dautwitzii*.

On page 64, vol. II, under *Stetsonia coryne*, insert: W. B. Alexander wrote, under date of March 7, 1921, as follows:

"Noticing your statement that the fruit of *Stetsonia coryne* is unknown, I obtained a ripe specimen at La Rioja for you and am sending it by parcel post."

This we describe as follows:

Oblong, 6 cm. long, glabrous, bearing scattered scales, these 5 mm. broad, 1 mm. high, each with a cartaceous tip and a denticulate margin; seeds numerous, small, 1.5 mm. long, flattened, pitted; hilum large, basal.

On page 65, vol. II, under *Stetsonia coryne*, add to illustrations: Thomas, Zimmerkultur Kakteen 11, as *Cereus coryne*.

On page 66, vol. II, under *Escontria chiotilla*, add to illustrations: Möllers Deutsche Gärt. Zeit. 29: 438. f. 13; Flöralia 42: 389, as *Cereus chiotilla*.

On page 69, vol. II, under *Pachycereus pringlei*, insert: The distribution of *Pachycereus pringlei* in northern Sonora is not well defined. Dr. MacDougal has recently visited northwestern Sonora and states that he saw it along the route between Altar and Port Libertad to within a hundred miles of the United States boundary. Prospectors and ranchers also speak of it as being abundant in the valley of the Asuncion or Altar River some miles to the northward. He writes of it as follows:

"On the whole, however, my chief interest was centered on the sowesa or *Pachycereus pringlei*. We began to get into this about 85 miles from the Gulf, and in the region below a thousand feet it attains perfectly tremendous size, as you will see from some photographic prints."

Figure 243 is from a photograph obtained by Dr. MacDougal at Port Libertad, Sonora, May 4, 1923.




FIG. 242.—*Cephalocereus collinsii*.

Also add to illustrations: Zeitschr. Ges. Erdk. **1916**: f. 6, in part; Contr. U. S. Nat. Herb. **16**: 131, A; 132, A; Karsten and Schenck, Vegetationsbilder **13**: pl. 13, as *Pachycereus calvus*; Contr. U. S. Nat. Herb. **16**: pl. 131, B, as *P. titan*; Ann. Rep. Smith. Inst. **1908**: 553. f. 17.




FIG. 243.—*Pachycereus pringlei*.

On page 71, vol. II, under *Pachycereus pecten-aboriginum*, add to illustrations: Engler and Drude, Veg. Erde **13**: 297. f. 9; Watson, Cact. Cult. ed. 2. 246. f. 92 ed. 3. f. 17; Karsten and Schenck, Vegetationsbilder **1**: pl. 48, as *Cereus pecten-aboriginum*; Karsten and Schenck, Vegetationsbilder **13**: pl. 14; Contr. U. S. Nat. Herb. **16**: pl. 132, B.

On page 71, vol. II, under *Pachycereus gaumeri*, add additional characters:

Ribs sometimes only 3, thin, 3 to 4 cm. high; areoles sometimes 2.5 cm. apart; fruit becoming dry, globose, 3 to 4 cm. in diameter, scales at base of fruit small, becoming long and foliaceous above, fleshy at base but tips thin and soon drying black; axils of scales felted, with a cluster of about 8 very short black spines; seeds numerous, brown, 4 mm. long.

The above description is drawn from specimens sent by Dr. Gaumer to Washington in June 1922.

On page 73, vol. II, under *Pachycereus chrysomallus*, add to illustrations: Reiche, Elem. Bot. 226. f. 162, as *Pilocereus*; Möllers Deutsche Gärt. Zeit. **29**: 297. f. 5; U. S. Dept. Agr. Bur. Pl. Ind. Bull. **262**: pl. 13, f. 1, as *Pilocereus fulviceps*; Flörealia **42**: 377; Belg. Hort. **3**: pl. 57, as *P. chrysomallus*; Möllers Deutsche Gärt. Zeit. **25**: 473. f. 5, No. 1.

On page 74, vol. II, under *Pachycereus marginatus*, insert: The two varieties, *Cereus marginatus monstrosus* (Monatsschr. Kakteenk. **19**: 62. 1909) and *C. marginatus cristatus* (Monatsschr. Kakteenk. **4**: 194. 1894) occur in the trade.

Also add to illustrations: West Amer. Sci. **13**: 6; Möllers Deutsche Gärt. Zeit. **25**: 472. f. 2, No. 15; De Laet, Cat. Gén. f. 23; Möllers Deutsche Gärt. Zeit. **29**: 355. f. 11; Bot. Jahrb. Engler **58**: Beibl. **129**: 27. f. 9, as *C. marginatus*; Remark, Kakteenfreund 7; Karsten and Schenck, Vegetationsbilder **1**: pl. 43; 48, as *Cereus gemmatus*; Ann. Rep. Smiths. Inst. **1908**: pl. 11, f. 2.

On page 76, vol. II, under *Pachycereus lepidanthus*, insert: Since the appearance of volume II, we have received flowers of this species from Wilhelm Weingart, which show a very close likeness to those of *Escontria chiotilla*. The fruit of the latter, however, is a juicy edible berry, while that of the former is described as dry. The illustrations here printed may lead to the rediscovery of this rare plant.

Figure 244 is from a photograph of a plant grown in Washington, showing a joint as it came from the field and also the young growth as developed in the greenhouse; figure 245 is from a photograph of two flowers and a spine-cluster.




FIG. 244.—*Pachycereus lepidanthus*.

On page 78, vol. II, under *Leptocereus leonii*, insert: Specimens collected in June 1923, by Brother Leon and Dr. Roig on Loma de Somorrostro, Jamaica, Havana Province, Cuba, show that the fruit of this species becomes 6.5 cm. long by 5.5 cm. thick, when fully mature. The fruits are borne near the ends of the branches, 3 or 4 close together.

On page 76, vol. II, under *Cereus tetazo*, insert: *Pilocereus tetetzo cristatus* Weber (Schumann, Gesamtb. Kakteen 176. 1897) is only a name and so is *Cephalocereus tetetzo* (Monatsschr. Kakteenk. 19: 73. 1909) and *Cereus tetetzo* and *C. tetetzo* (Monatsschr. Kakteenk. 17: 79. 1907).




FIG. 245.—*Pachycereus lepidanthus*.

Also insert: *Illustration*: Bull. Soc. Nat. Acclim. 52: 55. f. 14, as *Cereus tetetzo*.

On page 82, vol. II, under *Eulychnia spinibarbis*, add to illustrations: Engler and Drude, Veg. Erde 8: pl. 5, f. 11, as *Cereus coquimbamus*.

On page 86, vol. II, under *Lemaireocereus hollianus*, add to illustrations: Bull. Soc. Nat. Acclim. **52**: 45. f. 9, as *Cereus baxosus*.

On page 86, vol. II, under *Lemaireocereus hystrix*, add the synonym: *Cactus americanus* Vitman, Summa Pl. **3**: 209. 1789.

Insert: *Cactus americanus* is based on Bradley's illustration (Hist. Succ. Pl. 12) which De Candolle referred to *Cereus eburneus*, but as the plant came from the West Indies it is perhaps better referred to *Lemaireocereus hystrix*.

Also insert: We have recently obtained from N. E. Brown a photograph of Haworth's *Cereus hystrix*, with the date, "Oct. 24, 1824."

On page 87, vol. II, under *Lemaireocereus griseus*, add to illustrations: Monatsschr. Kakteenk. **24**: 5, as *Cereus eburneus*; Ann. Rep. Smiths. Inst. **1908**: pl. 9, f. 5.

On page 89, vol. II, under *Lemaireocereus eichlamii*, add to the description:

Fruit globular, about 5 cm. in diameter, becoming dry, not edible, thin-skinned, filled with numerous large seeds, the surface bearing scattered areoles, these densely short-felted with clusters of short spines subtended by small, ovate, acute scales; seeds black, 4 to 5 mm. long with a prominent hilum.

Insert: This plant is much used for hedges in Salvador and was obtained there by Mr. Paul C. Standley in the vicinity of Sonsonate, altitude 220 to 300 meters, March 1922 (No. 22328), but was not seen in the wild state. It is called there órgano. This species heretofore has been known only from Guatemala and was not known to us in fruit; this differs from that of the other species of *Lemaireocereus* in being rather dry with very large seeds.

Also insert: Figure 246 is from a photograph of the plant, sent by F. Eichlam in 1909 to Washington, which flowered in 1918.




FIG. 246.—*Lemaireocereus eichlamii*.

On page 91, vol. II, under *Lemaireocereus chende*, add to illustrations: Grässner, Haupt-Verz. Kakteen **1912**: 3, as *Cereus del moralii*.

On page 96, vol. II, under *Lemaireocereus weberi*, add to illustrations: Floralia **42**: 388, as *Cereus candelabrum*.

On page 96, vol. II, insert:

14a. *Lemaireocereus benecke* (Ehrenberg).

Cereus benecke Ehrenberg, Bot. Zeit. **2**: 835. 1844.

Cereus farinosus Haage in Salm-Dyck, Allg. Gartenz. **13**: 355. 1845.

Cereus benecke farinosus Salm-Dyck, Cact. Hort. Dyck. 1849. 49. 1850.

Piptanthocereus benecke Riccobono, Boll. R. Ort. Bot. Palermo **8**: 226. 1909.

Plants 4 to 5 meters high, much branched; branches 6 to 7 cm. in diameter, the growing tips very glaucous; ribs 7 or 8, strongly tuberculate, obtuse, separated by narrow intervals; areoles small, circular, borne on the upper side of the tubercle, brown to black-felted; spines x to 7, acicular, the longest sometimes 2.5 cm. long, brown to black; flowers night-blooming, small, 4 cm. long, greenish brown without; inner perianth-segments rose-colored to white (?); ovary globose, glaucous, tuberculate, its areoles brown-felted and bearing 3 to 7 acicular spines, the longest sometimes 2.5 cm. long and brown to black; fruit about 2 cm. in diameter, somewhat tubercled, bearing clusters of spines at the areoles, red; pericarp thick, somewhat fleshy; pulp disappearing, leaving the large seeds loose, these escaping by a basal pore as in *Oreocereus* and many of the *Echinocactanae*.

Type locality: Mexico on red lava beds.

Distribution: Central Mexico.

In volume 11 of *The Cactaceae* (p. 18), we described this plant under *Cereus* but with the statement that it was not a true *Cereus*; we were not then able to refer it to any known genus. At that time we knew little about the flowers and nothing accurate about the ovary and fruit. In 1921 Professor K. Reiche sent us some living plants from Iguala, the station from which Dr. Rose obtained his plants in 1905. These contained some old withered flowers and some well-developed ovaries which have enabled us to refer the plant to *Lemaireocereus*.

Figure 247 is from a photograph of K. Reiche's plant, slightly reduced, showing the top of a branch bearing an old flower and a half-ripe fruit.




FIG. 247.—*Lemaireocereus beneckeii*.

On page 98, vol. 11, under *Lemaireocereus thurberi*, add to illustrations: Journ. N. Y. Bot. Gard. 3: f. 13, as *Cereus thurberi*; Bull. U. S. Nat. Mus. 56: pl. 8, f. 2; Karsten and Schenck, Vegetationsbilder 13: pl. 15; 21, f. A; Contr. U. S. Nat. Herb. 16: pl. 125, A; Amer. Bot. 20: 88.

On page 108, vol. 11, under *Bergerocactus emoryi*, add to illustrations: Cact. Journ. 1: 59; Gartenwelt 11: 498, as *Cereus emoryi*.

On page 111, vol. 11, under *Wilcoxia poselgeri*, add to illustrations: Remark, Kakteenfreund 6; Deutsche Garten-Zeitung 1886: f. 25, as *Cereus tuberosus*.

On page 111, vol. II, under *Wilcoxia striata*, insert: According to T. S. Brandege (under date of June 8, 1921), the flowers of *Wilcoxia striata* are nocturnal.

On page 112, vol. II, under *Peniocereus greggii*, add to illustrations: Amer. Gard. 11: 474, as *Cereus greggii*; Journ. Wash. Acad. 12: 329. f. i; Succulenta 4: 71.

On page 113, vol. II, insert:

2. *Peniocereus johnstonii* Britton and Rose, Journ. Wash. Acad. 12: 329. 1922.

A climbing or clambering plant, up to 3 meters long, with a very large fleshy root sometimes weighing 14 pounds; stems and branches 3 to 5-angled, the young growth not pubescent; spines 9 to 12, brown to black, glabrous; upper radial spines short, stubby, swollen at base, nearly black, the two lower light brown, elongated, bristle-like, reflexed; central spines 1 to 3, subulate, 4 to 8 mm. long; flower (only an old flower seen) about 15 cm. long; perianth-segments about 3 cm. long; the lower and outer ones bearing tawny hairs and long bristles; flower-tube slender, with prominent areoles on knobby projections and bearing tawny wool and bristly spines; fruit ovoid to oblong, about 6 cm. long, bearing prominent clusters of black spines, dry (?), many-seeded; seeds oblong, 3 mm. long or more, black, shining; seedling dark purple; cotyledons very thick, triangular.

Type locality: San Josef Island, off the east coast of southern Lower California.

Distribution: Southern Lower California.

This plant was always found growing up through bushes of *Olneya tesota*.

Illustrations: Journ. Wash. Acad. 12: 330. f. 2 Succulenta 4: 73.

Figure 248 shows a branch, old flowers, and seeds of the type specimen.




FIG. 248.—*Peniocereus johnstonii*, showing branch, old flower, and seeds.

On page 113, vol. II, under *Dendrocereus nudiflorus*, insert: In 1922 Dr. L. H. Bailey sent us two photographs and some stem-sections (No. 806) which he had obtained from the Botanic Garden at Roseau, Dominica. It grows as a low, rounded, much branched bush with the outer joints often pendent. Mr. Joseph Jones, curator of the Botanic Garden wrote that the group is made up of six plants which have not been cut back or interfered with in any way and have experienced two hurricanes without having a piece broken off. One of our colleagues, Dr. William R. Maxon, who had rediscovered this plant some years ago in Cuba, suggested that the plant grown in Dominica might be that species; a careful study of our material convinces us that he is correct. *Dendrocereus nudiflorus*, however, is naturally a large tree with a very definite trunk and a large, much branched top. An explanation of this inconsistency is that the Dominican plant was doubtless grown from cuttings, causing it to assume this bushy habit, a phenomenon also observed in other cacti.

Also insert: *Cereus undiflorus* is a misspelling, used by Sauvalle (Fl. Cuba 59. 1873) and reprinted in the Index Kewensis (1: 493).

Figure 249 is from one of the photographs sent us by Dr. Bailey.

On page 116, vol. II, under *Machaerocereus eruca*, add to illustrations: Journ. Intern. Gard. Club 3: 641; Karsten and Schenck, Vegetationsbilder 13: pl. 16, as *Cereus eruca*.



FIG. 249.—*Dendrocereus nudiflorus*.

On page 117, vol. II, under *Machaerocereus gummosus*, add to illustrations: Cact. Journ. 2: 107, as *Cereus gummosus*; Zeitschr. Ges. Erdk. 1916: f. 6, in part; Karsten and Schenck, Vegetationsbilder 13: pl. 17, f. A.

On page 119, vol. II, under *Nyctocereus serpentinus*, add to illustrations: Watson, Cact. Cult. 67: f. 16; ed. 3. f. 12, as *Cereus serpentinus*.

On page 119, vol. II, under *Nyctocereus guatemalensis*, add to illustrations: Monatsschr. Kakteenk. 31: 41, as *Cereus hirschtianus*.

On page 123, vol. II, under *Acanthocereus pentagonus*, add to illustrations: De Laet, Cat. Gén. f. 32, as *Cereus baxaniensis*; Monatsschr. Kakteenk. 32: 21, as *C. princeps*.

On page 125, vol. II, insert the following:

3a. *Acanthocereus floridanus* Small, sp. nov.

Stems and branches diffusely spreading or reclining, 3 to 10 meters long, stout: joints prominently 3 to 5-angled, but mostly 3-angled, dark green, often forming impenetrable thickets: areoles remote, with mostly 4 to 7 slender or subulate spines, the central one often 1 to 2 cm. long: ovary stout-trumpet-shaped, 8 to 10 cm. long, with few large, separated tubercled areoles at the base, bearing mostly 3 to diverging spines, those on the upper part usually with one spine each; outer perianth-segments deltoid to triangular-lanceolate or lanceolate-subulate and almost linear, the longer ones 3.5 to 4 cm. long, acuminate; inner perianth-segments broadly linear, 3.5 to 4.5 cm. long, about six times as long as wide, broadly acuminate; filaments adnate more than halfway up from the base of the hypanthium; anthers less than 2.5 mm. long.

Hammocks, along or near the coast, southern peninsular Florida, adjacent islands, and Florida Keys. Type collected by J. K. Small, on Key Largo, December 1917 and 1918; preserved in the herbarium of the New York Botanical Garden.

This Florida plant has been referred by us to *A. pentagonus*, but specimens recently collected by Dr. Small, including good flowers, which we had not seen before, indicate it to be a distinct species, characterized by its much shorter perianth and more spiny ovary.

Illustrations: Britton and Rose, Cactaceae 2: 123. f. 182; 124. f. 184, as *Acanthocereus pentagonus*.

On page 129, vol. II, under *Heliocereus speciosus*, add to the illustrations: Herb. Génér. Amat. 4: pl. 244; Colla, Hort. Ripul. pl. 10; Bonpl. Descr. Pl. Rar. pl. 3, as *Cactus speciosus*;

Maund, Bot. 1: pl. 12, as *C. speciosus lateritius*; Curtis's Bot. Mag. 49: pl. 2306 Edwards's Bot. Reg. 6: pl. 486, as *C. speciosissimus*; Edwards's Bot. Reg. 19: pl. 1596, as *C. speciosissimus lateritius*; Newman, Illustr. Bot. 209; Abh. Bayer. Akad. Besch. Cact. 2: pl. 3, f. 5; De Laet, Cat. Gén. f. 24, as *Cereus speciosus*; Lindley, Veg. King. ed. 3. 746. f. 498; Curtis's Bot. Mag. 67: pl. 3822; The Garden 53: 153, as *C. speciosissimus*; Illustr. Hort. 32: pl. 548, as *C. speciosissimus hoveyi*; Sci. Amer. 124: 492, as *Heliocereus mallisoni*; Van Géel, Sert. Bot. 1: 116, as *Cactus speciosissimus*.

On page 129, vol. II, under *Heliocereus cinnabarinus*, add: *Illustration*: Monatsschr. Kakteenk. 32: 54, 55, as *Cereus cinnabarinus*.

On page 129, vol. II, under *Heliocereus amecamensis*, add to illustrations: Rother, Praktischer Leitfaden Kakteen 74, as *Cereus amecamensis*.

On page 132, vol. II, under *Trichocereus spachianus*, add to illustrations: Remark, Kakteenfreund 5, as *Cereus spachianus*.

On page 133, vol. II, under *Trichocereus pasacana*, insert: The name *Cephalocereus pasacana* (Engler and Prantl, Pflanzenfam. 3^{6a}: 182. 1894) has been used for this plant.



FIG. 250.—*Borzicactus fieldianus*.



FIG. 251.—*Borzicactus fieldianus*.

On page 136, vol. II, *Trichocereus macrogonus*, add: *Illustrations*: Garten-Zeitung 4^a: 182. f. 8, as *Cereus macrogonus*.

On page 140, vol. II, under *Trichocereus coquimbicus*, add to illustrations: Engler and Drude, Veg. Erde 8: pl. 16, as *Cereus nigripilis*.

On page 140, vol. II, under *Trichocereus terscheckii*, insert: This cactus is the only timber found in the region of the Puna and in the western mountains of Argentina that can be utilized in any form. It is employed on a large scale in the mines for timbering the galleries, if these happen to be dry. It is called cardón.

Add: *Illustration*: Sci. Amer. 124: 492.

On page 143, vol. II, under *Trichocereus candicans*, insert: The names *Cereus gladiatus vernaculatus* Monville (Labouret, Monogr. Cact. 327. 1853) and *C. gladiatus courantii* (Förster, Handb. Cact. ed. 2. 833. 1885) were given as synonyms of *C. candicans*.

Cereus candicans dumesnilianus is figured and briefly described in the Gardeners' Chronicle (III. 26: 415. f. 132). It is an upright plant with long, straight spines; the flowers are large and pure white. It flowered in the collection of Justus Corderoy.

On page 144, vol. II, under *Trichocereus schickendantzii*, add to illustration: Möllers Deutsche Gärt. Zeit. 25: 475. f. 7, No. 16, as *Echinopsis schickendantzii*.

On page 146, vol. II, under *Echinopsis catamarcensis*, add: Illustration: Möllers Deutsche Gärt. Zeit. 25: 475. f. 7, No. 19.

On page 149, vol. II, under *Harrisia eriophora*, add to illustration: Journ. N. Y. Bot. Gard. 11: 234. f. Roig. Cact. Fl. Cub. pl. [5], as *Harrisia undata*.

Insert: *Cactus peruvianus jamaicensis* appears in Grisebach's Flora (Fl. Brit. W. Ind. 301. 1860) as a synonym of *Cereus eriophorus*, but refers to *Harrisia gracilis*.

On page 151, vol. II, under *Harrisia nashii*, add: Illustration: Descourtilz, Fl. Med. Antill. 1: pl. 66, as *Cactus divaricatus*.


On page 151, vol. II, under *Harrisia gracilis*, add the synonym: *Cactus subrepandis* Sprengel, Syst. 2: 495. 1825.

Add to illustrations: Förster, Handb. Cact. ed. 2. f. 139; Blühende Kakteen 2: pl. 84; Watson, Cact. Cult. 85. f. 28; ed. 3. f. 19; Dict. Gard. Nicholson Suppl. 220. f. 255, as *Cereus repandus*; Addisonia 2: pl. 61.

On page 154, vol. II, under *Harrisia aboriginum*, add: Illustration: Journ. N. V. Bot. Gard. 22: pl. 253.

On page 155, vol. II, under *Harrisia martinii*, add to illustrations: Addisonia 2: pl. 68.

On page 157, vol. II, under *Harrisia bonplandii*, add to illustrations: Monatsschr. Kakteenk. 25: 3, as *Cereus bonplandii*.



FIGS. 252, 253, 254.—*Borzicactus fieldianus*.

On page 163, vol. II, insert:

9. *Borzicactus fieldianus* sp. nov.

Forming thickets 3 to 6 meters high, the branches elongated, at first erect or ascending but sometimes becoming pendent or even prostrate; ribs few, perhaps only 6 or 7, stout, broad, 1 to 2 cm. high, depressed between the areoles and on young shoots and appearing as tubercled; areoles large, circular, short-lanate and spiny, with a depression extending upward from its upper side to constriction of rib; spines 6 to 10, white, subulate, very unequal, the longest ones 5 cm. long or longer; flowers several, from near tip of branches, but with only one from an areole, with a cylindric tube 6 to 7 cm. long and a very narrow limb; ovary and flower-tube bearing ovate, acute scales, 1 to 3 mm. long, these with long brown hairs in their axils; flower-tube within glabrous below its throat, bearing many stamens 4 cm. long; perianth-segments red, 1 cm. long; stamens exerted only beyond the perianth-segments, if at all; ovary globular, perhaps somewhat tuberculate, with scattered, long-hairy areoles; fruit probably fleshy, globular to ovoid, 2 cm. in diameter.

Collected by Macbride and Featherstone on gravelly river bluffs, eastern exposure at Huaraz, Peru, altitude about 2,600 meters, October 6, 1922 (No. 2519).

This very interesting plant we have named in honor of Captain Marshall Field, a patron of science, who financed the Botanical Expedition of 1922 to South America, sent out by the Field Museum of Natural History.

Figure 250 shows the habit of the plant, 251 a flowering branch, and figures 252 to 254 show flower, rib, and fruit.

On page 164, vol. II, under *Carnegiea gigantea*, add to illustrations:* Remark, Kakteenfreund 19, as *Pilocereus giganteus*; Nat. Geogr. Mag. 41: 373, as giant cactus; Tribune Hort. 4: 243; Journ. N. Y. Bot. Gard. 3: f. 15, 16, 17; 5: 173. f. 27; 6: f. 31, 32; Gartenwelt 8: 485; 11: 498; Schelle, Handb. Kakteenk. f. 20, 21; Cact. Mex. Bound. frontispiece; Bull. U. S. Nat. Mus. 56: pl. 8, f. 1; Useful Wild Pl. U. S. Canada opp. 112; Gartenflora 54: 589. f. 70; Gard. Chron. II. 20: 265. f. 39; Rev. Hort. IV. 3: 343. f. 20; Wiener Ill. Gart. Zeit. 11: 216. f. 47; Watson, Cact. Cult. 76. f. 22; Balt. Cact. Journ. 1: 67; Blanc, Cacti 30. No. 120; Carnegie Institution of Washington 6: pl. 1; De Laet, Cat. Gén. f. 26; Monatsschr. Kakteenk. 32: 87, as *Cereus giganteus*; Contr. U. S. Nat. Herb. 16: pl. 7; Amer. Bot. 26: 136; Stand. Cycl. Hort. Bailey; 1: Pl. 3; 2: f. 819; Nat. Geogr. Mag. 44: 171.

On page 167, vol. II, for *Binghamia melanostele*, substitute for this name:

***Binghamia multangularis* (Willdenow).**

Cactus multangularis Willdenow, Enum. Pl. Suppl. 33. 1813.

Cereus multangularis Haworth, Suppl. Pl. Succ. 75. 1819.

Echinocereus multangularis Rümpler in Förster, Handb. Cact. ed. 2. 825. 1885.

Cephalocereus melanostele Vaupel, Bot. Jahrb. Engler 50: Beibl. 111: 12. 1913.

Binghamia melanostele Britton and Rose, Cactaceae 2: 167. 1921.

Insert: We have recently obtained a photograph of Haworth's plant bearing the date "Oct. 29, 1824." A careful comparison of this photograph with photographs and specimens obtained by Dr. Rose in Peru in 1914 convinces us that this is the same plant as *Cephalocereus melanostele* which we referred to *Binghamia*.

Figure 255 is from a photograph of Haworth's plant, from N. E. Brown of Kew.

On page 171, vol. II, under *Oreocereus celsianus*, add to illustrations: Karsten and Schenck, Vegetationsbilder 7: pl. 42; Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 11, as *Pilocereus celsianus*; Möllers Deutsche Gärt. Zeit. 25: 473. f. 5, No. 6, as *P. kranzleri*;† Watson, Cact. Cult. 146. f. 57, as *P. bruenowii*; Balt. Cact. Journ. 1: 133, as *P. fossulatus*; Monatsschr. Kakteenk. 31: 123; 32: 9; Gartenflora 62: f. 55, as *Cereus straussii*; Möllers Deutsche Gärt. Zeit. 25: 475. f. 5, No. 15, as *P. williamsii*; Amer. Mus. Journ. 16: 39; Bull. Pan Amer. Union 42: 408.

On page 174, vol. II, under *Cleistocactus baumannii*, add to illustrations: Deutsches Mag. Gart. Blumen. 1851: pl. opp. 48, as *Cereus tweediei*; Jard. Fleur. 1: pl. 48; De Laet, Cat. Gén. f. 25; Blanc, Cacti 24. f. 2; West Amer. Sci. 13: 8, as *Cereus colubrinus*.

On page 177 vol. II, under *Lophocereus schottii* add to illustrations Karsten and Schenck Vegetationsbilder 13: pl. 18, in part




FIG. 255.—*Binghamia multangularis*.

* Some of the illustrations cited here and on pages 166 and 167 do not have the technical name of the plants.

† This name is credited to Rümpler but he gives the spelling as *Pilocereus kanzleri*.

Contr. U. S. Nat. Herb. **16**: pl. 126, B, as *Lophocereus australis*; Möllers Deutsche Gart. Zeit. **25**: 473. f. 5, No. 8; Cycl. Amer. Hort. Bailey **3**: f. 1803; Schumann, Gesamtb. Kakteen f. 7, 8; Nachtr. f. 8; Monatsschr. Kakteenk. **11**: 10; **18**: 101, as *Pilocereus schottii*; West Amer. Sci. **13**: 16, as *Cereus sargentianus*; Rep. Mo. Bot. Gard. **16**: pl. 4, 5, 6, 7, 8, as *Cereus schottii*; Contr. U. S. Nat. Herb. **16**: pl. 125, B, as *Lophocereus schottii*; Thomas, Zimmerkultur Kakteen **17**, as *Pilocereus sargentianus*.


On page 180, vol. II, under *Myrtillocactus geometrizans*, add to illustrations: Reiche, Elem. Bot. **228**. f. 164; Engler and Drude, Veg. Erde **13**: f. 31; Möllers Deutsche Gart. Zeit. **25**: 482. f. 13; West Amer. Sci. **13**: 15; Zeitschrift Sukkulenk. **1**: 31, as *Cereus geometrizans*.

On page 180, vol. II, under *Myrtillocactus cochal*, add to illustrations: Thomas, Zimmerkultur Kakteen **13**, as *Cereus cochal*.

On page 183, vol. II, insert the following:

39. NEOABBOTTIA Britton and Rose, Smiths. Misc. Coil. **72**⁹: 2. 1921.

A tree-like cactus with a smooth, upright, terete trunk and a much branched top, the branches strongly winged or ribbed, normally from distal end of preceding branch, but sometimes from below tip and usually in the same plane; ribs thin and high, very spiny; flowers nocturnal, small, tubular, with a narrow limb, borne several together at distal end of a terminal branch from a small, felted cephalium; perianth persisting on the ovary; perianth-tube and ovary bearing small scales with short wool and an occasional bristle in their axils; perianth-segments very small; throat of flower a little broadened at top, bearing many stamens; style slender; fruit oblong, turgid, nearly naked, deeply umbilicate; seed minute, black, muricate.



FIGS. 256, 257.—Flower and fruit of *Neoabbottia paniculata*. Natural size.

Type species: *Cactus paniculatus* Lamarck.

A monotypic genus of Hispaniola, dedicated to Dr. W. L. Abbott, a patron of natural history.

1. Neoabbottia paniculata (Lamarck) Britton and Rose, Smiths. Misc. Coll. **72**⁹: 3. 1921.

Cactus paniculatus Lamarck, Encycl. **1**: 540. 1783.

Cereus paniculatus De Candolle, Prodr. **3**: 466. 1828.

Plant 6 to 10 meters high or higher; trunk woody, 30 cm. in diameter, the wood close-grained, yellowish white; bark of trunk 1.5 cm. thick, brown, not spiny in age, smooth; branches 4 to 6 cm. broad, strongly 4-ribbed, occasionally 6-ribbed or winged; ribs thin, 1.5 to 2.5 cm. high, their margins somewhat crenate, areoles borne at base of sinuses, 1.5 to 2 cm. apart; spines 12 to 20, acicular, brownish to gray, 2 cm. long or less; cephalium 1 to 1.5 cm. in diameter, becoming elongated and angled; flowers straight, cm. long, with a limb about 3 cm. broad, tube 6 to 7 mm. long, about 18 mm. in diameter, with walls 5 to 6 mm. thick; inner perianth-segments greenish white, short-oblong, about 1 cm. long, obtuse; throat 18 mm. long, covered with numerous filaments, these

with a knee near base and pressing against style; stamens and style included; ovary and flower-tube tubercled, the former with short tubercles, the latter with oblong ones (sometimes 1.5 cm. long), each ending in a depressed areole subtended by a minute scale; areoles bearing a tuft of brown felt and an occasional brown bristle; fruit oblong in outline, 6 to 7 cm. long, 4 to 5.5 cm. in diameter, turgid, nearly naked; rind green, thick, hard; seeds rounded above, cuneate at base, with a large lateral depressed hilum.



FIG. 258.—*Neoabbottia paniculata*.

Type locality: Haiti.

Distribution: Hispaniola.

This plant was described by Plumier as follows: "*Melocactus arborescens*, tetragonus, fore ex albedo." This description was repeated by Tournefort, with the addition of a single word, in 1719. Plumier's drawing of this plant was published long after his death by Burmann as plate 192 of the *Plantarum Americanum*, and upon this plate Lamarck based his *Cactus paniculatus*, which De Candolle a little later took up as *Cereus paniculatus*. Ever since, the plant has usually passed under the latter name, with an occasional reversal to the earlier one.

Until recently, the species has been known only from this old illustration and these brief descriptions. It was collected near Port-au-Prince, Haiti, on the Cul-de-sac, by Dr. W. L. Abbott and Mr. E. C. Leonard, April 1920 (No. 3500); also at the same locality by Mr. H. M. Pilkington, December 1920; also a single branch by Dr. Paul Bartsch at Thomazeau in 1917 (No. 221). The Abbott and Leonard material consists of wood-sections and herbarium specimens of branches, flowers, fruit, and seeds, supplemented by living specimens and by fruit and flowers in formalin, together with several habit photographs.

In habit it resembles *Dendrocereus*, its branches resemble *Acanthocereus*, and the small limb of the flower resembles *Leptocereus*; but the plant differs from all of these in bearing several flowers at the ends of terminal branches and in developing a kind of cephalium. In the last respect it approaches *Neoraimondia*, near which we would place it in our present classification.

Illustrations: Smiths. Misc. Coll. 72^o: pl. 1 to 4; pl. 2, f. 1, 2; Bull. Amer. Mus. Nat. Hist. 33: 31. f. 11.

Figures 256 and 257 show the flower and fruit; figure 258 shows the top of a tree; figures 259 and 260 show the plant in its natural surroundings; figure 223*a*, page 248, is a reproduction of Plumier's plate.

On page 187, vol. II, under *Hylocereus undatus*, add to illustrations: De Laet, Cat. Gen. f. 31; Tribune Hort. 4: pl. 140; Blanc, Cacti 37. No. 346; Ann. Inst. Roy. Hort.

Fromont 2 pl. 1, f. D; Gartenwelt 11: 101; Watson, Cact. Cult. ed. 3. pl. opp. 29; Rev. Hort. Belg. 40: after 184; Meehans' Monthly 6: 5; West Amer. Sci. 13: 5; Gartenflora 55: f. 2, as *Cereus triangularis*; De Tussac, Fl. Antill. 4: pl. 26, as *Cactus triangularis*; Stand. Cycl. Hort. Bailey 3: pl. 57, as *Hylocereus tricostatus*; Cañizares, Jard. Bot. Inst. Habana 98, as *H. triangularis*.

On page 189, vol. 11, under *Hylocereus lemairei*, add to illustrations: Blühende Kakteen 3: pl. 173, as *Cereus lemairei*.

On page 191, vol. 11, under *Hylocereus napoleonis*, add to illustrations: Hartinger, Parad. 2: 1, as *Cereus napoleonis*.



Figs. 259 and 260.—*Neoabbotia paniculata*.

On page 192, vol. 11, under *Hylocereus triangularis*, insert: The name *Cactus anizogonus* of English gardens is given as a synonym of *Cereus triangularis* by Rümpler (Förster, Handb. Cact. ed. 2. 764. 1885).

On page 192, vol. 11, under *Hylocereus trigonus* insert: *Cereus triqueter* Haworth (Syn. Pl. Succ. 181. 1812) is some species of *Hylocereus* near *H. trigonus*. If really from South America, as stated by Haworth, it may be the same as *H. lemairei*.

On page 194, vol. 11, under *Hylocereus* sp., insert after first paragraph: This species of *Hylocereus* from the Guianas should be studied in connection with *Cereus scandens* Salm-Dyck (Cact. Hort. Dyck. 1849. 219. 1850), which is said to have come from Guiana. The variety *C. scandens minor* Boerhaave (Monatsschr. Kakteenk. 1: 82. 1891) is only mentioned.

After page proof had been read, some fine specimens of a *Hylocereus* were received from Surinam through Gerold Stahel, which we describe as follows:

Stems much elongated, 3-angled, 2 to 6 cm. broad, bluish or whitened, somewhat glaucous; ribs often thin; margins of ribs not horny, nearly straight, areoles distant, sometimes 6 cm. apart; spines brown, 2 or 3, very short, much swollen at base.

In the shape, number, and size of spines this specimen resembles *H. lemairei*, but differs from it in the whitened stems. We do not know its flowers.

On page 197, vol. II, under *Selenicereus grandiflorus*, insert: *Cereus haitiensis* Hortus is cited by Schelle (Handb. Kakteenk. 89. 1907) as a synonym of *C. grandiflorus*.

Cereus grandiflorus flemingii Rümpler (Förster, Handb. Cact. ed. 2. 751. 1885; *C. flemingii*, Monatsschr. Kakteenk. 3: 109. 1893) is said to be a hybrid between *C. grandiflorus* and *C. speciosissimus*.

Add to illustrations: Fl. Serr. 3: pl. 1-2, as *Cereus grandifloro-speciosissimus*; Balt. Cact. Journ. 1: 56, as queen of the night; Remark, Kakteenfreund 8; Gartenflora 42: 541. f. 110; 64: 90. f. 22; Gartenwelt 16: 613; 19: 18; Gard. Chron. III. 14: 187. f. 36; Thomas, Zimmerkultur Kakteen 15; Tribune Hort. 4: pl. 139; Blanc, Cacti 32; De Laet, Cat. Gén. f. 29; Fl. Serr. 3: 233-234; Weinberg, Cacti 8; Knippel, Kakteen pl. 1; Goebel, Pflanz. Schild. 1: pl. 2, f. 5; Möllers Deutsche Gärt. Zeit. 14: 340 to 343; 20: 561, as *Cereus grandiflorus*; Cañizares, Jard. Bot. Inst. Habana 100.

On page 198, vol. II, under *Selenicereus urbanianus*, add to illustrations: Gartenwelt 12: 255, as *Cereus urbanianus*; Roig, Cact. Fl. Cub. pl. [3] f. 2; pl. [4], as *Selenicereus maxonii*.

On page 199, vol. II, under *Selenicereus coniflorus*, insert: Dr. J. K. Small finds this plant naturalized in pinelands near the Everglades, west of Halenville, Florida.

On page 200, vol. II, under *Selenicereus pteranthus*, also add to illustrations: Garden 13: 291; Monatsschr. Kakteenk. 31: 71; Watson, Cact. Cult. 63. f. 15; ed. 3. f. 10; Gartenflora 41: f. 23, 24, as *Cereus nycticalus*.

Add the synonym: *Cereus nycticalus peanii* Beguin in Riccoboni, Boll. R. Ort. Bot. Giard. Col. Palermo 8: 252. 1909.

On page 202, vol. II, under *Selenicereus boeckmannii*, add to illustration: Blühende Kakteen 3: pl. 175, 176, as *Cereus boeckmannii*.

On page 202, vol. II, under *Selenicereus macdonaldiae*, add the synonym: *Cereus grandiflorus macdonaldiae* Blanc, Cacti 34.

Also insert: *Cereus kewensis* Worsley (Journ. Roy. Hort. Soc. 39: 92. 1913) is said to be a "garden hybrid between *C. macdonaldiae* and probably *C. nycticalus*."

Also add to illustrations: Blanc, Cacti 34. No. 206, as *Cereus grandiflorus macdonaldiae*; Monatsschr. Kakteenk. 30: 107; Gartenwelt 16: 537; Möllers Deutsche Gärt. Zeit. 25: 488. f. 22, No. 6, as *Cereus macdonaldiae*, Blühende Kakteen 3: pl. 166, 167, as *Cereus grusonianus*.

Insert: *Cereus rothii* Weingart (Monatsschr. Kakteenk. 32: 146. 1922) is of this relationship. It is a new name for the plant from South America called *Cereus macdonaldiae* by Spegazzini; we have not seen it.

On page 204, vol. II, under *Selenicereus hamatus*, add to illustrations: Tribune Hort. 4: pl. 140; Floralia 42: 371, as *Cereus rostratus*.

On page 209, vol. II, insert the following:

17. *Selenicereus nelsonii* (Weingart).

Cereus nelsonii Weingart, Zeitschrift Sukkulantenkunde 1: 33. 1823.

A slender, much branched vine, 1 to 1.5 cm. in diameter, giving off occasional aerial roots; ribs 6 or 7, low, somewhat tubercled; areoles small, circular, about 1 cm. apart; spines about 12, acicular, white to yellowish, 5 to 7 mm. long; length of flower including ovary and closed perianth

about 20 cm.; outer perianth-segments linear, pointed, reddish brown, the inner perianth-segments narrowly lanceolate, 7 cm. long, 12 to 15 mm. broad, acute; filaments numerous, weak, white; style long and slender, exerted beyond the withering perianth; stigma-lobes slender, white, entire; scales on the ovary and flower-tube minute, 1 to 1.5 mm. long, reddish brown, bearing white felt and white bristles in their axils; fruit crowned by the withering perianth, globular, 2 to 2.5 cm. in diameter, reddish, bearing numerous, small, circular areoles, these with clusters of acicular spines sometimes 1 cm. long.

Type locality: Southern Mexico.

Distribution: Mexico, but range not known.

We have had this plant under observation since 1914 when cuttings were sent us by C. Z. Nelson, an enthusiastic grower of cacti at Galesburg, Illinois, who obtained it from southern Mexico from Dr. J. L. Slater. This plant made two flowers during the week of May 17, 1922; the fruit ripens very slowly and did not mature until October 10, 1922.

According to Wilhelm Weingart, the same species has long been grown by Frantz de Laet at Contich, Belgium, also from Mexican material.

Illustration: Zeitschrift Sukkulente. 1: 33, as *Cereus nelsonii*.

Figure 261 shows a branch bearing a newly matured fruit.




FIG. 261.—*Selenicereus nelsonii*.

On page 216, vol. II, under *Werckleocereus tonduzii*, add: *Illustration:* Monatsschr. Kakteenk. 31: 85, as *Cereus tonduzii*.

On page 218, vol. II, under *Aporocactus leptophis*, add to illustrations: Ann. Fl. Pom. 1839: pl. 43, as *Cereus leptophis*.

On page 219, vol. II, under *Aporocactus flagelliformis*, add to illustrations: Tribune Hort. 1: pl. 4, as *Cereus serpentinus*; Fl. Antill. 1: pl. 67, as *cierge queue de souris*; Cact. Journ. 1: 82; Rother, Praktischer Leitfaden Kakteen 57; Remark, Kakteenfreund 6; Watson, Cact. Cult. ed. 3. f. 11; Floralia 42: 371; Gartenwelt 15: 637; Blanc, Cacti 27. No. 104; Amer. Gard. 11: 527 as *Cereus flagelliformis*; Cact. Journ. 1: 125; 2: 34; 2: 153, as *C. flagelliformis cristatus*.

Insert: *Cereus smithianus* is a hybrid listed by Sweet (Hort. Brit. ed. 2. 237. 1830) which the Index Kewensis refers to *C. smithii*, a generic hybrid already referred to.

On page 221, vol. II, under *Aporocactus martianus*, add to the illustrations: Thomas, Zimmerkult. Kakteen 14, as *Cereus martianus*.

CORRECTIONS AND ADDITIONS TO VOLUME III.

On page 6, vol. III, under *Echinocereus scheeri*, add to illustrations: Thomas, Zimmerkultur Kakteen 27.

On page 7, vol. III, under *Echinocereus salm-dyckianus*, add to illustrations: Thomas, Zimmerkultur Kakteen 25.

On page 11, vol. III, under *Echinocereus polyacanthus*, add to illustrations: Förster, Handb. Cact. ed. 2. 212. f. 19, as *Cereus polyacanthus*; Florialia 42: 376, as *Echinocereus polyacanthus* var.; Thomas, Zimmerkultur Kakteen 29; De Laet, Cat. Gén. f. 34, 37, 38; Möllers Deutsche Gärt. Zeit. 36: 145. f. 111; Succulenta 5: 74.

On page 12, vol. III, under *Echinocereus acifer*, add to illustrations: Thomas, Zimmerkultur Kakteen 26, as *Echinocereus acifer trichacanthus*; Blühende Kakteen 3: pl. 179, as *E. durangensis*.

On page 14, vol. III, under *Echinocereus coccineus*, add to illustrations: Pac. R. Rep. 4: pl. 4, f. 1 to 3, as *Cereus phoeniceus*.

On page 17, vol. III, under *Echinocereus viridiflorus*, add the synonym: *Cereus viridiflorus minor* Engelm., Proc. Amer. Acad. 3: 278. 1856.

On page 21, vol. III, under *Echinocereus blanckii*, add to illustrations: Watson, Cact. Cult. 70. f. 18, as *Cereus blankii*; Watson, Cact. Cult. 68. f. 17; ed. 3. f. 13, as *C. berlandieri*.

On page 22, vol. III, under *Echinocereus pentalophus*, add to illustrations: Watson, Cact. Cult. 78. f. 23, as *Cereus leptacanthus*; Watson, Cact. Cult. 83. f. 27; ed. 3. f. 18, as *C. procumbens*; Balt. Cact. Journ. 2: 218, as *Echinocereus procumbens*.

Also add the synonym: *Cereus propinquus subarticulatus* Pfeiffer in Förster, Handb. Cact. 373. 1846.

On page 23, vol. III, under *Echinocereus cinerascens*, insert: *Echinocactus deppii* Link and Otto (Steudel, Nom. ed. 2. 1: 536. 1840) was given in error for *Echinocereus deppii*.

On page 25, vol. III, under *Echinocereus reichenbachii*, insert: Watson, Cact. Cult. ed. 3. f. 14, as *Cereus caespitosus*; West Amer. Sci. 7: 237; 13: 14; Gartenflora 23: pl. 813, as *C. pectinatus*; Remark, Kakteenfreund 17; Balt. Cact. Journ. 2: 218, as *Echinocereus caespitosus*.

On page 37, vol. III, under *Echinocereus enneacanthus*, add to illustrations: Bull. Univ. Tex. 60: pl. 11, f. 1, as *Cereus longispinus*; Watson, Cact. Cult. 75. f. 21; ed. 3. f. 15, as *C. enneacanthus*.

Also insert: *Echinocereus saltillensis* is offered for sale by Haage and Schmidt, 1920, page 75.

Mr. C. R. Orcutt has called our attention to the following varieties which have been omitted: *Cereus engelmannii* var. *albispinus* Cels, var. *caespitosus*, var. *fulvispinus* Cels, var. *pfersdorffii* Heiden, all of which are listed by him (Orcutt, Rev. Cact. 1: 13. 1897),

On page 45, vol. III, insert: The name *Cactus bertini* was given for this plant when awarded a silver medal soon after its discovery (Hort. Franc. 11. 5: 222).

On page 45, vol. III, *Rebutia minuscula*, add to illustrations: Succulenta 3: 96; Thomas, Zimmerkultur Kakteen 34; Kaktusy 25, as *Echinocactus minusculus*.

On page 48, vol. III, under *Chamaecereus silvestrii*, add to illustrations: Blühende Kakteen 3: pl. 168, as *Cereus silvestrii*.

On page 48, vol. III, substitute for *Echinopsis deminuta*:

5. *Rebutia deminuta* (Weber).

Echinopsis deminuta Weber, Bull. Mus. Hist. Nat. Paris 10: 386. 1904.

Through the kindness of J. J. Verbeek Wolthuys, we have been able to examine a flower of this plant which shows that it belongs to the genus *Rebutia*.

On page 54, vol. III, under *Lobivia pentlandii*, add to illustrations: Watson, Cact. Cult. ed. 3. f. 32, as *Echinopsis pentlandii*.

On page 59, vol. III, insert the following:

21. *Lobivia famatimensis* (Spegazzini).

Echinocactus famatimensis Spegazzini, Anal. Soc. Cient. Argentina 92: 44. 1921.

Solitary or in clusters, short-cylindric, 3 to 3.5 cm. high, 2.5 to 2.8 cm. in diameter, strongly umbilicate at apex; ribs 24, low, obtuse, somewhat tuberculate; areoles approximate; spines small, appressed, whitish; flowers solitary, from the side near the middle, about 3 cm. long.

Type locality: Near Famatima, Argentina, altitude 2,000 to 3,000 meters.

Distribution: Province of La Rioja, Argentina.

Illustration: Anal. Soc. Cient. Argentina 92: f. , as *Echinocactus famatimensis*.

On page 64, vol. ii, under *Echinopsis multiplex*, add to illustrations: Watson, Cact. Cult. ed. 3. f. 16, as *Cereus multiplex*; Rev. Hort. 48: 13. f. 1, as *Echinocactus multiplex*; Rev. Hort. 48: 13. f. 2, as *E. multiplex cristata*; Gard. Chron. III. 56: 145. f. 60.

On page 65, vol. III, under *Echinopsis oxygona*, add to illustrations: Thomas, Zimmerkultur Kakteen 23, as *Echinopsis oxygona inermis*; Succulenta 5: 85.

On page 66, vol. III, under *Echinopsis eyriesii*, add to illustrations: Rother, Praktischer Leitfaden Kakteen 47, as *Echinopsis triumphans*; Remark, Kakteenfreund 9; Rother, Praktischer Leitfaden Kakteen 45, 106.

On page 67, vol. III, under *Echinopsis turbinata*, add to illustrations: Watson, Cact. Cult. 131. f. 50; ed. 3. f. 30; Gard. Chron. III. 16: 625. f. 79, as *Echinopsis decaisneana*; Floralia 42: 374, as *E. gemmata*.

On page 67, vol. III, under *Echinopsis tubiflora*, add to illustrations: Thomas, Zimmerkultur Kakteen 20, as *Echinopsis tubiflora rohlandii*; Rother, Praktischer Leitfaden Kakteen 44, as *E. zuccariniana*.

On page 72, vol. III, under *Echinopsis leucantha*, add to illustrations: Thomas, Zimmerkultur Kakteen 21.

On page 74, vol. III, under *Echinopsis bridgesii*, insert: *Illustration:* Möllers Deutsche Gärt. Zeit. 25: 475. f. 7, No. 18, as *Echinopsis salmiana*.

On page 75, vol. III, under *Echinopsis formosa*, add to illustrations: Monatsschr. Kakteenk. 32: 149.

On page 76, vol. III, *Echinopsis formosissima*, insert: *Cereus formosissimus* Weber (Dict. Hort. Bois 471. 1896) was cited by Weber as a synonym of this species.

On page 80, vol. III, under *Ariocarpus retusus*, add the synonyms: *Cactus areolus* Kuntze, Rev. Gen. Pl. 1: 260. 1891; *Cactus pulvilliger* Kuntze, Rev. Gen. Pl. 1: 260. 1891.

Add to illustrations: Monatsschr. Kakteenk. 23: 66, 67, as *Ariocarpus trigonus*.

On page 82, vol. III, under *Ariocarpus kotschoubeyanus* insert: *Anhalonium kotschubeyi* Lemaire (Salm-Dyck, Cact. Hort. Dyck. 1849. 5. 1850), given as a synonym of *A. sulcatum*, is to be referred here.

On page 83, vol. III, under *Ariocarpus fissuratus*, add to illustrations: Bull Univ. Texas 60: pl. 11, f. 2, as *Ariocarpus fissuratus*; Gard. Chron. III. 12: 789. f. 130; Watson, Cact. Cult. 161. f. 61, as *Mammillaria fissurata*; Watson, Cact. Cult. ed. 3. f. 6, as *Anhalonium engelmannii*; Remark, Kakteenfreund 10; Bait. Cact. Journ. 1: 27; 2: 247, as *Anhalonium fissuratum*; Rother, Praktischer Leitfaden Kakteen 35.

On page 85, vol. III, under *Lophophora williamsii*, add to illustrations: Sci. Amer. 124: 492, as mescal button; Rother, Praktischer Leitfaden Kakteen 36; Remark, Kakteenfreund 11; Karsten and Schenck, Vegetationsbilder 2: pl. 20, B; Thomas, Zimmerkultur Kakteen 30; Monatsschr. Kakteenk. 31: 187, as *Echinocactus williamsii*; Balt. Cact. Journ. 1: 71; 2: 247; Watson, Cact. Cult. ed. 2. 243. f. 91; ed. 3. f. 7, as *Anhalonium williamsii*; Succulenta 2: 3; 4: 7.

On page 91, vol. III, under *Pediocactus simpsonii*, add to illustrations: Wiener Obst. Zeit. 2: 90. f. 13; Remark, Kakteenfreund 13, as *Echinocactus simpsonii*.

On page 106, vol. III, under *Hamatocactus setispinus*, add to illustrations: Schulz, 500 Wild Fl. San Antonio pl. 12, as *Echinocactus setispinus*.

On page 107, vol. III, under *Strombocactus disciformis*, add to illustrations: Remark, Kakteenfreund 12, as *Echinocactus turbiniformis*.

On page 123, vol. III, at end of *Echinofossulocactus*, insert the following:

Echinocactus tetracentrus Lemaire (Cact. Gen. Nov. Sp. 31. 1839) has not been identified. It is to be referred to one of the species of *Echinofossulocactus*.

Echinocactus barcelona (Cact. Journ. 2: 79, 175, 191) was offered for sale by F. A. Walton.

On page 124, vol. III, under *Ferocactus stainesii*, insert: *Echinocactus pilosus canescens* Scheidweiler is listed in Index Bibliographique (286. 1887).

On page 129, vol. III, under *Ferocactus lecontei*, insert: *Echinocactus leopoldii* (Belg. Hort. 25: 132. 1876) was only briefly described when awarded a prize in Belgium. Schumann referred it as a synonym of *E. cylindraceus*, while De Laet, who saw the plant, thought it was a form of *Ferocactus lecontei*. It is misspelled in Volume III.

Add to illustrations: Wiener Ill. Gart. Zeit. 11: pl. 3, in part; Thomas, Zimmerkultur Kakteen 32; Watson, Cact. Cult. ed. 3. f. 26, as *Echinocactus lecontei*.

On page 130, vol. III, under *Ferocactus acanthodes*, add to illustrations: Thomas, Zimmerkultur Kakteen 40, as *Echinocactus cylindraceus*.

On page 132, vol. III, insert:

11a. *Ferocactus johnstonianus* Britton and Rose, sp. nov.

Plants simple, short-cylindric, 6 dm. high or less, up to 3.5 dm. in diameter; ribs 24 to 31, with margins undulate; areoles elliptic, rather closely set; spines 20 or more, subulate, very much alike, none hooked, slightly spreading and more or less outwardly recurved, 7 cm. long or less, yellow to brownish yellow, annulate; flowers including ovary 5 cm. long; perianth-segments narrow, yellowish, or the outer ones tinged with red, short-acuminate, the margins slightly erose; filaments yellowish below, becoming reddish above; stigma-lobes 8 to 13, flesh-colored; scales on the ovary orbicular; fruit small, 2.5 cm. in diameter, the seed dehiscing by a large pore at the base; seeds angled, black, pitted, 2 mm. long; hilum small, circular, depressed, white.

Collected by Ivan M. Johnston at Angel de la Guardia Island, Lower California May 2, 1921 (Nos. 3394, type, and 3395).

This species is perhaps nearest *Ferocactus diguetii* but is much smaller and has fewer ribs, many more spines in a cluster, and yellow flowers.

On page 140, vol. III, under *Ferocactus viridescens*, add to illustrations: Blühende Kakteen 3: pl. 177, as *Echinocactus viridescens*.

On page 143, vol. III, under *Ferocactus latispinus*, add to illustrations: Remark, Kakteenfreund 13, as *Echinocactus cornigerus flavispinus*.

Also add the note: *Cactus cornigereus* Mociño and Sessé (De Candolle, Prodr. 3: 461. 1828) is given as a synonym of *Echinocactus cornigerus*.

On page 144, vol. III, under *Ferocactus hamatacanthus*, add to illustrations: Rother, Praktischer Leitfaden Kakteen 106, as *Echinocactus longihamatus*.

On page 148, vol. III, under *Echinomastus erectocentrus*, insert: Mr. C. R. Orcutt has called our attention to the fact that in *Echinomastus erectocentrus* the fruit opens by splitting down one side, and in this respect differs from our generic description. This character in the description was drawn from a study of the fruits of *E. intertextus*, the only species in this genus of which we know much about the fruit. He also states that the fruit of *Astrophytum myriostigma* splits open on one side, an observation we had not recorded.

On page 150, vol. III, under *Echinomastus unguispinus*, add to illustrations: Thomas, Zimmerkultur Kakteen 31; Schelle, Handb. Kakteenk. 200. f. 132, as *Echinocactus unguispinus*.

On page 155, vol. III, under *Gymnocalycium denudatum*, insert: *Echinocactus denudatus multiflorus* (Monatsschr. Kakteenk. 14: 178) is only a name.

Also add to illustrations: Thomas, Zimmerkultur Kakteen 41, as *Echinocactus denudatus*.

On page 157, vol. III, under *Gymnocalycium saglione*, add to illustrations: Thomas, Zimmerkultur 37, as *Echinocactus saglionis*.

On page 158, vol. III, under *Gymnocalycium gibbosum* add to illustrations: Van Géel, Sert. Bot. 1: 113, as *Cactus gibbosus*.

On page 159, vol. III, under *Gymnocalycium brachyanthum*, insert: *Illustration*: Möllers Deutsche Gärt. Zeit. 36: 145. No. 11, as *Echinocactus brachyanthus*.

On page 161, vol. III, under *Gymnocalycium monvillei*, add to illustrations: Möllers Deutsche Gärt. Zeit. 36: 145. f. 1, as *Echinocactus monvillei*.

On page 168, vol. III, under *Echinocactus grusonii*, add to illustrations: Watson, Cact. Cult. ed. 3. f. 23; Rother, Praktischer. Leitfaden Kakteen 30; Deutsche Garten-Zeitung 28. f. 6; Zeitschrift Sukkulentk. 1: 15.

On page 168, vol. III, under *Echinocactus ingens*, add to illustrations: Remark, Kakteenfreund 14.

On page 171, vol. III, under *Echinocactus visnaga*, add to illustrations: Balt. Cact. Journ. 2: 181.

On page 181, vol. III, insert the following paragraphs:

Echinocactus acutispinus Hildmann (Deutsche Garten-Zeitung 1886: 116. f. 27. 1886) was described and figured, but the plant is a small, barren one which we have not been able to associate with any described species. It came from Mexico and may be one of the species of *Echinocactus*.

Echinocactus cylindricus Hortus (Forbes, Hort. Tour Germ. 152) was described as cylindrical, with 12 or 13 ribs, the radial spines white and the central ones light brown. It was introduced from Mexico in 1836. It can not be identified.

On page 182, vol. III, under *Homalocephala texensis*, add to illustrations: Schulz, 500 Wild Flowers of San Antonio, pl. 13 in part as *Echinocactus texensis*.

Echinocactus darrahii Schumann (Monatsschr. Kakteenk. 12: 21. 1902) is only mentioned.

Echinocactus dicracanthus Hortus (Forbes, Journ. Hort. Tour Germ. 160) is only a name.

Echinocactus inflatus Gillies (Steudel, Nom. ed. 2. 1: 536. 1840) seems never to have been published. Steudel simply states that it was from Chile.

Echinocactus praegnacanthus Förster (Handb. Gartenz. 17: 160. 1861) is a plant from Chile which has never been identified.

Echinocactus purpureus (Monatsschr. Kakteenk. 5: 106. 1895) is listed by Schumann as in Gruson's Garden.

Echinocactus rhodanthus (Forbes, Journ. Hort. Tour Germ. 151) is only a name.

On page 182, vol. III, under *Astrophytum myriostigma*, add to illustrations: Remark, Kakteenfreund 11; Balt. Cact. Journ. 1: 82.

On page 185, vol. III, under *Astrophytum capricorne*, add to illustrations: Remark, Kakteenfreund 12, as *Echinocactus capricornis*.

On page 188, vol. III, under *Malacocarpus tephracanthus*, add the synonym: *Echinocactus sellowii tetracanthus* Lemaire in Schumann, Monatsschr. Kakteenk. 18: 150. 1908.

Insert: *Echinocactus buchheimianus* Haage in Quehl (Monatsschr. Kakteenk. 9: 74. 1899) has been described briefly but its flower and fruit are unknown. It is said to resemble *E. sellowii*.

Add to illustrations: Schumann, Gesamtb. Kakteen f. 15, as *Cereus tephraacanthus*.

On page 193, vol. III, under *Malacocarpus concinnus*, add to illustrations: Succulenta 3: 22, 48, as *Echinocactus concinnus*.

On page 193 vol. III, under *Malacocarpus scopia*, add to illustrations: Rother, Praktischer Leitfaden Kakteen 107, as *Echinocactus scopia cristatus*; Kaktusy 26, as *Echinocactus scopia*; Kaktusy 27, as *E. scopia cristata*; Kaktusy 28, as *E. scopia rubra*.

On page 195 vol. III, under *Malacocarpus linkii*, insert: *Echinocactus ottonis linkii* Hortus (Förster, Handb. Cact. ed. 2. 554. 1885) is given as a synonym of *E. linkii*.

On page 196, vol. III, under *Malacocarpus ottonis*, add to illustrations: Succulenta 3: 56, as *Echinocactus ottonis tenuispinus*; Rother, Praktischer Leitfaden Kakteen 34, as *E. ottonis*.

On page 198, vol. III, under *Malacocarpus erinaceus*, add to illustrations: Watson, Cact. Cult. 98. f. 32, as *Echinocactus corynodes*.

On page 200, vol. III, under *Malacocarpus mammulosus*, add to illustrations: Rother, Praktischer Leitfaden Kakteen 30, as *Echinocactus submammulosus*.

On page 202, vol. III, under *Malacocarpus haselbergii*, add to illustrations: Succulenta 3: 31, as *Echinocactus haselbergii*.

On page 205, vol. III, under *Malacocarpus leninghausii*, add to illustrations: Succulenta 3: 39, as *Echinocactus leninghausii*.

On page 207, vol. III, under *Hickenia microsperma*, add to illustrations: Succulenta 3: 71; 5: pl. 1; Thomas, Zimmerkultur Kakteen 35, as *Echinocactus microspermus*.

On page 237, vol. III, insert the following:

19. *Cactus oaxacensis* sp. nov.

Globular to ovoid, 12 to 15 cm. thick, with a small, low crown only 2 to 3 cm. high and 3 to 4 cm. broad; ribs 11 to 15, prominent, usually rounded; radial spines 8 to 12, subulate, more or less recurved at first, reddish brown but grayish in age, 2 cm. long or less; central spines 1 or sometimes 2, erect or porrect; flowers slender, about 2 cm. long, dark rose; filaments and style light yellow; fruit thick-clavate, 2 to 4.5 cm. long, scarlet, shiny; seeds small, black.

This plant was illustrated and mentioned in the place here cited (Cactaceae 3: 237. f. 249) but was not given a specific name. Since then C. R. Orcutt reports finding it at Salina Cruz and Dr. B. P. Reko sends us a photograph and flowers obtained by him in 1923, while Dr. J. A. Purpus re-collected it in 1923 (type) and has sent us living plants.

Illustration: Cactaceae 3: 236. f. 249, as *Cactus* sp.

Figure 262 is from a photograph of the plant sent us by Dr. Reko.

On page 238, vol. III, insert: *Melocactus ellemeetii* Miquel (Nederl. Kruidk. Arch. 4: 336. 1858) and *M. pachycentrus* Suringar (Verh. Akad. Wettensch. Amst. II. 8: 28. 1901) have not been identified.

On page 238, vol. III, at end of *Cactus* add: *Cactus aculeatissimus* is listed by Steudel (Nom. 131. 1821) credited to Zeyher and cited by the Index Kewensis, but it has never been identified.

Cactus tuna major is used by Roxburgh (Hort. Beng. 37. 1814).




Fig. 262.—*Cactus oaxacensis*.

Cactus reptans Willdenow (Ann. Hort. Berol. Suppl. 33. 1813) was taken up in this work by mistake as *Cereus reptans*.

Cactus neglectus Dehnhardt (Rivist. Napol. 1. 3: 166.), according to the Index Kewensis, is a species of *Pereskia*.

Cereus erinaceus is credited to Haworth by Steudel (Nom. ed. 2. 1: 334. 1840) and said to come from the West Indies. Steudel must have had in mind *Cactus erinaceus* Haworth; if so, the plant is from South America.

Cereus torrellianus (Monatsschr. Kakteenk. 20: 42. 1910) is probably a misspelling for *C. tonelianus*.

Figure 263, shown below, gives a typical Arizona landscape in which *Carnegiea gigantea* is the dominant plant.



INDEX.

This index covers the four volumes. The large capitals occurring before the figures indicate the particular volume. Pages of principal entries are in heavy-faced type.

- Aaron's beard, I, 175
Acanthocereus, 11, 1, 2, 15, 121-126, 207, 281
albicaulis, II, 122, 125, 126
brasiliensis, II, 122, 125, 126
colombianus, II, 122
floridanus, IV, 276
horridus, II, 122, 123
occidentalis, II, 122, 125
pentagonus, II, 122, 123, 124, 191; IV, 276
subinermis, II, 122, 125
Acanthorhopsalis, IV, 208, 211-213
crenata, IV, 211, 212
micrantha, IV, 211
monacantha, IV, 211, 212
Acutangles, II, 122
Agave, I, 117
Agave cactus, III, 108
Agave lophantha, III, 109
Agua-colla, II, 135
Ahoplocarpus, I, 11
Airampo, I, 161
Alati, IV, 186
Alfilerillo, I, 26
Algae, IV, 65
Alicocha, III, 40
Ammophilae, I, 45, 211; IV, 265
Ancistracanthi, III, 45
Ancistrocactus, IV, 3-5
brevihatus, IV, 4, 5
megarhizus, IV, 4
scheeri, IV, 4, 5
Angulosae-tetragonae, IV, 168
Anhalonium, III, 80, 83
abides pulvilligerum, III, 81
areolusum, III, 80
aselliforme, IV, 59
elongatum, III, 80, 81
engelmannii, III, 83; IV, 286
fissipedum, III, 82
furfuratum, III, 83; IV, 286
furfuraceum, III, 80
heteromorphum, III, 83
jordanianum, III, 85
kotschoubeyanum, III, 82
kotschubeyanum, III, 82
kotschubeyi, IV, 286
lewinii, III, 84, 85
prismaticum, III, 80, 81
pulvilligerum, III, 80, 81
retusum, III, 80
rungei, III, 84
subnodosum, III, 84
sulcatum, III, 82; IV, 286
trigonum, III, 80, 81
turbiniiforme, III, 106
visnaga, III, 84
williamsii, 111, 84, 85; IV, 286
Anomali, II, 22
Aporocactus, II, 183, 217-221
baumannii, II, 174
colubrinus, II, 174
conzattii, II, 217, 220, 221
flagelliformis, II, 217, 218, 219, 221; IV, 284
flagriformis, II, 217, 218, 219
leptophis, II, 217, 218; IV, 284
martianus, II, 217, 220, 221; IV, 284
Apple, I, 9
Arequipa, III, 77, 100, 101
leucotricha, III, 101
myriacantha, III, 101
Aria, III, 80
Ariocarpus, III, 77, 78, 80-83, 84, 93
arelliformis, IV, 59
Ariocarpus—continued,
fissuratus, III, 80, 81, 83; IV, 23, 286
furfuraceus, III, 80, 81, 184
kotschoubeyanus, III, 80, 82; IV, 286
kotschubeyanus, III, 82
lloydii, III, 83
mcdowellii, III, 82
prismaticus, III, 80
pulvilligerus, III, 80
retusus, III, 80, 81; IV, 286
sulcatus, III, 82
trigonus, III, 80, 81; IV, 286
williamsii, III, 84
Arrojadoa, II, 2, 170, 171, 178
penicillata, II, 170, 171
rhodantha, II, 170
Astrophytum, III, 78, 167, 182-185
asterias, III, 182, 183, 184
capricorne, III, 182, 184; IV, 288
glabrescens, III, 185
myriostigma, III, 182, 183, 185; IV, 287, 288
ornatum, III, 182, 185
prismaticum, III, 182
Aulacothele sulcolanatum, IV, 38
Aurantiacae, I, 45, 74, 106
Austrocactus, III, 3, 44, 45
bertinii, III, 44
Ayrampo, I, 135
Azurae, II, 4
Bande du sud, II, 151
Barbados gooseberry, I, 10
Barrel cacti, III, 131
Bartschella, IV, 3, 57-59
schumannii, IV, 58
Basilares, I, 45, 118, 193; IV, 262
Bauhinia, II, 92
Bavoso, II, 86
Beaucarnea, I, 117
Beaver-tail, I, 120
Bergerocactus, II, 2, 107, 108; III, 4
emoryi, II, 107, 108; IV, 274
Bergerocereus, II, 108
Bigelovianae, I, 44, 58
Binghamia, II, 2, 167-169; IV, 279
acrantha, II, 167, 168
melanostele, II, 167, 168; IV, 279
multangularis, IV, 279
Bisnaga, III, 134
Blade apple, I, 10
Borzacactus, II, 2, 159-164, 173; III, 103
acanthurus, II, 159, 161
aurivillus, II, 159, 163, 226
decumbens, II, 159, 162
fieldianus, IV, 277, 278
humboldtii, II, 159, 163
icosagonus, II, 159, 160
morleyanus, II, 159, 160, 161
plagiostoma, II, 159, 163
sepium, II, 159, 160
ventimigliac, II, 159, 160
Bottle cactus, II, 60
Brachycereus, II, 2, 120, 121
thouarsii, II, 120, 121
Brain cactus, III, 90
Brasilienses, I, 45, 209
Brasilopuntia, I, 209
Breebee, II, 18
Bromeliads, II, 42; IV, 183
Browningia, II, 2, 63, 64
candelaris, II, 63, 64
Bullsucker, I, 43, 116
Button cactus, III, 93
Cabeça branca, II, 30
Cabeça de frade, III, 221, 235
Cabeza del viego, III, 28
Cactaceae, I, 3, 8; II, 1, 211; IV, 3
Cactales, I, 8
Cactanae, II, 1; III, 216; IV, 273
Cacti, I, 5, 6, 7, 9, 28, 33, 39, 42, 49, 66, 67, 80, 87, 94, 95, 111, 126, 137, 151, 216, 219, 220; II, 1, 2, 3, 8, 9, 19, 21, 23, 25, 28, 32, 65, 67, 77, 82, 113, 116, 117, 121, 122, 124, 127, 145, 147, 159, 165, 166, 167, 169, 173, 176, 177, 178, 182, 183, 185, 189, 196, 212, 214, 216, 217, 224; III, 3, 4, 6, 32, 33, 42, 46, 61, 63, 64, 70, 90, 91, 100, 108, 109, 123, 166, 167, 172, 179, 202, 207, 212, 213, 221, 224, 225, 226, 236; IV, 3, 6, 14, 19, 33, 48, 51, 53, 58, 64, 70, 71, 100, 107, 118, 134, 145, 155, 177, 183, 185, 186, 191, 205, 206, 208, 211, 213, 215, 216, 219, 220, 249, 250, 253, 256, 275, 284
Cactier rouge, III, 232
Cactier rouge main, III, 226
Cactodendron, I, 42, 43
Cactus, I, 5, 6, 8, 9, 23, 30, 32, 34, 35, 43, 49, 87, 88, 93, 107, 113, 114, 116, 120, 121, 146, 164, 177, 186, 207, 210, 215, 216, 217, 221; II, 1, 5, 9, 19, 23, 25, 40, 42, 58, 61, 62, 64, 65, 70, 71, 72, 76, 87, 88, 90, 92, 102, 107, 111, 113, 115, 124, 125, 135, 140, 147, 150, 151, 153, 158, 159, 163, 164, 165, 166, 167, 170, 172, 179, 181, 191, 192, 195, 210, 212, 221, 222, 224; III, 3, 6, 7, 23, 29, 31, 32, 44, 45, 54, 60, 78, 90, 91, 104, 106, 107, 109, 123, 142, 155, 166, 170, 173, 175, 182, 186, 193, 208, 212, 213, 215, 216, 220-238; IV, 3, 4, 6, 22, 24, 35, 53, 57, 60, 71, 79, 95, 98, 141, 156, 162, 163, 172, 177, 186, 201, 203, 204, 249, 250, 253, 261, 277, 280, 289
abnormis, II, 12
acanthophlegma, IV, 107
acanthostephes, IV, 41
acicularis, IV, 171
aciculatus, IV, 132
ackermannii, IV, 198
actinopleus, IV, 172
aculeatissimus, III, 238; IV, 289
aeruginosus, IV, 88
affinis, IV, 88
alatus, IV, 205, 213, 243
albisetosus, IV, 270
albisetus, IV, 270
alpinus, IV, 165
alteolens, III, 218
alternatus, IV, 79
amabilis, IV, 172
ambiguus, II, 118, 119; III, 98; IV, 75
americanus, IV, 273
Cactus—continued,
amoenus, III, 222, 232
ancistracanthus, IV, 15
ancistrius, IV, 131
ancistrodes, IV, 149
anguineus, IV, 134
anizogonus, IV, 282
arbores, I, 209, 210
areolusos, IV, 286
argenteus, IV, 172
atratus, III, 97
atorruber, IV, 172
atrosanguineus, IV, 172
aulacanthus, IV, 171
aulacothele, IV, 30
aurantiacus, I, 107
aurantiiformis, III, 238
aureiceps, IV, 114
aureus, II, 105
auricomus, IV, 118
auroreus, IV, 118
ayrampo, IV, 261
bahiensis, III, 222, 234
barbatus, IV, 144
barlowii, IV, 172
bellatulus, IV, 165
beneckeii, IV, 171
berteri, III, 97, 98
bertini, IV, 285
bicolor, III, 105
biglandulosus, IV, 31
bihamatus, IV, 140
bispinus, IV, 108
bleo, I, 17, 63
bocanusus, IV, 147
bockii, IV, 78
bolivianus, IV, 256
bonplandii, I, 223
bradypus, I, 121; II, 27
brandegei, IV, 73, 74
brasiliensis, I, 209
brevimamma, IV, 31
brevisetus, IV, 172
broadwayi, III, 217, 222, 225, 229
brownii, IV, 33
brunneus, IV, 15
bulbispinus, IV, 255
caesius, 111, 222, 232, 233, 234, 235
caespitius, IV, 165
calcaratus, IV, 48
californicus, I, 8
campachianus, IV, 253
candelaris, II, 63, 145
canescens, IV, 113
capillaris, IV, 122, 123
caripensis, II, 124; IV, 226
carneus, IV, 88
cassithoides, IV, 227
caudatus, IV, 175
celsianus, IV, 112
centricirrhos, IV, 78
centrispinus, IV, 5
cephalophorus, IV, 41
ceratocentrus, IV, 32
chiloensis, II, 137
chinensis, I, 156, 157
chlorocarpus, II, 224
chrysacanthus, IV, 122
cirrhifer, IV, 91
clavatus, IV, 255
clavus, IV, 30
coccinellifer, I, 179, 224; IV, 252
coccineus, III, 79
cochenillifer, I, 34, 35, 173; IV, 253
columnaris, IV, 99, 107
communis, III, 224, 225, 226
compactus, IV, 36

Cactus—continued,
 compressus, I, 127; IV, 91
 conicus, IV, 70
 conodeus, IV, 17
 conoideus, IV, 17
 conopseus, IV, 78
 coquimbatus, II, 88, 139
 corniferus, IV, 39
 cornigerus, IV, 287
 corollarius, IV, 172
 coronarius, IV, 176
 coronatus, III, 224; IV, 172, 176
 corrugatus, I, 95; III, 179
 crassispinus, IV, 122
 crebrispinus, IV, 18
 crinitus, IV, 150
 crispatus, III, 116
 crocidatus, IV, 87
 cruciformis, IV, 215
 cruciger, IV, 113
 curassavicus, I, 102; IV, 257
 curvispinus, III, 203; IV, 172
 cylindraceus, IV, 127
 cylindricus, I, 63, 77; IV, 140, 141, 176, 231
 dasyacanthus, IV, 55
 dealbatus, IV, 110
 decipiens, IV, 131
 decumanus, I, 80
 densispinus, IV, 119, 120
 densus, IV, 134
 dentatus, IV, 214, 241
 depressus, III, 177; IV, 132, 140
 diadema, IV, 78
 dillenii, I, 162, 163
 disciformis, III, 106
 discolor, IV, 132
 divaricatus, II, 151; IV, 78, 278
 divergens, IV, 78
 dolichocentrus, IV, 106
 dyckianus, IV, 107
 eborinus, IV, 173
 eburneus, I, 95; II, 225
 echinaria, IV, 136
 echinocactodes, IV, 17
 echinocactoides, IV, 17
 echinocarpus, IV, 254
 echinus, IV, 18, 42
 ehrenbergii, IV, 78
 elatior, I, 153
 elegans, IV, 107, 205
 elephantidens, IV, 32
 elongatus, I, 179; IV, 134
 emoryi, IV, 255
 engelmannii, IV, 27, 28
 ensiformis, IV, 201
 epidendrum, IV, 227
 erectacanthus, IV, 173
 erectus, IV, 32
 eriacaanthus, IV, 127
 erinaceus, III, 198; IV, 290
 eschanzieri, IV, 156
 euchlorus, IV, 173
 euphorbioides, II, 33
 eximius, IV, 118
 exsudans, IV, 31
 fascicularis, II, 141
 fasciculatus, IV, 162, 229
 fellneri, IV, 173
 ferox, I, 199, 200, 206
 ficus-indica, I, 177
 fimbriatus, I, 13; II, 87, 151
 fischeri, IV, 95
 flagelliformis, II, 217, 218, 219
 flavescens, IV, 166
 flavispinus, II, 60
 floribundus, III, 97
 foersteri, IV, 78
 foliosus, I, 105; III, 179
 formosus, IV, 90
 foveolatus, IV, 83
 fragilis, I, 193
 frutescens, I, 27

Cactus—continued,
 fulvispinus, IV, 176
 fulvispinosus, II, 50
 funalis, I, 231
 funkii, IV, 93
 fuscatus, IV, 122
 gabbii, IV, 73, 74
 garipensis, IV, 226
 geminatus, IV, 173
 geminispinus, IV, 98
 gibbosus, III, 18, 159, 177; IV, 173, 288
 glabratus, IV, 173
 gladiatus, IV, 78
 glanduliger, I, 31
 glaucus, I, 78
 glochidiatus, IV, 149
 glomeratus, IV, 124
 goodridgii, IV, 158
 gracilis, II, 151; IV, 136
 grahamii, IV, 155, 156
 grahlianus, III, 209
 grandicornis, IV, 173
 grandiflorus, II, 196, 197, 198; IV, 17, 251
 granulatus, IV, 125
 griseus, IV, 166
 guilleminianus, IV, 131
 gummiifer, IV, 74
 gummiiferus, IV, 74
 haageanus, IV, 110
 haematactinus, IV, 173
 halei, IV, 22
 hamatus, IV, 140
 harlowii, III, 221, 222, 232
 haworthianus, IV, 124
 haworthii, II, 44
 heinei, IV, 166
 helicteres, IV, 167
 hemisphaericus, IV, 75
 heptagonus, II, 43; III, 237
 heterocladus, I, 210
 heterogonus, III, 237
 heteromorphus, IV, 25
 hexacanthus, IV, 167
 hexagonus, II, 3, 4, 5, 43, 223; IV, 266
 heyderi, IV, 75
 heyderi hemisphaericus, IV, 75, 76
 horridus, I, 21; III, 202, 203; IV, 251, 259
 humboldtii, II, 163; IV, 130
 humifusus, I, 127
 humilis, I, 113
 hybridus, IV, 198
 hyptiacanthus, III, 156
 hystrix, II, 86; I, 78
 icosagonus, II, 160
 imbricatus, IV, 254
 incurvus, I, 173
 indicus, I, 156
 intertextus, IV, 134
 intortus, III, 220, 221, 222, 230, 231
 irregularis, I, 167
 isabellinus, I, 118
 italicus, I, 266
 jamacaru, II, 8
 jucundus, IV, 173
 kageneckii, II, 20
 karwinskianus, IV, 95
 kleinei, IV, 173
 klugii, I, 107
 kotschoubeyi, III, 82
 kotschubei, III, 82
 kotschubeyi, III, 82
 krameri, IV, 78
 kunthii, IV, 107
 lactescens, IV, 78
 laetus, II, 99
 lamarckii, III, 224, 225
 lanatus, II, 61, 62
 lanceolatus, I, 179
 langsdorffii, III, 199, 200
 lanifer, IV, 113
 lanuginosus, II, 49

Cactus—continued,
 lanuginosus aureus, II, 20
 lasiacanthus, IV, 128
 lasiacanthus denudatus, IV, 129
 latimamma, IV, 41
 latispinus, III, 143
 lecchii, II, 20
 lehmannii, IV, 31
 lemairei, III, 222, 225, 226
 leucocentrus, IV, 167
 leucodasys, IV, 173
 leucodictus, IV, 173
 leucotrichus, IV, 93
 linkeanus, IV, 118
 linkii, I, 121; III, 193
 lividus, IV, 173
 longimamma, IV, 163
 longisetus, IV, 91
 loricatus, IV, 168
 lucidus, I, 10
 ludwigii, IV, 76
 luteus, III, 238
 lyratus, IV, 217, 218
 macracanthus, III, 222, 226; IV, 79
 macrocanthos, III, 222
 macromeris, IV, 25
 macrothele, IV, 30
 maculatus, IV, 15
 magnimamma, IV, 77
 mallisonii, II, 219
 mammillaris, I, 4; IV, 53, 65, 70, 71, 166
 mammillaris glaber, IV, 70
 mammillaris lanuginosus, IV, 166
 mammillaris prolifer, IV, 124
 martianus, IV, 31
 maschalacanthus, IV, 93
 maximus, IV, 263
 maxonii, III, 221, 222, 227, 228
 megacanthus, IV, 78
 meiacanthus, IV, 84
 meissneri, IV, 107
 melaleucus, IV, 63
 melocactoides, III, 222, 228, 235
 melocactus, II, 29, 30 III, 220, 221, 222, 224, 233
 melocactus communis, III, 224
 mensarum, III, 238
 meonacanthus, III, 224
 mexicanus, II, 197
 micans, IV, 173
 micracanthus, IV, 173
 micranthus, IV, 239
 microceras, IV, 78
 microdasys, I, 120, 121
 micromeris, III, 93
 micromeris greggii, III, 93
 microthele, IV, 70, 109
 minimus, IV, 134
 mirabilis, IV, 118
 missouriensis, IV, 53
 missouriensis robustior, IV, 52
 missouriensis similis, IV, 52
 mitis, IV, 175
 moniliformis, I, 206, 207
 monocanthos, I, 156
 mucronatus, IV, 173
 muehlenpfordtii, IV, 112
 multangularis, II, 19; III, 139, 142; IV, 279
 multiceps, IV, 125
 multisectus, IV, 173
 mutabilis, IV, 93
 mystax, IV, 93
 nanus, I, 217
 napoleonis, II, 191
 neglectus, IV, 290
 neo-mexicanus, IV, 45
 neryi, III, 222, 236, 237
 neumannianus, IV, 77, 78

Cactus—continued,
 niger, II, 44; IV, 171
 nigricans, I, 153; IV, 116, 262
 niveus, IV, 98
 nivosus, IV, 71
 nobilis, III, 123, 141, 143, 158; IV, 98
 nopal, IV, 253
 notestini, IV, 53
 nudus, IV, 168
 oaxacensis, IV, 289
 obconella, IV, 106
 obliquus, IV, 173
 obtusipetalus, III, 222, 232
 obvallatus, III, 115; IV, 174
 octacanthus, IV, 31
 octogonus, II, 4; III, 238
 odieranus, IV, 122
 olorius, IV, 174
 oothele, IV, 174
 opuntia, I, 43, 115, 127, 128, 129, 163, 177
 opuntia inermis, I, 161, 162
 opuntia nana, I, 127, 128
 opuntia polyanthos, I, 115
 opuntia tuna, I, 157
 opuntiaeflorus, I, 27
 oreas, III, 222, 227
 ottonis, I, 121; III, 195, 196, 197; IV, 26
 ovatus, II, 20
 ovimamma, IV, 174
 ovoidea, I, 95
 oxypetalus, IV, 189
 pallescens, IV, 88
 palmeri, IV, 140
 paniculatus, II, 82; IV, 280, 281
 paradoxus, I, 209
 parasiticus, IV, 219, 225, 229
 parkinsonii, IV, 98
 parryi, IV, 254
 parvifolius, IV, 266
 parvimamma, IV, 70
 parvispinus, III, 237
 pazzanii, IV, 78
 pectinatus, IV, 34
 pendulinus, IV, 226
 pendulus, IV, 225, 227
 pentacanthus, IV, 78
 pentagonus, II, 15, 121, 123, 193
 pentlandii, I, 98
 pereskia, I, 9, 10, II; IV, 251
 persicanus, IV, 174
 peruvianus, II, 11, 13, 225; III, 234; IV, 266
 peruvianus jamaicensis, IV, 278
 pfeifferanus, IV, 39
 phaeacanthus, IV, 116
 phaeotrichus, IV, 174
 phellospermus, IV, 60
 phyllanthoides, IV, 204, 205
 phyllanthus, IV, 186, 187, 188, 197, 198
 phymatothele, IV, 76
 pictus, IV, 169
 pitajaya, II, 15, 123
 placentiformis, III, 219, 220
 plaschnickii, IV, 31
 plecostigma, IV, 169
 pleiocephalus, IV, 174
 polyanthos, I, 113, 115; IV, 259
 polycentrus, IV, 118
 polycephalus, IV, 113
 polyedrus, IV, 102
 polygonus, II, 47; IV, 101
 polymorphus, II, 139, 174
 polythele, IV, 88
 polytrichus, IV, 102
 pomaceus, IV, 118
 pondii, IV, 23
 portulacifolius, I, 23
 pottsi, IV, 136
 praelii, IV, 96

Cactus—continued,

pretiosus, IV, 118
 pringlei, IV, 114, 116
 prismaticus, II, 123; III, 80
 procerus, IV, 174
 prolifer, IV, 71, 124
 proliferus, IV, 124
 proteiformis, III, 238
 pruinosis, II, 88
 pseudococcinellifer, I, 153
 pseudomammillaris, IV, 132, 133
 pseudotuna, III, 238
 pubescens, IV, 257
 pugionacanthus, IV, 174
 pulchellus, IV, 132
 pulcher, IV, 160
 pulcherrimus, IV, 118
 pulvilliger, IV, 286
 purpureus, IV, 174
 pusillus, I, 105; IV, 124
 pycnacanthus, IV, 41
 pyramidalis, III, 222
 pyrrhocephalus, IV, 99
 pyrrhocroanthus, IV, 122
 quadrangularis, II, 124
 quadratus, IV, 175
 quadriflorus, IV, 254
 quadrispinus, IV, 88
 radians, IV, 36
 radians pectinoides, IV, 36, 37
 radiosus, IV, 43
 radiosus alversonii, IV, 46
 radiosus arizonicus, IV, 45
 radiosus chloranthus, IV, 43
 radiosus deserti, IV, 46
 radiosus neo-mexicanus, IV, 45
 recurvatus, IV, 27
 recurvispinus, IV, 51
 recurvus, III, 123, 141, 142; IV, 78
 reductus, III, 158
 regalis, IV, 266
 regius, IV, 174
 repandus, II, 17, 151, 152
 reptans, IV, 290
 reticulatus, IV, 264
 retusus, IV, 38
 raphidacanthus, IV, 15
 rhodanthus, IV, 122
 rhodanthus sulphureospinus, IV, 116, 122
 rhodocentrus, IV, 174
 robustior, IV, 2
 robustispinus, IV, 33
 rosa, I, 19, 20
 roseanus, IV, 22
 roseus, IV, 174
 rotundifolius, I, 27
 royenii, II, 50
 rueschianus, IV, 174
 ruestii, III, 221, 222, 227
 ruficeps, IV, 122
 rufidulus, IV, 174
 rufocroceus, IV, 174
 rutilus, IV, 169
 salicornioides, IV, 217, 218
 salm-dyckianus, IV, 39
 salmianus, I, 74
 salvador, III, 222, 228, 229
 saxatilis, IV, 169
 scepontocentrus, IV, 41
 schaeferi, IV, 112
 scheeri, IV, 28
 scheidweilerianus, IV, 148
 schelhasi, IV, 149
 schiedeanus, IV, 128
 schilinzkyanus, III, 210
 schlechtendalii, IV, 30
 schlumbergeri, IV, 269
 scolymoides, IV, 18, 39
 scolymoides sulcatus, IV, 48
 scopa, III, 193
 seemannii, IV, 170
 seidelii, IV, 174

Cactus—continued,

seitzianus, IV, 83
 sempervivi, IV, 86
 senilis, II, 25, 27; IV, 19
 sepium, II, 160
 sericeus, I, 134
 serpens, II, 163
 serpentinus, II, 118, 119
 setispinus, IV, 22
 setosus, IV, 88
 severinii, IV, 174
 similis, IV, 52
 solitarius, IV, 175
 sororius, IV, 170
 speciosissimus, II, 128, 129; IV, 277
 speciosissimus lateritius, II, 128; IV, 277
 speciosus, II, 127, 128; IV, 175, 205, 244, 277
 speciosus grandiflorus, IV, 201
 speciosus lateritius, IV, 277
 spectabilis, IV, 174
 sphacelatus, IV, 138
 sphaericus, IV, 61
 sphaerotrachus, IV, 130
 spinaureus, IV, 170
 spini, IV, 132, 133
 spinosissimus, I, 204; IV, 118, 119
 spinosus, IV, 175
 splendidus, IV, 253
 squarrosus, IV, 91
 stella-auratus, IV, 134
 stellaris, IV, 125
 stellatus, IV, 124, 125
 stellatus texanus, IV, 125
 stenocephalus, IV, 122
 stramineus, IV, 166
 strictus, I, 161; II, 44
 strobiliformis, IV, 54
 stueberi, IV, 122
 subangularis, IV, 91
 subcroceus, IV, 134
 subcurvatus, IV, 78
 subechinatus, IV, 134
 subinermis, I, 34
 subpolyedrus, IV, 105
 subquadriflorus, IV, 254
 subquadrifolius, I, 65; IV, 254
 subrepandus, IV, 278
 subtetragonus, IV, 88
 subulifer, IV, 174
 sulcatus, IV, 48
 sulcolanatus, IV, 37
 sulphureus, I, 134
 supertextus, IV, 107
 sylvestris, III, 238
 tectus, IV, 174
 tentaculatus, IV, 122
 tenuis, IV, 134, 215
 teres, IV, 227
 tetracanthus, IV, 106
 tetracentrus, IV, 78
 tetragonus, II, 9, 14
 tetrancistrus, IV, 60
 texanus, IV, 125
 texensis, IV, 76
 tomentosus, I, 73; IV, 175
 torquatus, IV, 245
 townsendii, III, 222, 234
 triacanthos, I, 112
 triacanthus, IV, 91
 triangularis, II, 183, 187, 188, 192, 193, 194, 212; IV, 282
 triangularis aphyllus, II, 187
 triangularis foliaceus, II, 192
 trichotomus, III, 238
 trigonus, II, 192
 triqueter, II, 192; IV, 243
 truncatus, IV, 177, 179, 180
 tuberculatus, I, 214
 tunculosus, IV, 54
 tuna, I, 113, 114, 163
 tuna elatior, I, 153

Cactus—continued,

tuna major, IV, 289
 tuna nigricans, I, 153
 tunicatus, I, 65
 turbinatus, III, 106
 uberiformis, IV, 63
 umbrinus, IV, 164
 uncinatus, IV, 140
 undulosus, II, 123
 urumbaba, I, 156, 157
 urumbella, I, 157
 varimamma, IV, 175
 versicolor, IV, 78
 verticillatus, III, 238
 vetulus, IV, 130
 villifer, IV, 102
 villosus, III, 103
 virens, IV, 95
 viridis, IV, 96
 viviparus, IV, 43, 44, 45
 vrieseanus, IV, 174
 vulpinus, IV, 118
 webbianus, IV, 87
 wegneri, IV, 175
 wildianus, IV, 143
 winkleri, IV, 41
 woburnensis, IV, 100
 wrightii, IV, 152
 xanthotrachus, IV, 93
 zegschwitzii, IV, 175
 zehntneri, III, 222, 235, 236
 zephyranthodes, IV, 159
 zepnickii, IV, 175
 zinniaeflorus, I, 21
 Cactuses, II, 116
 Calibanus caespitosus, IV, 132
 Camussa, I, 191
 Candebobe, II, 96
 Cane cactus, I, 43
 Capilliformes, IV, 220
 Cardon, II, 70, 96; III, 232; IV, 277
 Cardon grande, II, 140
 Cardoncillo, II, 112
 Carnegiea, II, 2, 164-167, 178
 gigantea, II, 133, 164, 165, 166, 178; IV, 155, 279, 290
 Cassuthae, IV, 220
 Cassytha, IV, 219
 baccifera, IV, 227
 filiformis, IV, 219, 225
 polysperma, IV, 227
 Cephalocactus, III, 85
 Cephalocereus, I, 116; II, 1, 3, 13, 25-60, 178, 224, 225; III, 6, 7; IV, 268
 alensis, II, 26, 55
 arrabidae, II, 26, 42, 43, 136; IV, 268
 bahamensis, II, 26, 38
 bakeri, II, 39, 40
 barbadensis, II, 26, 44, 45, 46
 brasiliensis, II, 26, 57
 brooksonianus, II, 26, 49
 californicus, II, 224
 catingicola, II, 26, 49, 56; IV, 269
 chrysacanthus, II, 26, 48
 chrysomallus, II, 72
 collinsii, IV, 269, 270
 colombianus, II, 26, 34, 55, 56
 columna, II, 76
 columna-trajani, II, 76
 cometes, II, 26, 51, 52
 compressus, II, 193
 deeringii, II, 26, 38, 39
 delaetii, III, 6
 dybowskii, II, 25, 30, 58
 euphorbioides, II, 25, 33
 exerens, II, 42
 fluminensis, II, 25, 29, 33, 57; IV, 268
 fouachianus, II, 51
 gaumeri, II, 26, 47
 gounellei, II, 25, 34, 35; III, 236
 hermentianus, II, 58; IV, 270

Cephalocereus—continued,

hoppenstedtii, II, 25, 27, 225
 keyensis, II, 26, 40
 lanuginosus, II, 18, 26, 49, 50; IV, 269
 leucocephalus, II, 26, 52, 53; IV, 269
 leucostele, II, 25, 36, 37, 39, 60
 macrocephalus, II, 25, 31, 75, 76; IV, 268
 maxonii, II, 26, 48, 53
 melanostele, II, 167; IV, 279
 melocactus, II, 29, 30, 8
 millsapghii, II, 26, 45, 46
 monoclonos, II, 26, 40, 41
 moritzianus, II, 26, 41, 42
 nobilis, II, 26, 44, 45; IV, 269
 palmeri, II, 26, 53
 pasacana, IV, 277
 pentaedrophorus, II, 25, 31
 phaeacanthus, II, 26, 57
 piahyensis, II, 26, 48, 49
 polygonus, II, 26, 47; IV, 269
 polylophus, II, 25, 32; IV, 268
 purpureus, II, 25, 28, 29
 purpusii, II, 26, 56; IV, 269
 robinii, II, 26, 39, 40
 robustus, II, 26, 51, 52
 royenii, II, 26, 46, 50; IV, 269
 russelianus, II, 25, 33, 34, 56
 sartorianus, II, 26, 53
 schlumbergeri, IV, 269
 scoparius, II, 26, 41
 senilis, II, 25, 27, 28, 31; III, 4, 6; IV, 268
 smithianus, II, 26, 36, 37
 swartzii, II, 26, 46, 47
 tetetzo, IV, 272
 tweedyanus, II, 26, 54, 55
 ulei, II, 26, 52, 58
 urbanianus, II, 26, 43
 zehntneri, II, 25, 35
 Cephaloidei, III, 8, 90, 176
 Cephalophora, II, 25
 Cephalophorus, II, 25
 columna-trajani, II, 76
 senilis, II, 27
 Ceratistes copiapensis, III, 186
 Cereanae, II, 1; III, 3; IV, 177
 Cereace, I, 8, 24, 215; II, 1, 111; III, 3; IV, 3
 Cerei, II, 59, 70
 Cereus, I, 8, 75, 151, 215, 217; II, 1, 3-21, 33, 41, 42, 47, 58, 59, 68, 77, 82, 87, 105, 108, 110, 118, 122, 127, 144, 145, 147, 152, 158, 163, 164, 170, 173, 192, 219, 221, 222, 223, 224, 225; III, 3, 8, 11, 23, 25, 45, 60, 76, 78, 79, 121, 132, 154, 166, 237; IV, 48, 64, 71, 180, 186, 189, 208, 209, 210, 211, 266, 267, 268, 274
 abnormis, II, 12
 acanthosphaera, II, 209
 acanthurus, II, 161
 acidus, II, 84
 acifer, III, 12
 aciniformis, III, 23
 ackermannii, IV, 198
 acranthus, II, 168
 acromelas, II, 59
 aculeatus, II, 21
 acutangulus, II, 123, 124, 157
 adscendens, II, 155, 156
 adustus, III, 23
 adustus radians, III, 24
 aethiops, II, 4, 16, 17, 18; IV, 266
 affinis, II, 14
 aggregatus, III, 14; IV, 47
 alacriportanus, II, 4, 6, 7

Cereus—continued,
 alamosensis, II, 169
 alatus, IV, 213, 243
 alatus crassior, IV, 214
 albertinii, II, 21
 albiflorus, II, 128
 albisetosus, II, 8
 albispinus, II, 37, 59, 118
 albispinus major, II, 59
 alensis, II, 5
 amalonga, IV, 267
 amalonga cristata, IV, 267
 amazonicus, II, 24
 ambiguus, II, 118
 ambiguus strictior, II, 118
 amblyogonus, II, 20
 amecaensis, II, 129
 amecanensis, II, 129; IV, 277
 americanus octangularis, II, 89
 americanus triangularis, II, 193
 amoenus, III, 33
 andalgalensis, III, 6
 andryanus, II, 59
 anguiniformis, II, 22, 174
 anguinus, II, 175
 angulosus, II, 32
 anisacanthus, II, 102
 anisacanthus ortholophus, II, 102
 anisacanthus subspiralis, II, 102
 anisitsii, II, 23
 anizogonus, II, 193
 anomalus, IV, 181
 antoinii, II, 201
 apiciflorus, II, 107
 aquicaulensis, II, 180
 aragonii, II, 92, 103
 aragonii palmatus, II, 92
 arboreus, II, 80
 arcuatus, II, 124
 arendtii, II, 154
 areolatus, II, 159
 arequipensis, II, 134, 145
 argentincensis, II, 4, 11, 12
 armatus, II, 50
 arrabidae, II, 42, 43
 arrigens, II, 180
 articulatus, I, 89
 assurgens, II, 77, 79
 atacanensis, II, 145
 atropurpureus, II, 154
 atrovirens, II, 21
 aurantiacus, II, 128
 aurantiacus superbus, II, 129
 auratus, IV, 268
 auratus genuinus, IV, 268
 auratus intermedius, IV, 268
 auratus mollissimus, IV, 268
 auratus pilosus, IV, 268
 aureus, II, 44, 105, 106, 107
 aureus pallidior, II, 44
 aurivillus, II, 173, 226
 aurora, II, 219
 azureus, II, 4, 15, 16
 azureus seidelii, II, 15
 baccilerus, IV, 227
 bahamensis, II, 38
 bajanensis, II, 124
 bakeri, II, 39
 balansaei, II, 157, 158
 barbatus, II, 51
 baumannii, II, 173, 174, 217
 baumannii colubrinus, II, 174
 baumannii flavispinus, II, 174
 baumannii smaragdiflorus, II, 174
 bavosus, II, 86; IV, 273
 baxaniensis, II, 123, 124; IV, 276
 baxaniensis ramosus, II, 124
 belieuli, II, 96
 beneckeii, II, 18; IV, 273
 beneckeii farinosus, II, 18; IV, 273

Cereus—continued,
 bergerianus, II, 72
 berlandieri, III, 20, 21; IV, 285
 bertini, III, 44
 bertinii, III, 44, 45
 beysiegelii, II, 223
 bififormis, IV, 201, 202
 bifrons, II, 128
 bigelovii, III, 8, 9
 bigelovii zuniensis, III, 14
 biolleyi, II, 215
 blankii, III, 20
 blankii, III, 21; IV, 285
 boeckmannii, II, 202; IV, 283
 bolivianus, II, 136
 bonariensis, II, 7
 bonplandii, II, 157, 158; IV, 278
 bonplandii brevispinus, II, 157
 bonplandii pomanensis, II, 155
 brachiatus, II, 86
 brachypetalus, II, 67
 bradtianus, I, 215
 bradypus, II, 27
 brandegeei, III, 34
 brandii, II, 14
 breviflorus, II, 82, 83
 brevispinulus, II, 200
 brevistylus, II, 66
 bridgesii, II, 134
 bridgesii brevispinus, II, 134
 bridgesii lageniformis, II, 134
 bridgesii longispinus, II, 134
 brookii, II, 151
 brooksianus, II, 49
 caesius, II, 4, 13, 15
 caespitosus, III, 25, 26; IV, 285
 caespitosus castaneus, III, 25
 caespitosus major, III, 25
 caespitosus minor, III, 25
 calcaratus, II, 193
 californicus, I, 8; II, 224
 callicanthus, II, 197
 callicochi, III, 182
 calvescens, II, 11
 calvus, II, 69, 70
 candelabrus, II, 126
 candelabrum, II, 96; IV, 273
 candelaris, II, 63
 candicans, II, 142, 143; III, 28; IV, 277
 candicans courantii, II, 142
 candicans dumesnilianus, II, 143; IV, 277
 candicans gladiatus, II, 142
 candicans robustior, II, 142
 candicans spinosior, II, 143
 candicans tenuispinus, II, 142
 caracore, IV, 266
 caripensis, II, 124; IV, 226
 castaneus, II, 84
 catamarcensis, II, 146
 catingae, IV, 269
 catingicola, II, 56; IV, 269
 cauchinii, II, 8
 caudatus, II, 20; IV, 175
 cavendishii, II, 21, 22; IV, 268
 celsianus, II, 171, 172
 chacoanus, II, 224
 chalybaeus, II, 4, 16, 17, 223
 chende, II, 90, 91
 chichiipe, II, 89, 90
 childsi, IV, 266
 chilensis, II, 137, 138, 139
 chilensis acidus, II, 84
 chilensis breviflorus, II, 83
 chilensis brevispinulus, II, 139

Cereus—continued,
 chilensis eburneus, II, 139
 chilensis flavescens, II, 139
 chilensis fulvibarbis, II, 139
 chilensis funkianus, II, 138
 chilensis heteromorphus, II, 137
 chilensis linnaei, II, 139
 chilensis nigripilis, II, 140
 chilensis panhoplites, II, 137
 chilensis polygonus, II, 137
 chilensis poselgeri, II, 137
 chilensis pycnanthus, II, 137
 chilensis quisco, II, 139
 chilensis spinosior, II, 139
 chilensis zizkaanus, II, 137, 224
 chilensis zizkeanus, II, 224
 chiloensis, I, 79 II, 84, 137, 138
 chiloensis lamprochlorus, II, 133
 chiotilla, II, 6, 66; IV, 270
 chloranthus, III, 16, 17
 chlorocarpus, II, 224
 chotaensis, II, 163
 chrysacanthus, II, 48
 chrysomallus, II, 72
 cinerascens, III, 23
 cinerascens crassior, III, 23
 cinerascens tenuior, III, 23
 cinnabarinus, II, 129; IV, 277
 cirrhiferus, III, 23
 clavarioides, I, 73
 clavatus, II, 94
 claviformis, III, 43
 coccineus, II, 127, 210, 211, 212; III, 14; IV, 48
 cochal, II, 180; IV, 280
 coerulescens, II, 17, 59; IV, 266
 coerulescens fulvispinus, II, 17
 coerulescens landbeckii, II, 17
 coerulescens longispinus, II, 17
 coerulescens melanacanthus, II, 17
 coeruleus, II, 17
 cognatus, II, 123
 colombianus, II, 55
 colubrinus, II, 174, 217; IV, 279
 colubrinus flavispinus, II, 174
 colubrinus smaragdiflorus, II, 175
 columna-trajani, II, 76
 columnaris, II, 224
 colvillii, II, 14
 cometes, II, 52
 compressus, II, 192, 193
 concinnus, II, 21
 concolor, III, 25, 26
 conformis, II, 103
 conglomeratus, III, 39
 conicus, II, 33
 coniflorus, II, 199
 conoideus, III, 13, 14
 coquimbanus, II, 83, 139, 225; IV, 272
 coracare, II, 10
 coryne, II, 64, 65; IV, 270
 cossyrensis, II, 152
 crenatus, II, 59; IV, 192
 crenulatus, II, 49, 59
 crenulatus gracilior, II, 49
 crenulatus griseus, II, 87
 crimsonii, II, 219
 crispatus, IV, 245
 crispatus crenulatus, IV, 244
 crispatus laevior, IV, 244
 cruciformis, IV, 213, 216
 ctenoides, III, 19

Cereus—continued,
 cubensis, II, 149
 cumengei, II, 116
 cupulatus, II, 74
 curtisii, II, 44
 cyaneus, IV, 266
 cylindricus, I, 77
 damacaro, II, 21
 damazioi, II, 159
 dasyacanthus, III, 19
 dasyacanthus minor, III, 31
 dasyacanthus neo-mexicanus, II, 19
 dautwitzii, II, 61, 62
 davisii, II, 154
 dayamii, II, 4, 11
 decagonus, II, 59
 decandollii, II, 13
 decorus, II, 20
 decumbens, II, 162
 deficiens, II, 94
 de laguna, II, 20
 del morali, II, 90, 91; IV, 273
 denudatus, III, 155
 deppei, III, 23
 devauxii, II, 128
 dichroacanthus, III, 97, 98
 diguetii, II, 111
 divaricatus, II, 151
 divergens, II, 151
 donatii, II, 203
 donkelarii, II, 200; IV, 230
 donkelarii, II, 200
 donkelarii, II, 200
 dubius, III, 3
 duledevantii, II, 139
 dumesnilianus, II, 143, 159
 dumortieri, II, 102
 dusenii, III, 197
 dussii, II, 123
 dybowskii, II, 30
 dyckii, II, 92, 93
 eburneus, II, 20, 87, 89, 99, 103, 224, 225; III, 43; IV, 273
 eburneus clavatus, II, 94
 eburneus monstrosus, II, 88
 eburneus polygonus, II, 87, 88
 edulis, II, 89
 ehrenbergii, III, 41
 elegans, II, 139; IV, 216
 emoryi, II, 108; IV, 274
 engelmannii, II, 38
 engelmannii albispinus, IV, 285
 engelmannii caespitosus, IV, 285
 engelmannii chrysocentrus, III, 38
 engelmannii fulvispinus, IV, 285
 engelmannii pfersdorffii, IV, 285
 engelmannii variegatus, III, 38
 enneacanthus, III, 36; IV, 285
 enriquezii, II, 88
 erectus, II, 49, 152
 erectus cristatus, II, 193
 erectus maximus, II, 13
 ericomus, II, 4
 erinaceus, IV, 290
 eriocarpus, II, 145
 eriophorus, II, 149, 202; IV, 278
 eriophorus laeteviridis, II, 149
 eruca, II, 114, 115, 116; IV, 276
 erythrocephalus, III, 7
 estrellensis, II, 216
 euchlorus, II, 21, 22; IV, 268
 euphorbioides, II, 33
 exerens, II, 42, 43
 extensus, II, 191
 eyriesii, II, 159; III, 65

Cereus—continued,

farinosus, II, 18; IV, 273
 fascicularis, II, 141
 fendleri, III, 35, 36
 fendleri pauperculus, III, 35
 fercheckii, II, 140
 fernambucensis, II, 14
 ferox, II, 30
 fimbriatus, II, 151
 flagelliformis, II, 218, 219;
 IV, 21, 284
 flagelliformis cristatus, IV,
 284
 flagelliformis funkii, II, 219
 flagelliformis leptophis, II,
 218
 flagelliformis mallisoni, II,
 219
 flagelliformis minor, II, 218,
 219; IV, 230
 flagelliformis nothus, II, 219
 flagelliformis scotii, II, 219
 flagelliformis smithii, II, 219
 flagelliformis speciosus, II,
 219
 flagriformis, II, 219, 220
 flavescens, II, 20; III, 212;
 IV, 71
 flavicomus, II, 52
 flaviflorus, III, 15
 flavispinus, II, 20, 60
 flavispinus hexagonus, II,
 224
 flemingii, IV, 283
 flexuosus, IV, 116
 floccosus, II, 50, 51
 fluminensis, II, 29
 foersteri, II, 53
 forbesii, II, 7
 formosissimus, IV, 286
 formosus, II, 14
 formosus monstrosus, IV, 266
 fouchianus, II, 50
 foveolatus, III, 98
 fulgens, II, 210
 fulgidus, II, 210
 fulvibarbis, II, 139
 fulviceps, II, 72
 fulvispinosus, II, 50
 fulvispinus, II, 140
 funkii, II, 137
 galapagensis, II, 146, 147
 garambello, II, 180
 geminisetus, II, 224
 gemmatus, II, 74; III, 66,
 67; IV, 271
 geometrizans, II, 20, 179,
 180; IV, 280
 geometrizans cochal, II, 180
 geometrizans pugioniferus, II,
 179, 180
 geometrizans quadrangula-
 rispinus, II, 179, 180
 geometrizans quadranguli-
 spinus, II, 180
 ghesbreghtii, II, 60
 gibbosus, III, 158
 giganteus, II, 135, 164, 167;
 IV, 279
 gilliesii, III, 75
 gilvus, II, 137
 glaber, II, 216
 gladiator, II, 179
 gladiator geometrizans, II,
 180
 gladiatus, II, 142
 gladiatus courantii, IV, 277
 gladiatus vernaculatus, IV,
 277
 gladiger, II, 87, 88, 180
 gladiiger, II, 87
 gladilger, II, 87
 glaucescens, II, 59
 glaucus, II, 8, 15
 glaucus speciosus, II, 14
 glaziovii, II, 109
 glomeratus, III, 15

Cereus—continued,

gloriosus, II, 51
 glycimorphus, III, 23
 gonacanthus, III, 10
 gonzalezii, II, 77, 214
 gracilis, II, 19, 147, 151, 209
 gracilis scandens, II, 198
 grandifloro-speciosissimus,
 IV, 283
 grandiflorus, II, 197, 198,
 199; IV, 282, 283
 grandiflorus affinis, II, 197
 grandiflorus barbadensis, II,
 198
 grandiflorus callicanthus, II,
 197
 grandiflorus flemingii, IV, 283
 grandiflorus grusonianus, II,
 197
 grandiflorus haitiensis, II,
 197
 grandiflorus hybridus, II,
 210
 grandiflorus macdonaldiae,
 IV, 283
 grandiflorus major, II, 198
 grandiflorus maximiliani, II,
 197
 grandiflorus maynardii, II,
 210
 grandiflorus mexicanus, II,
 197
 grandiflorus minor, II, 197
 grandiflorus ophites, II, 197
 grandiflorus ruber, II, 210
 grandiflorus schmidtii, II, 197
 grandiflorus speciosissimus,
 II, 210
 grandiflorus spectabilis, II,
 197
 grandiflorus uranos, II, 197
 grandiflorus viridiflorus, II,
 197
 grandis, II, 14
 grandis gracilior, II, 14
 grandis ramosior, II, 14
 grandispinus, II, 87
 greggii, II, 112, 113, 122;
 IV, 275
 greggii cismontanus, II, 112
 greggii roseiflorus, II, 112
 greggii transmontanus, II,
 112, 113
 grenadensis, II, 4, 18, 223
 griseus, II, 87
 grossei, II, 174
 grusonianus, II, 203; IV, 283
 guasabara, IV, 266
 guatemalensis, II, 89, 119
 guelichii, II, 158
 gummatosus, II, 117
 gummosus, II, 117
 gummosus, II, 116, 117; IV,
 276
 haageanus, II, 19
 haematuricus, II, 8
 haitiensis, IV, 283
 hamatus, II, 203, 205
 hankeanus, II, 3, 7, 8
 hansii, II, 128
 hassleri, II, 211
 haworthii, II, 44
 hayni, III, 102, 103
 hempelianus, II, 136
 heptagonus, III, 237
 hermannianus, II, 17
 hermentianus, II, 8
 heteracanthus, II, 224
 heteromorphus, II, 137
 hexaedrus, III, 10
 hexagonus, II, 4, 5, 9, 13,
 223, 224; III, 238; IV, 266
 hexangularis, II, 14
 hildmannii, II, 103
 hildmannianus, II, 4, 6
 hirschtianus, II, 119; IV,
 276

Cereus—continued,

hoffmannseggii, III, 98
 hollianus, II, 8, 86
 hondurensis, II, 199
 hoogendorpii, II, 22, 5
 hookeri, IV, 197, 198
 hoppenstedtii, II, 27; IV,
 268
 horizontalis, II, 20
 horrens, II, 195
 horribarbis, II, 8
 horridus, II, 5, 9
 houletii, II, 52, 53; IV, 269
 huascha, II, 142; III, 57
 huascha flaviflorus, II, 142
 huascha flaviformis, II, 142
 huascha rubriflorus, III, 6
 hutchinsii, III, 8
 humboldtii, II, 163
 humilis, II, 209, 210
 humilis major, II, 210
 humilis minor, II, 209
 humilis myriacaulon, II, 210
 humilis rigidior, II, 210
 huottii, III, 63
 hyalacanthus, II, 173, 176
 hybridus, II, 210
 hypogaeus, II, 106, 107
 hystrix, II, 86, 87, 99, 103;
 IV, 273
 ianthothele, IV, 210
 ianthothelus, IV, 210
 icosagonus, II, 360
 ictidurus, II, 224
 imbricatus, I, 63
 incrassatus, II, 21
 incrustans, II, 74
 incrustatus, II, 74
 incurvispinus, III, 72
 inermis, II, 207, 208; III, 183
 inermis laetevirens, II, 208
 ingens, III, 171
 insularis, II, 23
 intricatus, II, 143
 inversus, II, 191
 iquiquensis, II, 83
 irradians, II, 202
 isogonus, II, 160, 161
 jacquin, II, 21
 jalapaensis, II, 199
 jamacaru, II, 4, 5, 6, 8, 9,
 15, 18; IV, 266
 jamacaru caesius, II, 15
 jamacaru glaucus, II, 9
 janthothele, IV, 210
 jasmineus, III, 66
 jenkinsonii, II, 128
 jenkinsonii verus, II, 128
 jconostle, II, 93
 josselineus, II, 129
 jubatus, II, 52
 jusburtii, II, 157, 158
 kageneckii, II, 20
 kalbreyerianus, II, 118
 karstenii, II, 5, 207, 208
 karwinski, II, 21
 kerberi, II, 170
 kewensis, II, 40; IV, 283
 knightii, IV, 215
 knippelianus, III, 32
 kostratus, II, 205
 kunthianus, II, 200, 201
 labouretianus, II, 8
 laetevirens, II, 8
 laetevirens caesina, II, 15
 laetus, II, 99
 laevigatus, II, 88
 laevigatus guatemalensis, II,
 89
 lagenaeformis, II, 134
 lamprochlorus, II, 132, 133
 lamprochlorus salinicolus, II,
 133
 lamprospermus, II, 4, 10
 lanatus, II, 61, 62
 lanceanus, II, 191
 landbeckii, II, 17; IV, 266

Cereus—continued,

langlassei, II, 20
 laniceps, II, 173, 175
 lanuginosus, II, 4; IV, 71
 lanuginosus aureus, II, 20
 lanuginosus glaucescens, II,
 49
 lasianthus, II, 134
 lateribarbus, II, 76
 lateritius, II, 128
 latifrons, IV, 188, 189
 lauterbachii, II, 22
 leccii, II, 20
 lecanus, III, 9
 lehmannii, IV, 31
 leiocarpus, II, 50
 lemairei, II, 189; IV, 282
 lemoinei, II, 189
 leonensis, III, 20
 leonii, II, 78
 lepidanthus, II, 76
 lepidotus, II, 4, 5, 6; IV, 284
 leptacanthus, III, 22; IV,
 285
 leptophis, II, 218; IV, 284
 leucanthus, II, 72
 leucostele, II, 36, 37
 limensis, II, 20
 lindbergianus, II, 211
 lindenzweigianus, II, 23
 lindmanii, II, 211
 linkii, III, 195
 linnaei, II, 137
 lividus, II, 8, 9
 lividus glaucior, II, 9
 longicaudatus, II, 205
 longifolius, II, 20
 longipedunculatus, II, 21
 longisetus, III, 42
 longispinus, II, 137; III, 37;
 IV, 285
 lormata, II, 21; III, 237; IV,
 267
 lumbricoides, IV, 230
 lutescens, II, 44
 macdonaldiae, II, 202, 203;
 IV, 283
 macracanthus, III, 43
 macrocephalus, II, 31
 macrogonus, II, 43, 130, 136;
 IV, 277
 macrostibas, II, 181, 182
 maelenii, IV, 8
 magnus, II, 159
 malletianus, II, 145
 mallisonii, II, 219
 mamillatus, III, 41
 margaritensis, II, 4, 18, 224
 marginatus, II, 69, 74; IV,
 197, 271
 marginatus cristatus, IV, 271
 marginatus gemmatus, II, 74
 marginatus gibbosus, II, 74
 marginatus monstrosus, IV,
 271
 mariculi, II, 209
 maritimus, III, 15
 marmoratus, II, 23
 martianus, II, 219, 220, 221;
 IV, 284
 martinii, II, 155, 190
 martinii perviridis, II, 155
 maximiliani, II, 197
 maxonii, II, 48
 maynardae, II, 210
 maynardii, II, 210
 megalanthus, II, 212
 melanacanthus, II, 17
 melanotrichus, II, 68
 melanoma, II, 109, 110
 melocactus, II, 29
 mendory, II, 17
 merkeri, III, 5
 mexicanus, II, 129; IV, 200
 micranthus, IV, 211
 microsphaericus, II, 159; IV,
 181

Cereus—continued,

militaris, II, 73
 militaris californicus, II, 86
 millspaughii, II, 45
 minor, II, 218
 minutiflorus, II, 195
 miravallensis, II, 213
 mirbelii, II, 74
 mixtecensis, II, 89, 90
 moeninghoffii, II, 219
 mojavensis, III, 8, 2
 mojavensis zuniensis, III, 14
 mollis, II, 44
 mollis nigricans, II, 44
 monacanthus, II, 155, 190
 moniliformis, I, 206, 217
 monoclonos, II, 13, 41
 monstrosus, II, 12
 monstrosus minor, II, 12
 monstrosus, II, 12
 montevidensis, III, 98
 montezumae, II, 143
 monvilleanus, II, 173
 moritzianus, II, 41, 42
 moritzianus pfeifferi, II, 42
 multangularis, II, 19, 20, 30
 111, 142, 143; IV, 279
 multangularis albispinus, II, 20
 multangularis limensis, II, 20
 multangularis pallidior, II, 19
 multangularis prolifer, II, 20
 multangularis rufispinus, II, 21
 multicostatus, III, 9
 multiplex, III, 64; IV, 286
 multiplex cristatus, III, 64
 multiplex monstrosus, III, 64
 myosurus, IV, 215
 myosurus tenuior, IV, 216
 myriacaulon, II, 209
 myriophyllus, II, 143
 nanus, I, 217; II, 19
 napoleonis, II, 185, 187, 191;
 IV, 282
 nashii, II, 151
 nelsonii, IV, 283, 284
 nesioticus, II, 120, 121
 neumannii, II, 119
 nickelsii, II, 32
 niger, II, 44
 niger gracilior, II, 44
 nigricans, II, 44
 nigripilis, II, 139, 140 IV,
 277
 nigrispinus, II, 17
 nitens, II, 132
 nitidus, II, 123
 nobilis, II, 44
 northumberlandia, II, 4
 northumberlandianus, II, 4
 nothus, II, 201
 nudiflorus, II, 113, 114
 nycticalus, II, 200, 201, 216
 nycticalus, II, 196, 199; IV,
 283
 nycticalus armatus, II, 199
 nycticalus gracilior, II, 201
 nycticalus maximiliani, II,
 197, 201
 nycticalus peanii, IV, 283
 nycticalus viridior, II, 201
 nyriacaulon, II, 209
 obtusangulus, II, 22; IV, 181
 obtusus, II, 4, 13, 14, 15, 16
 ocamponis, II, 184, 185
 ochracanthus, II, 20
 octacanthus, III, 13
 octogonus, II, 39
 olfersii, II, 33
 oligolepis, II, 225
 olivaceus, II, 225
 opites, II, 21
 orcuttii, II, 70
 ottonis, III, 196
 ovatus, I, 91; II, 20

Cereus—continued,

oxygonus, III, 64
 oxypetalus, IV, 188, 189
 pachyrhizus, II, 4, 10
 pacificus, III, 12
 palmeri, II, 177
 paniculatus, II, 82; IV, 280,
 281
 panoplaeatus, II, 82, 137,
 138
 papillosus, III, 19
 paradisiacus, II, 198
 paradoxus, I, 209
 paraguayensis, II, 6
 parasiticus, IV, 226
 parviflorus, II, 173, 176
 parvisetus, II, 173
 parvulus, IV, 181
 pasacana, II, 133; III, 74, 76
 patagonicus, III, 197
 paucispinus, III, 10, 14
 paxtonianus, II, 21, 22
 peanii, II, 201
 pecten-aboriginum, II, 70,
 71; IV, 271
 pectinatus, III, 29; IV, 285
 pectinatus armatus, III, 24
 pectinatus centralis, III, 149,
 130
 pectinatus laevior, III, 30
 pectinatus rigidissimus, III,
 27
 pectinatus spinosus, III, 24
 pectiniferus, III, 29
 pellucidus, II, 79, 122, 133
 penicillatus, II, 171
 pensilis, III, 8
 pentaedrophorus, II, 31
 pentagonus, II, 123, 195,
 210, 213
 pentagonus glaucus, II, 31
 pentaplophorus, II, 31
 pentaplophus, III, 21, 22
 pentaplophus leptacanthus, III,
 22
 pentaplophus radicans, II, 210;
 III, 21
 pentaplophus simplex, III, 21,
 22
 pentaplophus subarticulatus,
 III, 21, 22
 pentapterus, II, 213
 pepinianus, II, 137
 perlucens, II, 4, 13
 pernambucensis, II, 4, 14,
 15, 213; IV, 266
 perotetti, II, 9
 perrottetiannus, II, 4, 6
 peruvianus, II, 3, 4, 5, 11,
 13, 133, 147; IV, 266
 peruvianus alacriportanus,
 II, 6
 peruvianus brasiliensis, II,
 13
 peruvianus cristatus, II, 12
 peruvianus monstrosus, II,
 12, 13; IV, 266
 peruvianus monstrosus minor,
 II, 13
 peruvianus monstrosus, II,
 223,
 peruvianus monstrosus
 nanus, II, 12
 peruvianus spinosus, II, 13
 peruvianus tortuosus, II, 12
 peruvianus tortuosus, II, 12
 perviridis, IV, 267
 pfeifferi, II, 42
 pfersdorffii, II, 117
 phaeacanthus, II, 57
 phatnospermus, II, 24
 philippii, II, 105
 phoeniceus, III, 14; IV, 48,
 285
 phoeniceus inermis, III, 14
 phoeniceus pacificus, III, 12
 phyllanthoides, IV, 203

Cereus—continued,

phyllanthoides albiflorus, IV,
 205
 phyllanthoides curtisii, IV,
 205
 phyllanthoides guillardieri,
 IV, 203
 phyllanthoides jenkinsonii,
 IV, 205
 phyllanthoides stricta, IV,
 205
 phyllanthoides vandesii, IV,
 205
 phyllanthus, IV, 187, 188
 phyllanthus marginatus, IV,
 188
 piauhensis, II, 49
 pitahaya variabilis, IV, 267
 pitajaya, II, 13, 14, 15, 123
 plagiotoma, II, 163
 platycarpus, IV, 242
 platygonus, II, 11, 156, 157,
 159
 pleiognon, III, 43
 plumieri, II, 191
 polyacanthus, III, 11, 15; IV,
 285
 polychaetus, II, 17
 polygonatus, II, 88
 polygonus, II, 47
 polylophus, II, 32, 33; IV,
 268
 polymorphus, II, 138, 139
 polyptychus, IV, 269
 polyrhizus, II, 185
 pomanensis, II, 135, 158
 pomanensis grossei, II, 135
 portoricensis, II, 150
 poselgeri, II, 111
 poselgerianus, III, 20
 pottsii, II, 112
 princeps, II, 123; IV, 276
 pringlei, II, 68, 69, 70
 prismaticus, II, 123, 212
 prismatiformis, II, 14
 procumbens, III, 22; IV, 285
 propinquus, III, 21, 22
 propinquus subarticulatus,
 IV, 283
 proteiformis, III, 238
 pruinatus, II, 21
 pruinatus, II, 88, 89
 pseudosonorensis, II, 169
 pteranthus, II, 200
 pterocaulis, IV, 237
 pterogonus, II, 201, 213
 pugionifer, II, 96
 pugioniferus, II, 179, 180
 pugioniferus quadranguli-
 spinus, II, 180
 pulchellus, III, 33
 purpusii, II, 184
 pycnacanthus, II, 137, 138
 quadrangularis, II, 124
 quadrangulispinus, II, 180
 quadricostatus, II, 8
 queretaroensis, II, 96, 97
 quintero, II, 139
 quisco, II, 137
 radicans, II, 195
 ramosus, II, 123
 ramulosus, IV, 240
 recurvus, III, 142
 reductus, III, 158, 159
 reflexus, II, 43; IV, 266
 regalis, II, 20
 regeli, II, 155; IV, 266
 reichenbachianus, III, 23
 reichenbachianus castaneus,
 III, 25
 repandus, II, 4, 17, 18, 151,
 152, 224; IV, 278
 repandus laetevirens, II, 149
 reptans, II, 8, 195; IV, 290
 resupinatus, II, 87
 retroflexus, II, 43
 rhodacanthus, III, 79

Cereus—continued,

rhodanthus, II, 170
 rhodocephalus, II, 158
 rhodoleucanthus, II, 21, 22;
 IV, 268
 rhombeus, IV, 244
 rigidispinus, II, 103
 rigidissimus, III, 28
 rigidus, II, 210
 robustior, III, 28
 robustus, II, 21
 roemerii, III, 13, 14
 roetteri, III, 31
 roezlii, IV, 268
 rogallii, II, 21
 roridus, II, 89
 rosaceus, II, 201
 roseanus, II, 198
 rostratus, II, 203, 204, 205;
 IV, 283
 rothii, IV, 283
 royeri, II, 44, 30
 royerii armatus, II, 40, 51
 royerii floccosus, II, 51
 ruber, II, 223
 ruferi, II, 219
 ruferi major, II, 219
 ruficeps, II, 75
 rufispinus, III, 23
 russellianus, IV, 184, 185
 russellianus, II, 33
 salm-dyckianus, III, 7
 salmianus, III, 7, 74
 salpingensis, II, 21
 sanborgianus, III, 34
 santiaguensis, II, 131
 sargentianus, II, 177, 178;
 IV, 280
 saxicola, II, 21, 22
 saxicola anguiniformis, II, 22
 scandens, IV, 282
 scandens minor, II, 197; IV,
 282
 scheeri, III, 6
 schelhasii, III, 66
 schenckii, II, 180
 schickendantzii, II, 144
 schmidtii, II, 197
 schoenemannii, II, 21
 schomburgkii, II, 191
 schottii, II, 177, 178; IV,
 280
 schottii australis, II, 177
 schrankii, II, 127
 schumannii, II, 103
 scirius, III, 22
 sclerocarpus, II, 146, 147
 scolopendrii, IV, 188
 scopa, III, 193
 scoparius, II, 41
 seidelii, II, 15
 senilis, II, 27; IV, 268
 sepium, II, 160
 sericeus, I, 73
 serpens, I, 217; II, 163
 serpentinus, II, 20, 117, 118,
 119; IV, 276, 284
 serpentinus albispinus, II, 59,
 118
 serpentinus splendens, II, 21,
 118
 serpentinus stellatus, II, 118
 serpentinus strictior, II, 118
 serratus, II, 129
 serruliflorus, II, 131
 setaceus, II, 211, 212
 setaceus viridior, II, 211
 setiger, II, 129
 setosus, II, 34, 35; IV, 215,
 216
 silvestrii, III, 48, 49; IV, 283
 simonii, II, 169
 sirul, II, 123
 smaragdiflorus, II, 174, 173
 smithianus, IV, 284
 smithii, II, 218; IV, 284
 sonorensis, II, 169

Cereus—continued,

- spachianus, II, 131, 132; IV, 277
 spatulatus, II, 21
 speciosissimus, II, 128 129; IV, 277, 283
 speciosissimus albiflorus, II, 128
 speciosissimus aurantiacus, II, 128
 speciosissimus blindii, II, 128
 speciosissimus bodii, II, 129
 speciosissimus bollwillerianus, II, 128
 speciosissimus bowtrianus, II, 128
 speciosissimus candesii, II, 128
 speciosissimus coccineus, II, 127, 128
 speciosissimus colmariensis, II, 128
 speciosissimus curtisii, II, 128
 speciosissimus danielsii, II, 128
 speciosissimus devauxi, II, 128
 speciosissimus edesii, II, 128
 speciosissimus elegans, II, 128
 speciosissimus eugenia, II, 128
 speciosissimus finkii, II, 128
 speciosissimus gebvillerianus, II, 128
 speciosissimus gloriosus, II, 128
 speciosissimus grandiflorus, II, 128
 speciosissimus guillardieri, II, 128
 speciosissimus hansii, II, 128
 speciosissimus hitchensii, II, 128
 speciosissimus hitchensii hybridus, II, 128
 speciosissimus hitchensii speciosus, II, 128
 speciosissimus hoveyi, II, 128; IV, 277
 speciosissimus ignescens, II, 128
 speciosissimus jenkinsonii, II, 128
 speciosissimus kampmannii, II, 128
 speciosissimus kiardii, II, 128
 speciosissimus kobii, II, 128
 speciosissimus latifrons, II, 128
 speciosissimus lateritius, II, 128; IV, 198
 speciosissimus longipes, II, 128
 speciosissimus lothii, II, 128
 speciosissimus loudonii, II, 128
 speciosissimus macqueanus, II, 128
 speciosissimus maelenii, II, 128
 speciosissimus maurantianus, II, 128
 speciosissimus merckii, II, 128
 speciosissimus mexicanus, II, 128
 speciosissimus mittleri, II, 128
 speciosissimus mulhausianus, II, 128
 speciosissimus peacocki, II, 128

Cereus—continued,

- speciosissimus peintneri, II, 128
 speciosissimus rintzii, II, 128
 speciosissimus roidii, II, 128
 speciosissimus roseus albus, II, 128
 speciosissimus roseus superbus, II, 128
 speciosissimus roydii, II, 128
 speciosissimus sarniensis, II, 128
 speciosissimus seidelii, II, 128
 speciosissimus seitzii, II, 128
 speciosissimus selloi, II, 128
 speciosissimus smithii, II, 128
 speciosissimus superbus, II, 128
 speciosissimus suwaroffii, II, 128
 speciosissimus suwarowii, II, 128
 speciosissimus triumphans, II, 128
 speciosissimus unduliflorus, II, 128
 speciosissimus vitellinus, II, 128
 speciosus, II, 128, 129; IV, 277
 speciosus albiflorus, II, 128
 speciosus coccineus, II, 127
 spegazzinii, II, 23; IV, 268
 spegazzinii hassleri, IV, 268
 spinibarbis, II, 82
 spinibarbis flavidus, II, 144
 spinibarbis minor, II, 139
 spinibarbis purpureus, II, 139
 spinosissimus, II, 11
 spinulosus, II, 207
 splendens, II, 21, 118
 splendidus, IV, 198
 squamosus, II, 173, 176
 squamulosus, IV, 215
 squarrosus, II, 104
 steckmannii, II, 21
 stellatus, II, 92, 93, 169
 stenogonus, II, 4, 9, 10, 11
 stenopterus, IV, 190
 stolonifer, IV, 268
 stramineus, III, 40
 straussii, II, 171, 226; IV, 79
 striatus, II, 22, III
 strictus, II, 44
 strigosus, II, 143, 144
 strigosus intricatus, II, 143
 strigosus longispinus, II, 143
 strigosus rufispinus, II, 144
 strigosus spinosior, II, 144
 subinermis, III, 16
 subintortus, II, 19
 subintortus flavispinus, II, 19
 sublanatus, II, 43; IV, 268
 subrepandus, II, 151
 subsquamatus, II, 191
 subtortuosus, II, 174
 subuliferus, II, 137
 superbus, II, 128
 surinamensis, II, 13
 swartzii, II, 46, 87
 syringacanthus, I, 89
 tacaquirensis, II, 225
 tarijensis, II, 226
 taylorii, II, 153
 tellii, II, 21
 tenellus, II, 126
 tenuis, II, 19; IV, 215
 tenuispinus, IV, 215
 tephacanthus, III, 188; IV, 289
 tephacanthus bolivianus, II, 136
 terscheckii, II, 143

Cereus—continued,

- testudo, II, 213
 tetazo, II, 76; IV, 272
 tetezo, IV, 272
 tetetzo, IV, 272
 tetracanthus, II, 136
 tetragonus, II, 3, 4, 7, 9, 14
 tetragonus major, II, 9
 tetragonus minor, II, 14
 tetragonus ramosior, II, 9
 thalassinus, II, 5, 9
 thalassinus quadrangularis, II, 5
 thelegonoides, II, 131
 thelegonus, II, 130, 131
 thouarsii, II, 120, 121
 thurberi, II, 68, 97, 98, 99; III, 43; IV, 274
 thurberi littoralis, II, 97
 thurberi monstrosus, II, 98
 tilophorus, II, 43; IV, 268
 tinei, II, 152
 titan, II, 69, 70
 tonduzii, II, 77, 26; IV, 284
 tonelianus, II, 92; IV, 290
 torrellianus, IV, 290
 tortuosus, II, 154, 155
 tortus, II, 83
 treleasei, II, 93
 triangularis, II, 187, 188, 192, 193; IV, 282
 triangularis major, II, 187, 191
 triangularis pictus, II, 192
 triangularis undeanus, II, 193
 trichacanthus, II, 44
 trichocentrus, II, 21
 tricoatus, II, 187
 triglochidiatus, III, 10
 trigonodendron, II, 19; IV, 267
 trigonus, II, 186, 192
 trigonus costaricensis, II, 186
 trigonus guatemalensis, II, 184
 trigonus quadrangularis, II, 124
 trinitatensis, II, 189
 tripteris, IV, 268
 triqueter, IV, 282
 truncatus, IV, 177
 truncatus altensteinii, IV, 178
 tuberosus, II, 111; IV 8, 274
 tunicatus, I, 66
 tubiflorus, III, 67
 tunilla, II, 214
 tupizensis, II, 226
 turbinatus, III, 66
 tweediei, II, 174; IV, 279
 ulei, II, 52
 undatus, II, 149, 151, 152, 187
 undiflorus, IV, 275
 undulatus, II, 124
 undulosus, II, 123
 uranos, II, 197
 uranus nycticalus, II, 197
 urbanianus, II, 43, 198; IV, 283
 ureacanthus, II, 158
 uspensi, IV, 268
 vagans, II, 205
 validissimus, III, 63
 validus, II, 4, 7
 variabilis, II, 4, 13, 14, 123, 124
 variabilis glaucescens, II, 14
 variabilis gracilior, II, 14
 variabilis laetevirens, II, 14
 variabilis micracanthus, II, 14
 variabilis obtusus, II, 14
 variabilis ramosior, II, 14
 variabilis salm-dyckianus, II, 14

Cereus—continued,

- vasmeri, II, 123
 vaupellii, II, 202
 venditus, II, 192
 ventimiglae, II, 160
 verschaffeltii, II, 21
 victoriensis, II, 53
 violaceus, II, 44
 viperinus, II, 110
 virens, II, 43
 viridiflorus, III, 17, 18
 viridiflorus cylindricus, III, 17
 viridiflorus minor, IV, 285
 viridiflorus tubulosus, III, 17
 vulcan, II, 219
 vulnerator, IV, 266
 warmingii, II, 42
 weberbaueri, II, 141
 weberi, II, 95, 96
 weingartianus, II, 77, 78
 wercklei, II, 208
 wittii, II, 221, 222
 xanthocarpus, II, 4, 10
 xanthochaetus, II, 225
 ziczkaanus, II, 224
 zizkaanus, II, 224
 Chacoub, II, 187
 Chaetophorae, I, 174
 Chaffeyanae, I, 45, 213
 Chalote, III, 84
 Chamaecereus, III, 3, 48, 49
 silvestrii, III, 48; IV, 285
 Chaute, III, 8
 Chende, II, 91
 Chente, II, 91
 Chiapas, IV, 177, 203, 204
 nelsonii, IV, 203
 Chichibe, II, 90
 Chichipe, II, 90
 Chichituna, II, 90
 Chilotos, III, 93
 Chino, II, 91
 Chiotilla, II, 66
 Chique-chique, II, 35
 Chique-chique das pedras, II, 35
 Chirinola, II, 115
 Cholla, I, 43, 61
 Christmas cactus, IV, 178
 Cina, II, 169
 Clavarioides, I, 44, 72
 Clavatae, I, 4, 79, 84
 Cleistocactus, II, 2, 173-176; III, 60, 78, 79
 anguinus, II, 173, 175
 areolatus, II, 159
 aureus, II, 105
 baumannii, II, 163, 173, 174, 175, 226; IV, 279
 baumannii colubrinus, II, 174
 baumannii flavispinus, II, 174
 celsianus, II, 171
 chotaensis, II, 163
 colubrinus, II, 174
 humboldtii, II, 163
 hyalacanthus, II, 176
 icosagonus, II, 160
 kerberi, II, 170
 lanatus, II, 61
 laniceps, II, 175
 monvilleanus, II, 173
 parviflorus, II, 176
 parvisetus, II, 175
 rhodacanthus, III, 79
 sepium, II, 160
 serpens, II, 163
 smaragdiflorus, II, 173, 174, 175
 Cob cactus, III, 37
 Cochal, II, 180
 Cochemia, IV, 3, 19, 21-23, 65
 halei, IV, 21, 22
 hallei, IV, 22
 pondii, IV, 21, 23

- Cochemia*—*continued*,
 poselgeri, IV, 21, 22
 rosiana, IV, 22
 setispina, TV, 21, 22
Coerulescentes, II, 3, 59
 Cola de diablo, I, 26
Compresso-costati, II, 3
Consolea, I, 42, 43, 202
 catacantha, I, 208
 ferox, I, 206
 leucacantha, I, 175
 rubescens, I, 43, 208
 spinosissima, I, 204
Copado, II, 83
Copiapoa, III, 77, 85-90, 100
 cinerascens, III, 85, 88
 cinerea, III, 85, 86
 coquimbana, III, 85, 87
 echinoides, III, 85, 88
 marginata, III, 85, 86
 megarrhiza, III, 85, 89
Corotilla, I, 96
Corryocactus, II, 2, 66-68
 brachypetalus, II, 66, 67, 68
 brevistylus, II, 66, 67, 68
 melanotrichus, II, 66, 68
Coryphantha, III, 3, 45, 148;
 IV, 3, 4, 13, 23-51, 53,
 55, 57, 65, 165, 170, 175
 acanthostephes, IV, 41
 aggregata, IV, 25, 47, 154
 ancistracantha, IV, 15
 arizonica, IV, 25, 45
 aulacothele, IV, 30, 31
 brevimamma, IV, 31
 bumamma, IV, 24, 29, 33, 51
 calacarata, IV, 48
 chlorantha, IV, 25, 43, 44, 47
 clava, IV, 24, 30
 compacta, IV, 24, 36, 37
 connivens, IV, 24, 34
 conoidea, IV, 17
 conspicua, IV, 51
 cornifera, IV, 18, 25, 39
 cubensis, IV, 23, 25, 48
 daimonoceras, IV, 36
 dasyacantha, IV, 55
 deserti, IV, 25, 46
 durangensis, IV, 25, 42, 43
 echinoidea, IV, 24, 30
 echinus, IV, 25, 35, 42
 elephantidens, IV, 24, 32,
 33, 50
 engelmännii, IV, 27, 51
 erecta, IV, 24, 32
 exsudans, IV, 24, 31
 glanduligera, IV, 31
 grahamii, IV, 155
 guerkeana, IV, 24, 29
 heteromorpha, IV, 25
 heterophylla, IV, 25
 hookeri, IV, 51
 impexicoma, IV, 37
 lehmanni, IV, 30, 31
 loricata, IV, 168
 macromeris, IV, 23, 24, 25
 missouriensis, IV, 53
 muchlenpfordtii, IV, 24, 28,
 29, 34
 neo-mexicana, IV, 25, 44, 45
 nickelsae, IV, 23, 24, 35, 36
 nivosa, IV, 71
 nuttallii, IV, 53
 octacantha, IV, 24, 30
 ottonis, IV, 24, 26
 pallida, IV, 25, 40
 palmeri, IV, 25, 39
 pectinata, IV, 24, 34, 35, 175
 poselgeriana, IV, 24, 2, 28, 29
 pottsii, IV, 137
 pycnacantha, IV, 25, 40, 41
 radians, IV, 24, 36, 37
 radiosa, IV, 43, 60
 raphidacantha, IV, 15
 recurvata, IV, 24, 27
 retusa, IV, 24, 38
 raphidacantha, IV, 15
Coryphantha—*continued*,
 robustispina, IV, 24, 33
 runyonii, IV, 24, 26
 salm-dyckiana, IV, 25, 38, 39
 scheeri, IV, 28
 schlechtendalii, IV, 30
 similis, IV, 52
 sublanata, IV, 51
 sulcata, IV, 25, 48, 49
 sulcolanata, IV, 24, 37, 50
 tuberculosa, IV, 54
 vivipara, IV, 25, 43, 44, 45,
 47
Coryphanthae, IV, 53
Coryphanthanae, II, 1; III, 77,
 90, 146, 149; IV, 3, 6,
 57, 62
 Cow's tongue, I, 164
 Crab cactus, IV, 178, 186
 Creeping devil cactus, II, 115
 Criniferae, I, 140, 176
 Crispatae, IV, 221
 Cruciformes, I, 208
 Cubenses, IV, 25
 Cuija, I, 149
 Curassavicae, I, 45, 102, 104,
 106, 193
 Curuntilla, IV, 266
 Cuscuta baccifera, IV, 227
 Cycad, III, 108
 Cylindropuntia, I, 32, 44, 45,
 46, 71, 75, 79, 84, 100, 142
 Daatoc, II, 88
 Deamia, II, 183, 212-214
 testudo, II, 209, 213, 214
 Deerhorn cactus, II, 113
 Dendrocreus, II, 2, 113, 114;
 IV, 281
 nudiflorus, II, 113, 114; IV,
 276
 Denmoza, III, 77, 78, 79
 rhodacantha, III, 79
 Devil's bead cactus, III, 182
 Devil's pincushion, III, 182
 Devil's root, III, 84
 Dildo, II, 224
 Dildoes, I, 105
 Dillenianae, I, 45, 159, 169
 Discocactus, III, 216-220; IV,
 70, 165
 alteolens, III, 216, 217, 218
 bahiensis, III, 217, 220
 besleri, III, 219
 hartmannii, III, 217
 heptacanthus, III, 217, 218
 insignis, III, 216, 219, 220
 lehmannii, III, 219
 linkii, III, 219
 placentiformis, III, 216, 217,
 218, 219; IV, 70, 165
 subnudus, III, 216, 217
 tricornis, III, 218
 zehntneri, III, 217, 218, 220
 Disisocactus, IV, 201
 biformis, IV, 202
 Disocactus, IV, 177, 201-203
 biformis, IV, 201, 202
 eichlamii, IV, 202, 203
 Dissimiles, IV, 221
 Diurna, IV, 186
 Dolichothele, IV, 3, 61-64, 6
 longimamma, IV, 61, 62, 63
 sphaerica, IV, 61, 62
 uberiformis, IV, 61, 63, 64
 Dumpling cactus, III, 84
 Durasnilla, I, 175
 Eccremocactus, IV, 177, 204
 bradei, IV, 204
 Echinocactanae, II, 1; III, 3, 77-
 215; IV, 24, 57, 175, 274
 Echinocacti, IV, 6, 34, 58
 Echinocactus, I, 117; II, 1, 105,
 137, 171; III, 3, 32, 45, 48,
 49, 55, 60, 77, 78, 83, 85,
 90, 92, 93, 94, 98, 100,
 101, 104, 108, 109, 118,
 123, 128, 140, 142, 148,
 Echinocactus—*continued*,
 151, 152, 166-181, 186,
 187, 205, 206, 207, 211,
 216, 237; IV, 3, 4, 6, 10,
 12, 16, 24, 28, 29, 58,
 137, 165, 175, 288
 acanthion, III, 98, 120
 acanthodes, III, 129
 acanthostephes, IV, 41
 aciculatus, III, 198
 acifer, III, 111
 acifer spinosus, III, 111
 acroacanthus, III, 120
 acroacanthus, III, 121
 acuatus, III, 188
 acuatus arechavaletai, III,
 200
 acuatus corynodes, III, 198
 acuatus depressus, III, 198
 acuatus erinaceus, III, 198
 acuatus sellowii, III, 188
 acuatus spinosior, III, 188
 acuatus tetraacanthus, III,
 188
 acutangulus, III, 199
 acutatus, III, 188
 acutispinus, III, 179; IV, 288
 acutissimus, III, 97, 98, 99,
 103
 acutissimus cristatus, III, 98
 adversispinus, III, 121
 agglomeratus, III, 136, 143
 alamosanus, III, 137
 albatu, III, 112
 allardtianus, III, 114
 alteolens, III, 218
 amazonicus, III, 175
 ambiguus, III, 88
 ancylocanthus, III, 146
 anfractuus, III, 117, 123
 anfractuus ensiferus, III,
 114
 anfractuus laevior, III, 116
 anfractuus orthogonus, III,
 157
 anfractuus pentacanthus,
 115, 116
 anfractuus spinosior, III,
 117
 anisitsii, III, 159, 161
 apricus, III, 192
 arachnoideus, III, 176
 araneifer, III, 177
 araneolarius, III, 176
 arcuatus, III, 188
 arechavaletai, III, 196, 197,
 200, 201
 arizonicus, III, 127, 128
 arizonicus, III, 176
 armatus, III, 178, 238
 arrectus, III, 114
 arrigens, III, 114
 arrigens atropurpureus, III,
 114
 asterias, III, 183, 184
 aulacogonus, III, 169, 170
 aulacogonus diacopaulax, III,
 169
 aurantiacus, III, 102
 auratus, II, 143; III, 186
 aureus, II, 105; III, 168
 baldianus, III, 163, 164
 barcelona, IV, 287
 beguinii, III, 148, 149; IV,
 16
 berterii, III, 7
 biceras, III, 115
 bicolor, III, 43, 105; IV,
 11, 12
 bicolor bolansii, IV, 11
 bicolor montemorelanus, IV,
 12
 bicolor pottsii, IV, 12
 bicolor schottii, IV, 11
 bicolor tricolor, IV, 11
 bolansii, IV, 11, 12
 bolivianus, III, 88
Echinocactus—*continued*,
 brachiatus, III, 121
 brachyanthus, III, 159; IV,
 288
 brachycentrus, III, 121
 brachycentrus olygacanthus,
 III, 121
 brevihamatus, IV, 4, 5
 brevimmamus, IV, 31
 bridgesii, III, 88, 89
 buchheimianus, IV, 288
 buckii, IV, 8
 bukkii, IV, 8
 cachensis, III, 53
 cachetianus, III, 105, 106
 cachetianus orcuttii, III, 105
 caespitius, IV, 16
 caespitosus, III, 208, 211
 californicus, III, 130, 141
 campylacanthus, III, 140
 candicans, II, 143
 capricornis, III, 184, 185;
 IV, 288
 capricornis major, III, 185
 capricornis minor, III, 184,
 185
 castaneoides, III, 99
 castaniensis, III, 179; IV, 12
 catamarcensis, II, 146; III,
 197
 catamarcensis obscurus, III,
 197
 catamarcensis pallidus, III,
 197
 cataphractus, III, 208, 210
 celsianus, III, 158
 centetierius, III, 203, 204
 centetierius grandiflorus, III,
 203, 204
 centetierius major, III, 203,
 204
 centetierius pachycentrus, III,
 203
 cephalophorus, IV, 41
 ceratistes, II, 143; III, 176,
 178, 186
 ceratistes celsii, III, 186
 ceratistes melanacanthus, III,
 186
 ceratistes, III, 187
 ceratistes, III, 186
 cerebriformis, III, 179
 cereiformis, III, 121
 chereauianus, III, 54
 chilensis, III, 99
 chilensis confinis, III, 99
 chionanthus, III, 49, 58
 chlorophthalmus, III, 32
 chrysacanthion, III, 176
 chrysacanthus, III, 127
 cinerascens, III, 59, 87, 88, 89
 cinereus, III, 86
 cinnabarinus, III, 54, 59
 cinnabarinus spinosior, III,
 54
 clavatus, III, 99, 101
 clavus, IV, 30
 coccineus, III, 79
 columnaris, III, 86, 87
 concinnus, III, 192, 193,
 208; IV, 288
 concinnus joadii, III, 192
 concinnus tabularis, III, 193
 confertus, III, 179
 conglomeratus, III, 39, 88
 conimamma, IV, 38
 conoideus, IV, 17
 conothelos, IV, 13
 conquades, III, 199
 contractus, III, 161
 copiapensis, III, 88, 89, 186
 copoldi, III, 130
 coptonogonus, III, 109, 110,
 111
 coptonogonus major, III, 110
 coptonogonus obvallatus, III,
 115

- Echinocactus*—*continued*,
coquimbatus, III, 87
corniferus, IV, 13, 39
corniferus impexicomus, IV, 36
corniferus longisetus, IV, 39
corniferus muticus, IV, 39
corniferus nigricans, IV, 39
corniferus raphidacanthus, IV, 15
corniferus scolymoides, IV, 39
corniger flavispinus, III, 143
corniger rubrispinosus, III, 143
cornigerus, III, 116, 142, 143; IV, 287
cornigerus flavispinus, III, 143; IV, 287
cornigerus latispinus, III, 143; IV, 287
corrugatis, III, 179
corynacanthus, III, 168
corynodes, III, 187, 198, 199; IV, 289
corynodes erinaceus, III, 198
coulteri, III, 138
courantianus, III, 182
courantii, III, 188
courantii spinosior, III, 188
coxii, III, 197
crassihamatus, III, 144
crenatus, III, 98
criocereus, III, 180
crispatus, III, 115, 116, 117
crispatus cristatus, III, 117
crispatus horridus, III, 116
ctenoides, III, 19
cumingii, III, 59
cumingii flavispinus, III, 39
cumingii, III, 99
cupreatus, III, 96, 176
cupulatus, III, 176
curvicornis, III, 142
curvispinus, III, 100, 203
cylindraceus, III, 129, 130
cylindraceus albispinus, III, 130
cylindraceus longispinus, III, 130
cylindricus, IV, 288
dadaki, III, 179
damsii, III, 163
darrahi, IV, 288
debilispinus, III, 121
decaisnei, III, 66
deflexispinus, III, 145
delaetii, III, 164, 165
deminutus, III, 48
densus, IV, 134
denudatus, III, 155; IV, 288
denudatus andersohnianus, III, 155
denudatus bruennowianus, III, 163
denudatus bruennowii, III, 156
denudatus delaetianus, III, 155
denudatus delaetii, III, 155
denudatus flavispinus, III, 156
denudatus golzianus, III, 155
denudatus heuschkehlai, III, 156
denudatus heuschkelianus, III, 155, 156
denudatus intermedius, III, 156
denudatus meiklejohnianus, III, 155
denudatus multiflorus, IV, 288
denudatus octogonus, III, 155
- Echinocactus*—*continued*,
denudatus paraguayensis, III, 156
denudatus roseiflorus, III, 156
denudatus scheidelianus, III, 155
denudatus typicus, III, 155
denudatus wagnerianus, III, 153
denudatus wiedzianus, III, 155
deppii, IV, 285
depressus, III, 576, 235
dichroacanthus, III, 117
dichroacanthus spinosior, III, 117
dicracanthus, IV, 288
dietrichianus, III, 137
dietrichii, III, 118, 119
diguettii, III, 131, 132
disciformis, III, 106
dolichacanthus, III, 136
dolichocentrus, III, 136
drageanus, IV, 10
droegeanus, IV, 10
dumesnilianus, III, 79
durangensis, III, 152
ebenacanthus, III, 99
ebenacanthus affinis, III, 99
ebenacanthus intermedius, III, 99
ebenacanthus minor, III, 99
echidna, III, 136
echidne, III, 136
echidne gilvus, III, 136
echinatus, III, 177
echinoides, II, 138; III, 88, 89
echinoides pepinianus, II, 137
edulis, III, 170
ehrenbergii, IV, 9
elachisanthus, III, 205
electracanthus, III, 139
electracanthus capuliger, III, 138
electracanthus haematacanthus, III, 147
electracanthus rufispinus, III, 139
elegans, II, 139
elephantidens, IV, 32
ellemeetii, III, 121
ellipticus, IV, 11, 12
emoryi, III, 127, 128, 132, 133, 134
emoryi chrysacanthus, III, 127
emoryi rectispinus, III, 134
engelmannii, III, 38
ensiferus, III, 114
ensiferus pallidus, III, 114
equitans, III, 175
erectocentrus, III, 148, 149; IV, 16, 287
erectus, IV, 32
erinaceus, III, 198, 199, 200
erinaceus elatior, III, 199
escayachensis, III, 205
exsculptus, III, 97, 98, 110
exsculptus cristatus, III, 98
exsculptus dichroacanthus, III, 98
exsculptus elatior, III, 98
exsculptus foveolatus, III, 98
exsculptus fulvispinus, III, 98
exsculptus gayanus, III, 98
exsculptus tenuispinus, III, 98
exsculptus thrincogonus, III, 98
eyriesii, III, 60, 65, 66, 142
eyriesii glaucus, III, 66
falconeri, III, 127, 128
famatimensis, IV, 286
- Echinocactus*—*continued*,
farinosus, II, 19
fascicularis, II, 141
fennellii, III, 158
ferox, III, 158
fiebrigii, III, 46, 47
fiedlerianus, III, 87
fischeri, III, 178, 238
flavescens, III, 138
flavicomma, III, 179
flavispinus, III, 143, 145
flavovirens, III, 138, 147
flexispinus, III, 117, 144
flexuosus, III, 121
floricomus, III, 200, 201
fluctuosus, III, 121
fobeanus, III, 89
foersteri, III, 121
foliosus, III, 179
forbesii, III, 62
fordii, III, 126, 127
formosus, III, 3
formosus crassispinus, III, 75
fossulatus, IV, 6, 10
foveolatus, III, 98
fricii, III, 188, 189
froehlichianus, III, 203
fuscus, III, 99
galeottii, III, 168
gayanus, III, 98
gayanus intermedius, III, 98
geissei, III, 177
geissei albicans, III, 177
gemmaus, III, 66
gerardii, III, 147
ghiesbreghtii, II, 60; III, 185
gibbosus, III, 152, 158
gibbosus celsianus, III, 158
gibbosus cerebriformis, III, 158
gibbosus chubutensis, III, 158
gibbosus fennellii, III, 158
gibbosus ferox, III, 158
gibbosus leonensis, III, 158
gibbosus leucacanthus, III, 158
gibbosus leucodictyus, III, 158
gibbosus nobilis, III, 158, 159
gibbosus platensis, III, 163
gibbosus pluricostatus, III, 158
gibbosus polygonus, III, 158
gibbosus schlumbergeri, III, 158
gibbosus typicus, III, 158
gibbosus ventanicola, III, 158
gigas, III, 179
gilliesii, III, 75
gilvus, III, 136
glabrescens, III, 180
gladiatus, III, 119, 120, 123
gladiatus intermedius, III, 119
gladiatus ruficeps, III, 119
glanduligerus, IV, 31
glaucescens, III, 137
glaucus, III, 143, 213
globosus cristatus, III, 138
gracilis, III, 209
gracillimus, III, 209, 211
graessneri, III, 203
grahlianus, III, 209
grahlianus adustior, III, 209
grandicornis, III, 114
grandicornis fulvispinus, III, 114
grandicornis nigripinus, III, 114
grandis, III, 167, 169, 171
griseispinus, III, 122
grossel, III, 190
grusonii, III, 167, 168; IV, 288
grusonii azureus, III, 68
- Echinocactus*—*continued*,
guerkeanus, III, 134, 153
guyannensis, III, 98
haageanus, III, 77, 170, 185
haematacanthus, III, 147
haematanthus, III, 37
haematochroanthus, III, 145
hamatacanthus brevispinus, III, 143
hamatacanthus longihamatus, III, 143
hamatocanthus, III, 144
hamatus, III, 104, 177
hamulosus, III, 104
hankeanus, III, 99
hartmannii, III, 217
haselbergii, III, 201, 202; IV, 289
haselbergii cristatus, III, 202
hastatus, III, 111
hastatus fulvispinus, III, 111
haynei, III, 102, 103
bayou, III, 102, 103
helianthoidiscus, III, 107
helophorus, III, 169
helophorus laevior, III, 169
helophorus longifossulatus, III, 169
hemifossus, III, 180
hemifossus gracilispinus, III, 180
hemipelianus, III, 101
heteracanthus, III, 112, 117
heterochromus, IV, 12, 13
heteromorphus, IV, 23
hexacanthus, III, 122
hexaedrophorus, IV, 6, 7, 10
hexaedrophorus droegeanus, IV, 10
hexaedrophorus fossulatus, IV, 10
hexaedrophorus labouretianus, IV, 6
hexaedrophorus major, IV, 6
hexaedrophorus roseus, IV, 6
hexaedrophorus subcostatus, IV, 10
hexaedrus, IV, 13
heyderi, III, 122
heynei, III, 10
histrix, III, 138
hoffmanseggii, III, 98
holopteris, III, 185
hookeri, III, 122
horizontalis, III, 173
horizontalionus, III, 100, 167, 175, 177
horizontalionus centrispinus, III, 175
horizontalionus curvispinus, III, 173
horizontalionus obscurispinus, III, 173
horridus, III, 202
horripilus, IV, 16
horripilus erectocentrus, III, 149
horripilus longispinus, IV, 16
humilis, III, 89, 99
huottii, III, 63
hybocentrus, III, 203, 204
hybognus, III, 157
hybognus saglionis, III, 157
hylainacanthus, III, 179
hypocrateriformis, III, 200
hypocrateriformis spinosior, III, 200
hyptiacanthus, III, 154, 156, 157
hyptiacanthus eleutheracanthus, III, 156
hyptiacanthus megalotelus, III, 156
hyptiacanthus nitidus, III, 156
hystrichacanthus, III, 139
hystrichocentrus, III, 122

Echinocactus—*continued*,
 hystrioides, III, 122
 hystrix, II, 86; III, 138, 170
 inflatus, IV, 288
 ingens, III, 18, 167, 168,
 169, 170, 171; IV, 288
 ingens edulis, III, 170
 ingens grandis, III, 170
 ingens helophorus, III,
 ingens irroratus, III, 169
 ingens saltillensis, III, 172;
 IV, 28
 ingens subinermis, III, 170,
 172, 173
 ingens visnaga, III, 170, 171
 insculptus, IV, 7
 insignis, III, 145; IV, 287
 intermedius, III, 155
 interruptus, III, 98, 110
 intertextus, III, 149, 150,
 197
 intertextus dasyacanthus, III,
 149, 150; IV, 55
 intortus, III, 230
 intortus purpureus, III, 226
 intricatus, III, 180, 206
 intricatus longispinus, III,
 88
 irroratus, III, 170
 islayensis, III, 201
 joadi, III, 192, 193
 jeneschianus, II, 138
 johnsonii, III, 141
 johnsonii octocentrus, III,
 141
 joossensianus, III, 166
 jordanianus, III, 85
 juniori, III, 180
 juori, III, 180
 jussianus, III, 96
 jussieui, III, 96, 97
 jussieui cristatus, III, 96
 karwinskius, III, 169
 karwinskii, III, 169
 knippelianus, III, 211
 krausei, III, 150
 kunzei, III, 99
 kunzei brevispinus, III,
 100
 kunzei rigidior, III, 100
 kunzii, III, 99
 kurtzianus, III, 163
 labouretianus, IV, 7
 lamellosus, III, 113
 lamellosus fulvescens, III,
 114
 lancifer, III, 115, 118, 119,
 138
 langsdorfii, III, 199, 200
 laticostatus, III, 175
 latispinus, III, 180
 latispinus, III, 143
 latispinus flavispinus, III,
 143
 lecchii, II, 20
 lecomtei, III, 129
 lecontei, III, 129; IV, 287
 lecontei albispinus, III, 129
 lecontei hagei, III, 129
 lecontei, III, 129; IV, 256
 lecanus, III, 154
 lemarii, III, 226
 leninghausii, III, 204, 205;
 IV, 289
 leninghausii cristatus, III,
 205
 leonensis, III, 21, 158
 leopoldi, III, 130; IV, 287
 leucacanthus, III, 225; IV,
 8, 9
 leucacanthus crassior, IV, 8
 leucacanthus tuberosus, IV, 8
 leucanthus, III, 72
 leucocarpus, III, 198, 199
 leucodictyus, III, 158
 leucotrichus, III, 100, 101
 lewinii, III, 84
 limitus, III, 140

Echinocactus—*continued*,
 lindheimeri, III, 181
 lindleyi, III, 90
 linkeanus, III, 122
 linkii, III, 195; IV, 289
 linkii spinosior, III, 195
 longihamatus, III, 117, 144,
 145, 146; IV, 287
 longihamatus bicolor, III, 145
 longihamatus brevispinus, III,
 145
 longihamatus crassispinus,
 III, 145
 longihamatus deflexispinus,
 III, 145
 longihamatus gracilispinus,
 III, 144
 longihamatus hamatacanthus,
 III, 144
 longihamatus insignis, III,
 145
 longihamatus sinuatus, III,
 145, 146
 longispinus, III, 180
 lophothele, IV, 7, 8
 lophothele longispinus, IV, 8
 loratus, III, 155; IV, 168
 maassii, III, 202
 macdowellii, III, 151
 mackieanus, III, 158, 159
 macleanii, IV, 8
 macracanthus, III, 89, 169
 macracanthus cinerascens,
 III, 89
 macrocephalus, III, 122
 macrodiscus, III, 139, 140
 macrodiscus decolor, III, 139
 macrodiscus laevior, III, 139
 macrodiscus multiflorus, III,
 139, 140
 macromeris, IV, 25
 macrothele, IV, 30
 macrothele biglandulosus, IV,
 30
 macrothele lehmanni, IV, 30
 maelenii, IV, 8
 malletianus, III, 100, 177
 mammosus, III, 180
 mammosus, III, 122, 137
 mammosus, III, 138, 203,
 204
 mammosus, III, 200
 mammosus cristatus, III,
 200
 mammosus hircinus, III,
 200
 mammosus minor, III, 200
 mammosus pampeanus, III,
 200
 mammosus spinosior, III,
 200
 mammosus submammosus,
 III, 200
 mammosus typicus, III,
 200
 marginatus, III, 85, 86, 87
 marisianus, III, 105
 martinii, III, 189
 mathsonii, III, 144
 mcdowellii, III, 151
 megalothelos, III, 87, 162
 megarrhizus, IV, 3, 4
 melanacanthus, III, 186
 melanocarpus, III, 161, 162
 melanochnus, III, 86
 melmsianus, III, 117
 melocactiformis, III, 138,
 139
 melocactoides, III, 224
 meonacanthus, III, 224
 merbellii ornatus, III, 185
 merckeri, III, 180
 micracanthus, III, 180
 micromeris, III, 93
 microspermus, III, 176, 207,
 205; IV, 289
 microspermus brevispinus,
 III, 208

Echinocactus—*continued*,
 microspermus elegans, III,
 208
 microspermus erythranthus,
 III, 208
 microspermus macrancistrus,
 III, 207, 208
 microspermus thionanthus,
 III, 208
 mihanovichii, III, 153, 151
 minax, III, 169
 minax laevior, III, 169
 minusculus, III, 45, 46, 176;
 IV, 285
 minusculus cristatus, III, 46
 mirbeli, III, 185
 mirbeli ornatus, III, 185
 misleyi, III, 73
 mitis, III, 181
 molendensis, III, 201
 montevidensis, III, 180
 monvillei, III, 155, 161; IV,
 288
 mostii, III, 158
 muchlenpfordti, III, 104; IV,
 28, 29
 multangularis, III, 142
 multicosatus, III, 111
 multiflorus, III, 155, 156, 159
 multiflorus albispinus, III,
 159
 multiflorus hybopleurus, III,
 159
 multiflorus parisiensis, III,
 159
 multiplex, III, 64; IV, 286
 multiplex cristatus, IV, 286
 muricatus, III, 194, 195
 muricatus hortatani, III, 196
 mutabilis, III, 177
 myriacanthus, III, 101
 myriostigma, III, 182, 183
 myriostigma amabile, III,
 183
 myriostigma amabilis, III,
 183
 myriostigma amoenus, III,
 183
 myriostigma bedinghausi, III,
 153
 myriostigma bedinghausii,
 III, 183
 myriostigma begunii, III,
 183
 myriostigma bellus, III, 153
 myriostigma candidus, III,
 183
 myriostigma cereiformis, III,
 183
 myriostigma cinerascens, III,
 183
 myriostigma cinerascens
 brevispinus, III, 183
 myriostigma cinerascens
 crassispinus, III, 183
 myriostigma cinerascens
 longispinus, III, 183
 myriostigma cinerascens
 parvimaculatus, III, 183
 myriostigma columnaris, III,
 182, 183
 myriostigma conspicuum, III,
 183
 myriostigma conspicuum, III,
 183
 myriostigma cornutus, III,
 183
 myriostigma cornutus candi-
 dus, III, 183
 myriostigma crenatus, III,
 183
 myriostigma darrahii, III,
 183
 myriostigma delaeti, III, 183
 myriostigma diadematus, III,
 183
 myriostigma elegantissimus,
 III, 183

Echinocactus—*continued*,
 myriostigma erectus, III, 183
 myriostigma formosus, III,
 183
 myriostigma gardei, III, 183
 myriostigma glabrescens, III,
 183
 myriostigma hanburyi, III,
 183
 myriostigma hybridus, III,
 183
 myriostigma imperiale, III,
 183
 myriostigma imperialis, III,
 183
 myriostigma incanus, III,
 183
 myriostigma incomparabilis,
 III, 183
 myriostigma inermis, III, 183
 myriostigma insignis, III,
 183
 myriostigma jusberti, III,
 183
 myriostigma lapaixi, III, 183
 myriostigma lapaixii, III,
 183
 myriostigma laurani, III, 183
 myriostigma lesaunieri, III,
 183
 myriostigma lophothele, III,
 183
 myriostigma lophothele ce-
 reiformis, III, 183
 myriostigma martini, III,
 183
 myriostigma mirabile, III,
 183
 myriostigma mirabilis, III,
 183
 myriostigma nobilis, III, 183
 myriostigma nudus, III, 182,
 183
 myriostigma octagonus, III,
 183
 myriostigma octogonum, III,
 183
 myriostigma pentagonus, III,
 183
 myriostigma pictus, III, 183
 myriostigma princeps, III,
 183
 myriostigma quadratus, III,
 183
 myriostigma rebuti, III, 183
 myriostigma regale, III, 183
 myriostigma regalis, III, 183
 myriostigma regulare, III,
 183
 myriostigma regulare spino-
 sum, III, 183
 myriostigma robustum, III,
 183
 myriostigma schilinskyi, III,
 183
 myriostigma sehlinzkyi, III,
 183
 myriostigma schumannii, III,
 183
 myriostigma speciosus, III,
 183
 myriostigma spectabilis, III,
 183
 myriostigma spiralis, III, 183
 myriostigma splendens, III,
 183
 myriostigma variegatus, III,
 183
 myriostigma weberi, III, 183
 myriostigma zonatus, III,
 183
 napinus, III, 191
 netrelianus, III, 154
 neumannianus, III, 100
 neumannianus rigidior, III,
 100
 nidulans, IV, 9, 10
 nidus, III, 94, 207

Echinocactus—*continued*,
 niger, III, 96
 nigricans, III, 95, 96, 176
 nigripinus, III, 190, 191
 nobilis, III, 158
 nodosus, III, 17, 105
 nummularioides, III, 203
 obrepandus, III, 73
 obvallatus, III, 115, 116, 119
 obvallatus pluricostatus, III, 115
 obvallatus spinosior, III, 115
 occultus, III, 95
 ochroleucus, III, 122
 octacanthus, III, 122
 octogonus, III, 6
 odieri, III, 177
 odieri magnificus, III, 178
 odieri mebbesi, III, 178
 odieri spinis nigris, III, 178
 odierianus, III, 177
 olacogonus, III, 180
 oligacanthus, III, 121, 170
 orcuttii, III, 134, 135
 oreptilis, III, 180
 ornatus, III, 185
 ornatus glabrescens, III, 185
 ornatus mirbelii, III, 185
 orthacanthus, III, 138
 ottonianus, IV, 26
 ottonis, III, 142, 195, 196;
 IV, 289
 ottonis brasiliensis, III, 196
 ottonis linkii, IV, 289
 ottonis minor, III, 196
 ottonis pallidior, III, 195
 ottonis paraguayensis, III, 196
 ottonis pfeifferi, III, 196
 ottonis spinosior, III, 195
 ottonis tenuispinus, III, 195,
 196; IV, 289
 ottonis tortuosus, III, 195
 ottonis uruguayus, III, 196
 ourselianus, III, 159
 ourselianus albispinus, III,
 159
 oxyacanthus, III, 180
 oxygonus, III, 64, 65
 oxypterus, III, 138
 pachycentrus, III, 203, 204
 pachycornis, III, 178
 palmeri, III, 167, 172; IV, 28
 pampeanus, III, 200, 201
 pampeanus charruanus, III,
 200
 pampeanus rubellianus, III,
 200
 pampeanus subplanus, III,
 200
 papyracanthus, III, 91, 92
 paraguayensis, III, 156
 parryi, III, 175
 parvispinus, III, 237
 pauciareolatus, III, 188, 189
 pectinatus, III, 29
 pectiniferus, III, 29, 30
 pectiniferus laevior, III, 29
 pelachicus, III, 180
 peninsulae, III, 133, 134
 pentacanthus, III, 115, 116
 pentlandii, III, 49, 54, 55
 pepinianus, II, 137; III, 100
 pepinianus affinis, III, 89
 pepinianus echinoides, II,
 138
 peruvianus, III, 102
 pfeifferi, III, 137, 138
 pferdsdorffii, III, 139
 philippi, II, 105
 phoeniceus inermis, III, 14
 phyllacanthoides, III, 118
 phyllacanthus, III, 118
 phyllacanthus laevior, III,
 118
 phyllacanthus laevis, III, 118
 phyllacanthus macracanthus,
 III, 118

Echinocactus—*continued*,
 phyllacanthus micracanthus,
 III, 118
 phyllacanthus pentacanthus,
 III, 118
 phyllacanthus tenuiflorus, III,
 118
 phyllacanthus tricuspidatus,
 III, 117, 118
 phymatothele, IV, 8
 phymatothelos, IV, 8, 13
 piliferus, III, 125
 pilosus, III, 124, 125, 126,
 147
 pilosus canescens, IV, 287
 pilosus pringlei, III, 125
 pilosus steinesi, III, 124
 placentiformis, III, 219
 plaschnickii, IV, 30
 platensis, III, 163, 164
 platensis leptanthus, III, 163,
 164
 platensis parvulus, III, 163
 platensis quehlianus, III, 163
 platensis typicus, III, 163
 platyacanthus, III, 166, 167,
 171
 platycarpus, III, 180
 platycephalus, III, 181
 platyceras, III, 169
 platyceras laevior, III, 169
 platyceras minax, III, 169
 plicatilis, III, 180
 pluricostatus, III, 180
 poliacanthus, III, 198
 polyacanthus, III, 167
 polyancistrus, III, 167, 212,
 213, 214
 polycephalus, III, 167, 174
 polycephalus flavispinus, III,
 174
 polycephalus xeranthemoides,
 III, 173
 polygrhaphis, III, 103
 polyocentrus, III, 138
 polyraphis, III, 103
 polyrhaphis, III, 103
 porrectus, IV, 8
 posegerianus, IV, 13, 28
 pottsianus, IV, 136, 137
 pottsii, IV, 12, 13
 praegnacanthus, IV, 288
 pringlei, III, 125, 126
 pruinosis, III, 88, 89
 pseudo-cereus, III, 98
 pseudominusculus, III, 47
 pubispinus, III, 213
 pulchellus, III, 33
 pulcherrimus, III, 194
 pulverulentus, III, 178
 pumilus, III, 209, 210
 pumilus gracillimus, III, 209
 punctulatus, III, 180
 purpureus, IV, 288
 pycnacanthus, IV, 41
 pycnoxyphus, III, 139
 pygmaeus, III, 210, 211
 pygmaeus phaeodiscus, III,
 210
 pyramidatus, III, 90
 pyramidalis, II, 139
 quadrinatus, III, 122
 quehlianus, III, 163, 164
 radians, IV, 36
 radiatus, IV, 42
 radicans, IV, 36
 radiosus, IV, 43
 rafaelensis, III, 47
 raphidacanthus, III, 122; IV,
 15
 raphidocentrus, III, 122
 rebutii, III, 180
 rectispinus, III, 134
 recurvus, III, 141, 142, 143
 recurvus bicolor, III, 143
 recurvus latispinus, III, 143
 recurvus solenacanthus, III,
 143

Echinocactus—*continued*,
 recurvus spiralis, III, 142
 recurvus tricuspidatus, III,
 143
 reductus, III, 158
 reichei, III, 191
 reichenbachii, III, 25
 rettigii, III, 101
 retusus, III, 180
 rhodacanthus, III, 78, 79
 rhodacanthus coccineus, III,
 79
 rhodanthus, IV, 288
 rhodophthalmus, IV, 11, 12
 rhodophthalmus ellipticus,
 IV, 11, 12
 rinconadensis, IV, 7
 rinconensis, IV, 7
 robustus, III, 135, 136, 147
 robustus monstrosus, III,
 136
 robustus prolifer, III, 136
 rosaceus, III, 199
 rostratus, III, 59, 97, 98
 rotherianus, III, 206
 rubidus superbissimus, III,
 98
 rubripinus, III, 127
 saglionis, III, 197; IV, 288
 salmensis, IV, 28
 salm-dyckianus, III, 88; IV,
 39
 salmianus, III, 74, 222
 salmii, III, 180
 salpingophorus, III, 72
 saltensis, III, 5
 saltillensis, III, 172, 173;
 IV, 28
 sandillon, III, 186, 187
 sanjuanensis, III, 206
 santa-maria, III, 131
 saussierii, IV, 13
 scheeri, IV, 3, 4, 5, 28
 scheeri breviamatus, IV, 5
 scheerii, IV, 4
 schickendantzii, III, 87, 152,
 164, 165
 schillinkyanus, III, 210
 schillinkyanus grandiflorus,
 III, 210
 schlechtendalii, IV, 30
 schluengeri, III, 158
 schottii, IV, 11
 schumannianus, III, 189
 schumannianus longispinus,
 III, 189
 schumannianus nigrispinus,
 III, 191
 sclerothrix, III, 128
 scopa, III, 122, 193, 194,
 202; IV, 289
 scopa albicans, III, 193
 scopa candidus, III, 193, 194
 scopa candidus cristatus, III,
 193, 194
 scopa cristatus, III, 193, 194;
 IV, 289
 scopa ruberrimus, III, 193
 scopa rubra, IV, 289
 scopa rubrinus, III, 193
 sellowianus, III, 188, 189
 sellowianus tetracanthus, III,
 188
 sellowii, III, 188, 189; IV,
 288
 sellowii acutatus, III, 188
 sellowii courantii, III, 188
 sellowii macrocanthus, III,
 188, 189
 sellowii macrogonus, III,
 188, 189
 sellowii martinii, III, 188,
 189
 sellowii tetracanthus, IV,
 288
 sellowii turbinatus, III, 188,
 189
 sellowii typicus, III, 188

Echinocactus—*continued*,
 senilis, II, 27; III, 94
 sessiliflorus, III, 188, 189
 sessiliflorus pallidus, III, 188
 sessiliflorus tetracanthus, III,
 188
 setispinus, III, 104, 105, 106;
 IV, 287
 setispinus cachetianus, III,
 104
 setispinus hamatus, III, 104,
 105
 setispinus longihamatus, III,
 144
 setispinus longispinus, III,
 106
 setispinus martelii, III, 105
 setispinus micrensis, III, 105
 setispinus muehlenpfordtii,
 III, 105
 setispinus orcuttii, III, 105
 setispinus robustus, III, 144
 setispinus setaceus, III, 104,
 105
 setispinus sinuatus, III, 144
 setosus, III, 77
 sickmannii, III, 180
 sileri, III, 215
 similis, IV, 52
 simpsonii, III, 90, 91, 92
 IV, 286
 simpsonii minor, III, 90
 simpsonii robustior, III, 90,
 91
 sinuatus, III, 144, 146
 smithii, IV, 13
 soehrensii, III, 202, 203
 soehrensii albispinus, III,
 203
 soehrensii brevispinus, III,
 203
 soehrensii niger, III, 203
 solenacanthus, III, 142
 sparathacanthus, III, 180
 spectabilis, III, 136
 spagazzinii, III, 158, 190
 sphacelatus, IV, 138
 sphaerocephalus, III, 114
 spina-christi, III, 178, 238
 spiniflorus, III, 8, 78, 178,
 179
 spinosior, III, 213
 spinosissimus, III, 179; IV,
 118
 spinosus, III, 111
 spiralis, III, 141, 142, 143
 spiralis stellaris, III, 142
 stainesii, III, 124, 125, 180
 staplesiae, II, 27
 steinmannii, III, 47, 45
 stellaris, III, 142
 stellatus, III, 142, 152, 163
 stenocarpus, III, 163
 stenogoni, III, 118
 stenogonus, III, 117
 strausianus, III, 201
 streptocaulon, III, 86, 87
 strobiliformis, IV, 54
 stuckertii, III, 165
 suberinaceus, III, 158
 subgibbosus, III, 59, 94, 96,
 97, 98
 subglaucus, III, 213
 subgrandicornis, III, 181
 submammulosus, III, 200,
 201; IV, 289
 subniger, III, 100
 subporrectus, IV, 8
 subuliferus, III, 136
 sulcatus, III, 64
 sulcolanatus, IV, 37
 sulphureus, III, 123
 supertextus, III, 100
 tabularis, III, 193
 tabularis cristatus, III, 193
 tellii, III, 123
 tenuiflorus, III, 118
 tenuispinus, III, 166, 195, 196

- Echinocactus*—continued,
tenuispinus minor, III, 195
tenuispinus ottonis, III, 196
tenuissimus, III, 196
tephracanthus, III, 188
tephracanthus spinosior, III, 188
teretispinus, III, 123
terscheckii, III, 199
tetracanthus, III, 188, 189
tetracentrus, IV, 287
tetraxiphus, III, 112
texensis, III, 167, 181, 182;
IV, 288
texensis gourginsii, III, 181
texensis longispinus, III, 181
texensis treculianus, III, 145
theiacaanthus, III, 122, 137
theionacanthus, III, 122, 137
thelephorus, III, 181
theloides, IV, 8
thionanthus, III, 49, 57
thrinogonus, III, 97, 98
thrinogonus elatior, III, 97
thortusos, III, 195, 196
tortus, III, 185
towensis, III, 158
treculianus, III, 144
tribolacanthus, III, 123
tricolor, IV, 12
tricornis, III, 218
tricuspidatus, III, 117
trifurcatus, III, 123
trollietii, III, 150, 151
tuberculatus, III, 170
tuberculatus spiralis, III, 170
tuberisulcatus, III, 202, 203
tuberosus, IV, 9
tuberosus subporrectus, IV, 9
tubiflorus, III, 67
tulensis, IV, 8, 11
turbatus, III, 66
turbiniiformis, III, 106, 107;
IV, 287
uncinatus, III, 146
uncinatus wrightii, III, 146
undulatus, III, 117
unguispinus, III, 150, 151;
IV, 287
uruguayensis, III, 162
valparaiso, III, 98
vanderayeii, III, 136
vargasii, IV, 13
verutum, III, 118
victoriensis, III, 136
villiferus, III, 181
villosus, III, 103
villosus crenator, III, 103
violaciflorus, III, 114, 115
viridescens, III, 135, 140,
141; IV, 287
viridescens cylindraceus, III,
129
viridiflorus, III, 17
visnaga, III, 167, 170, 171;
IV, 288
viviparus, IV, 43
wangertii, II, 133
weberbaueri, III, 103
wegereri, III, 122
weinartianus, III, 206
whipplei, III, 167, 213
whipplei nanus, III, 213
whipplei spinosior, III, 213
wilhelmii, III, 181
williamsii, III, 83, 84, 85;
IV, 286
williamsii lewinii, III, 85
winkleri, IV, 41
wippmannii, III, 111
wislizeni, III, 123, 127, 128
wislizeni albispinus, III, 127
wislizeni albus, III, 129
wislizeni decipiens, III, 127
wislizeni latispinus, III, 128
wislizeni lecontei, III, 129
wislizeni phoeniceus, III, 129
- Echinocactus*—continued,
wislizeni purpureus, III, 128
wrightii, III, 146
xanthacanthus, III, 230, 231
xeranthemoides, III, 167, 173
xiphacanthus, III, 114
Echinocactuses, IV, 168
Echinocarpace, I, 44, 56
Echinocereaceae, II, 1; III, 3-
77; IV, 285
Echinocereus, I, 79, 215; II,
3, 104, 110; III, 3-44, 45,
48, 60, 104, 146, 150, 202,
211, 212; IV, 12, 48, 57,
162; IV, 285
acifer, III, 4, 12, 13, 14;
IV, 285
acifer brevispinulus, III, 12
acifer diversispinus, III, 12
acifer durangensis, III, 12
acifer tenuispinus, III, 12
acifer trichacanthus, III, 12,
13; IV, 285
adustus, III, 5, 23, 24
aggregattis, III, 14; IV, 47
amoenus, III, 5, 33
baileyi, III, 3, 5, 26, 27
barcelona, IV, 287
barcelona, III, 43
barthelowanus, III, 5, 41
berlandieri, III, 20, 21
bertinii, III, 45
bicolor, III, 43
blanckii, III, 5, 20, 21; IV,
285
blankii, III, 20
bolansii, III, 40
boliviensis, III, 43
brandegei, III, 5, 34
caespitosus, III, 25, 26; IV,
285
caespitosus castaneus, III, 25
caespitosus major, III, 25
candicans, II, 142; III, 28
candicans tenuispinus, II,
143
carnosus, III, 36
centralis, III, 149
chiloensis, II, 138
chloranthus, 4, 16
chlorophthalmus, III, 5, 32
cinerascens, III, 5, 23; IV,
285
cinerascens crassior, III, 23
cinnabarinus, III, 54
cirrhiferus, III, 23
cirrhiferus monstrosus, III,
23
clavatus, II, 106
claviformis, III, 43
coccineus, III, 4, 13, 14; IV,
285
conglomeratus, III, 5, 32,
39, 40
conoideus, III, 4, 11, 13
ctenoides, III, 5, 19, 20
dahliaeflorus, III, 44
dasyacanthus, III, 5, 18, 19,
31
degandii, III, 9
delatetii, III, 4, 6
deppei, III, 23; IV, 285
dubius, III, 5, 39
durangensis, III, 12; IV, 285
durangensis nigripinus, III,
12
durangensis rufispinus, III,
12
ehrenbergii, III, 5, 41
ehrenbergii cristatus, III, 42
emoryi, II, 108; III, 127
engelmannii, III, 5, 38
engelmannii albispinus, III,
38
engelmannii chrysocentrus,
III, 38
engelmannii fulvispinus, III,
38
- Echinocereus*—continued,
engelmannii pferdorffii, III,
38
engelmannii robustior, III, 38
engelmannii variegatus, III,
38
engelmannii versicolor, III,
38
enneacanthus, III, 5, 36, 39;
IV, 285
enneacanthus carnosus, III,
36
fendleri, III, 5, 9, 14, 35, 36,
37, 40
fitchii, III, 5, 30
flavescens, II, 20; III, 212
flaviflorus, III, 15, 21
galtieri, III, 43
gladiatus, II, 142
glycimorphus, III, 23
gonacanthus, III, 10
grahamii, III, 43
grandis, III, 4, 18
havermansii, III, 43
hempeli, III, 5, 34, 35
hexaedrus, III, 10
hildmannii, III, 36
huitcholensis, III, 4, 8
hypogaeus, II, 106
inermis, III, 14, 32
intricatus, II, 143
jacobyi, III, 35
knippelianus, III, 5, 32
krausei, III, 14
kunzei, III, 31
labouretianus, III, 17
labouretii, III, 17
lamprochlorus, II, 132
lecanus, III, 4, 9, 43
lecanus multicosatus, III, 9
leonensis, III, 20, 21
leptacanthus, III, 22
liebnerianus, III, 32
limensis, II, 20
lloydii, III, 5, 37
longisetus, III, 5, 42
luteus, III, 4, 16, 17
malibranii, III, 43
mamillatus, III, 5, 41, 42
mamillosus, III, 43
maritimus, III, 4, 15
merkeri, III, 5, 35, 36
mohavensis, III, 8
mojavensis, III, 4, 8
mojavensis zuniensis, III, 14
monacanthus, III, 10
multangularis, II, 19; IV,
279
multangularis limensis, II,
20
multangularis pallidior, II,
19
multicostatus, III, 9
neo-mexicanus, III, 4, 11, 13
octacanthus, III, 4, 13
orcuttii, III, 15
pacificus, III, 4, 11, 12
palmeri, III, 5, 34
papillosus, III, 5, 19
papillosus rubescens, III, 19
paucispinus, III, 10
paucispinus flavispinus, III,
10
paucispinus gonacanthus, III,
10
paucispinus hexaedrus, III,
10
paucispinus triglochidiatus,
III, 10
paucipina, III, 44
pectinatus, III, 29, 5, 30
pectinatus adustus, III, 24
pectinatus armatus, III, 24
pectinatus caespitosus, III,
25, 26
pectinatus candicans, III, 28
pectinatus castaneus, III, 26
pectinatus centralis, III, 149
- Echinocereus*—continued,
pectinatus chrysacanthus, III,
29
pectinatus cristatus, III, 30
pectinatus laevior, III, 30
pectinatus rigidissimus, III,
27, 28
pectinatus robustior, III, 28
pectinatus robustus, III, 27,
28
pectinatus rufispinus, III, 24
penicilliformis, III, 4
penisilis, III, 4, 8
pentaplophus, III, 5, 20, 21;
IV, 285
pentlandii, III, 54
perbellus, III, 5, 24, 25
persolatus, III, 44
phoeniceus, III, 14
phoeniceus albispinus, III,
14
phoeniceus conoideus, III, 13
phoeniceus inermis, III, 14
phoeniceus longispinus, III,
14
phoeniceus rufispinus, III,
14
pleiogonus, III, 9, 43
polyacanthus, III, 4, 9, 11,
12, 15; IV, 285
polycephalus, III, 44
poselgeri, III, 110, 111
poselgerianus, III, 20
princeps, III, 44
procumbens, III, 22; IV, 285
procumbens longispinus, III,
22
propinquus, III, 22
pulchellus, III, 5, 33
pulchellus amoenus, III, 33
radians, III, 23
raphicephalus, III, 44
reichenbachianus, III, 25
reichenbachii, III, 5, 25, 27,
28, 31; IV, 52, 285
rigidispinus, III, 28
rigidissimus, III, 5, 27, 28, 29
robustior, III, 28
roemerii, III, 13, 14
roetteri, III

- Echinocereus*—*continued*,
 thurberi, III, 43
 thwaitesii, III, 44
 trichacanthus, II, 44
 triglochidiatus, III, 4, 10
 trockyi, III, 44
 tuberosus, II, 111
 uehri, III, 44
 undulatus, III, 23
 uspenskii, III, 44
 viridiflorus, III, 3, 4, 16, 17,
 24, 31; IV, 285
 viridiflorus cylindricus, III,
 17
 viridiflorus gracilispinus, III,
 17
 viridiflorus major, III, 17
 viridiflorus tubulosus, III, 17
 turbergii, III, 5, 29
Echinofossulocactus, III, 78,
 109-123, 138; IV, 287
 albatrus, III, 110, 112
 anfractusus, III, 110, 117
 arrigens, III, 110, 113, 114
 confusus, III, 110, 120
 coptonogonus, III, 110, 121
 coptonogonus major, III,
 110
 cornigerus, III, 143
 cornigerus angustispinus, III,
 143
 cornigerus elatior, III, 143
 cornigerus rubrospinus, III,
 143
 crispatus, III, 110, 116
 dichroacanthus, III, 110,
 117
 echidne, III, 136
 ensiformis, III, 114
 gladiatus, III, 110, 119, 120
 grandicornis, III, 110, 114
 harrisii, III, 109
 hastatus, III, 110, 111
 helophora, III, 170
 helophora longifossulatus,
 III, 170
 heteracanthus, III, 110, 112
 ignotus-venosus, III, 109
 karvinskianus, III, 170
 lamellosus, III, 110, 113
 lancifer, III, 110, 118, 119
 lloydii, III, 110, 112, 113
 macracanthus, III, 170
 mirbelii, III, 185
 multicostatus, III, 110, 111,
 113
 obvallatus, III, 110, 115, 116
 oxypterus, III, 138
 pentacanthus, III, 110, 115
 pfeifferi, III, 137
 phyllacanthus, III, 110, 118,
 119
 phyllacanthus macracanthus,
 III, 118
 phyllacanthus micracanthus,
 III, 118
 platyceras, III, 169
 recurvus campylacanthus, III,
 142
 robustus, III, 135
 tricuspidatus, III, 110, 117
 turbiniiformis, III, 106
 vanderacyi, III, 136
 vanderacyi ignotus-longispi-
 nus, III, 136
 violaciflorus, III, 110, 114,
 115
 wippermannii, III, 110, 111
 zacatecasensis, III, 110, 113
Echinomastus, III, 78, 147-152
 dasycanthus, III, 148, 150,
 151; IV, 55
 durangensis, III, 148, 152
 erectocentrus, III, 148; IV,
 16, 175, 287
 intertextus, III, 148, 149,
 150; IV, 287
 madowellii, III, 148, 151
Echinomastus—*continued*,
 unguispinus, III, 148, 150,
 152; IV, 287
Echinomelocactus, III, 221,
 225, 231
 minor, IV, 71
Echinonyctanthus, III, 60
 decaisneanus, III, 66
 eyriesii, III, 65
 leucanthus, III, 72
 multiplex, III, 64
 nigrispinus, III, 67
 oxygonus, III, 64
 pictus, III, 66
 pulchellus, III, 33
 schelhasii, III, 66
 tubiflorus, III, 67
 tubiflorus nigrispinus, III, 67
 turbinatus, III, 66
 turbinatus pictus, III, 66
Echinopsis, II, 3, 105, 130,
 144, 158, 159, 219; III, 3,
 4, 45, 48, 49, 53, 60-77,
 78, 79, 100, 101
 achatina, III, 55
 albispina, III, 76
 albispinosa, III, 61, 67
 amoena, III, 33, 34
 ancistrophora, III, 61, 69
 apiculata, III, 63
 aurata, II, 143; III, 79,
 186
 aurea, III, 60, 61, 74
 baldiana, III, 61, 74
 beckmannii, III, 76
 boeckmannii, III, 76
 boutillieri, III, 76
 bridgesii, III, 61, 63, 74;
 IV, 286
 cachensis, III, 52
 caespitosa, III, 53
 calochlora, III, 61, 68
 campylacantha, III, 69, 72
 campylacantha brevispina,
 III, 72
 campylacantha leucantha, III,
 72
 campylacantha longispina,
 III, 72
 campylacantha stylodes, III,
 72
 campylacantha stylosa, III,
 72
 candicans, II, 142
 catamarcensis, II, 146; IV,
 278
 cavendishii, III, 54
 chereauiana, III, 54
 cinnabarina, III, 54, 55
 cinnabarina cheroniana, III,
 54
 cinnabarina cristata, III, 54
 cinnabarina scheeriana, III,
 54
 cinnabarina spinosior, III,
 54
 colmariensis, III, 55
 colmarii, III, 54, 55
 columnaris, III, 55
 cordobensis, III, 61, 69
 cristata, III, 73
 cristata purpurea, III, 73
 decaisneana, III, 66, 67; IV,
 286
 deminuta, III, 48; IV, 285
 droegeana, III, 67
 ducis paulii, III, 76
 dumeliana, II, 143
 dumesniliana, II, 143
 duvallii, III, 76
 elegans vittata, III, 55
 eyriesii, III, 61, 65, 76; IV,
 286
 eyriesii cristata, III, 65
 eyriesii duvallii, III, 65
 eyriesii flore-pleno, III, 66
 eyriesii glauca, III, 65
 eyriesii glaucescens, III, 65
Echinopsis—*continued*,
 eyriesii grandiflora, III, 6
 eyriesii inermis, III, 66
 eyriesii lagemannii, III, 64
 eyriesii major, III, 65
 eyriesii phylligera, III, 65
 eyriesii rosea, III, 65
 eyriesii tettavii, III, 65, 66
 eyriesii triumphans, III, 65,
 66
 eyriesii wilkensis, III, 64
 falcata, III, 66
 fiebrigii, III, 61, 70, 71
 fischeri tephroacantha, III, 77
 fobeana, III, 76
 forbesii, III, 60, 62, 63
 formosa, III, 60, 61, 75; IV,
 286
 formosa albispina, III, 75
 formosa gilliesii, III, 75
 formosa laevior, III, 75
 formosa rubripina, III, 75
 formosa spinosior, III, 75
 formosissima, III, 76; IV, 286
 gemmata, III, 66, 67; IV, 286
 gemmata cristata, III, 67
 gemmata decaisneana, III, 66
 gemmata schelhasii, III, 66
 gibbosa, III, 158
 gigantea, III, 76
 grandiflora, III, 67
 haageana, III, 185
 hempeliana, III, 101
 huottii, III, 60, 63
 intricatissima, 61, 73
 jamesiana, III, 66
 kuottii, III, 63
 lagemannii, III, 64, 65
 lamprochlora, II, 132
 laterita, III, 56
 leucantha, III, 61, 72, 98,
 140; IV, 286
 leucantha aurea, III, 72, 73
 leucantha salpingophora, III,
 72
 longispina, III, 76
 mamillosa, III, 61, 75
 maximiliana, III, 54
 maximiliana longispina, III,
 55
 melanacantha, III, 67
 melanopotamica, III, 72
 meyeri, III, 60, 61, 62
 mieckleyi, III, 75
 minuana, III, 60, 63
 minuscula, III, 45
 mirabilis, III, 60, 62
 misleyi, III, 73
 molesta, III, 61, 74
 muelleri, III, 76
 multiplex, III, 60, 61, 64;
 IV, 286
 multiplex cossa, III, 64
 multiplex cristata, III, 64
 multiplex monstrosa, III, 64
 multiplex picta, III, 64
 nigerrima, III, 65, 77
 nigricans, III, 76, 95
 nigrispina, III, 67
 nodosa, III, 105
 obliqua, III, 73
 obrepanda, III, 61, 73
 ochroleuca, III, 55
 octacantha, III, 13
 oxygona, III, 60, 61, 64, 65,
 76; IV, 286
 oxygona inermis, III, 65; IV,
 286
 oxygona subinermis, III, 65
 oxygona turbinata, III, 65
 paraguayensis, III, 76
 pectinata, III, 29, 30
 pectinata laevior, III, 30
 pectinata reichenbachiana,
 III, 25
 pentlandii, III, 51, 54, 55;
 IV, 285
 pentlandii achatina, III, 55
Echinopsis—*continued*,
 pentlandii albiflora, III, 55
 pentlandii cavendishii, III,
 54
 pentlandii coccinea, III, 54
 pentlandii colmari, III, 55
 pentlandii cristata, III, 55
 pentlandii elegans, III, 54
 pentlandii forbesii, III, 54
 pentlandii gracilispina, III,
 54
 pentlandii integra, III, 55
 pentlandii laevior, III, 54
 pentlandii levior scheeri, III,
 55
 pentlandii longispina, III,
 54, 55
 pentlandii maximiliana, III,
 54, 55
 pentlandii neuberti, III, 54
 pentlandii ochroleuca, III, 54
 pentlandii pferdorffii, III, 54
 pentlandii pyrantha, III,
 54
 pentlandii pyrantha, III, 55
 pentlandii radians, III, 54
 pentlandii scheeri, III, 54, 55
 pentlandii tricolor, III, 54
 pentlandii vitellina, III, 54
 pferdorffii, III, 55
 philippi, II, 105
 picta, III, 66
 polyacantha, III, 72
 polyphylla, III, 76
 poselgeri brevispina, III, 72
 poselgeri longispina, III, 72
 pseudominuscula, III, 47
 pudatiti, III, 6, 66
 pulchella, III, 33
 pulchella amoena, III, 33
 pulchella rosea, III, 34
 pygmaea, III, 47
 pyrantha, III, 76
 quehlii, III, 76
 reichenbachiana, III, 26
 rhodacantha, III, 79, 101
 rhodacantha aurea, III, 79
 rhodacantha gracilior, III, 79
 rhodotricha, III, 61, 71
 rhodotricha argentinensis,
 III, 71
 rhodotricha robusta, III, 71
 rhodotricha roseiflora, III, 71
 roehlandii, III, 65
 roehlandii, III, 64
 salm-dyckiana, III, 76
 salmiana, III, 74; IV, 286
 salmiana bridgesii, III, 74
 salpingophora, III, 72
 salpingophora, III, 69, 72
 salpingophora aurea, III, 72,
 73
 saltensis, III, 53
 saluciana, III, 76
 scheeri, III, 54, 55
 scheeriana, III, 55
 schelhasii rosea, III, 66
 schelhasii, III, 66
 schickendantzii, II, 144; IV,
 278
 scopa, III, 193, 194
 scopa candida cristata, III,
 194
 shaferi, III, 61, 69, 70
 silvestrii, III, 61, 68
 simplex, III, 72
 spegazziniana, III, 61, 69, 70
 spegazzinii, III, 71
 stylosa, III, 72
 sulcata, III, 64
 tacuarembense, III, 76
 tettavii, III, 66
 tougardii, III, 76
 tricolor, III, 54
 triumphans, III, 65; IV, 286
 tuberculata, III, 76
 tubiflora, III, 61, 6, 67; IV,
 286

- Echinopsis*—*continued*,
tubiflora nigripina, III, 67
tubiflora paraguayensis, III, 67
tubiflora rohlandii, III, 67;
 IV, 286
tubiflora rosea, III, 67
turbinata, III, 61, 66; IV, 286
turbinata picta, III, 66
undulata, III, 6, 76
valida, III, 62, 63
valida densa, III, 12, 13, 14
valida forbesii, III, 62
verschaffeltii, III, 63
wilkensii, III, 64
yacutulana, III, 72
zuccariniana, III, 67; IV, 286
zuccariniana cristata, III, 67
zuccariniana monstrosa, III, 67
zuccariniana nigripina, III, 67
zuccariniana picta, III, 67
zuccariniana rohlandii, III, 67
zuccariniana rosea, III, 67
zuccarinii, III, 67, 196
zuccarinii monstrosa, III, 67
zuccarinii nigripina, III, 67
zuccarinii picta, III, 67
zuccarinii robusta, III, 67
Elatae, I, 45, 152, 156
Elatiores, I, 45, 149, 152, 155, 156, 222
Epiphyllanae, II, 1; IV, 177-207
Epiphyllanthus, IV, 177, 180-182, 209, 220
candidus, IV, 180, 182
microsphaericus, IV, 180, 181
obovatus, IV, 180
obtusangulus, IV, 180, 181
Epiphyllum, I, 9; II, 127, 221; III, 7; IV, 177, 179, 180, 183, 185-201, 204, 220, 250
ackermannii, II, 128, 219; IV, 198, 199, 200
acuminatum, IV, 188, 189
aitoni, IV, 198
alatum, IV, 213, 243
album violaceum, IV, 200
altensteinii, IV, 177, 179
amabile, IV, 200
americanum, IV, 187
anguliger, IV, 187, 191
aurantiacum, IV, 200
biforme, IV, 202
brasiliense, IV, 200
bridgesii, IV, 185
candidum, IV, 182
carminatum, IV, 200
cartagense, IV, 187, 197
caudatum, IV, 187, 190
caulorhizum, IV, 192
chiapensis, IV, 203
ciliare, IV, 241
ciliatum, IV, 241
costaricense, IV, 193
crenatum, IV, 187, 192, 193, 199
crenulatum, IV, 244
crispatum, IV, 245
cruentum, IV, 200
darrahii, IV, 187, 190, 191
delicatum, IV, 177, 179
delicatum, IV, 177, 179
elegans, IV, 179
gaertneri, IV, 183, 184
gaertneri coccineum, IV, 184
gaertneri mackoyanum, IV, 184
gaillardae, I, 6; IV, 187, 188
gibsonii, IV, 178
gracilis, IV, 200
Epiphyllum—*continued*,
grande, IV, 188
grande superbum, IV, 200
grandiflorum, IV, 200
grandilobum, IV, 187, 192
guatemalense, IV, 187, 195
guedeneyi, IV, 178
harrissonii, IV, 200
hercule, IV, 200
hitchinii, IV, 198
hookeri, IV, 187, 197
hybridum, IV, 198
jenkensoni, IV, 198
latetium album, IV, 200
latifrons, IV, 189, 196
lepidocarpum, IV, 187, 194
macropterum, IV, 187, 192, 193, 194
mackoyanum, IV, 183, 184
maximum, IV, 200
multiflorum, IV, 200
nelsonii, IV, 203
obovatum, IV, 180, 181
obtusangulum, IV, 181
opuntoides, IV, 180, 181
oxypetalum, IV, 187, 188, 193
palidum roseum, IV, 200
phyllanthoides, II, 128; IV, 188, 198, 205
phyllanthus, IV, 186, 187, 188, 197
pittieri, IV, 187, 190, 194
platycarpum, IV, 242
pumilum, IV, 187, 189
purpurascens, IV, 177
purpureum, IV, 179
ramulosum, IV, 241
rhombium, IV, 244
roseum, IV, 200
rubrum violaceum, IV, 200
ruckeri, IV, 179
ruckerianum superbum, IV, 200
rueckerianum, IV, 185
russellianum, IV, 182, 184, 185
russellianum gartneri, IV, 184
russellianum gaertneri, IV, 183
russellianum rubrum, IV, 185
russellianum superbum, IV, 185
salmonium, IV, 178
salmonium marginatum, IV, 200
smithianum, II, 128
speciosum, IV, 200, 205
speciosum jenkensoni, IV, 198
speciosum lateritium, IV, 200
spectabile, IV, 178
spectabile coccineum, IV, 200
splendens, IV, 199
splendidum, IV, 198
stenopetalum, IV, 187, 196
strictum, IV, 187, 196, 197
thomasianum, IV, 193
translucens, IV, 200
tricolor, IV, 200
truncatum, IV, 21, 177, 178, 179, 180, 185, 186
truncatum albiflorum, IV, 178
truncatum altensteinii, IV, 177
truncatum amabile roseum, IV, 179
truncatum aurantiacum, IV, 178
truncatum bicolor, IV, 178
truncatum bridgesii, IV, 185
truncatum carmineum, IV, 179
truncatum coccineum, IV, 178
Epiphyllum—*continued*,
truncatum cruentum, IV, 178, 179
truncatum elegans, IV, 178, 180
truncatum gracile, IV, 179
truncatum grandidens, IV, 178
truncatum grandiflorum rubrum, IV, 179
truncatum harrissonii, IV, 179
truncatum lateritium album, IV, 179
truncatum magnificum, IV, 178
truncatum makoyanum, IV, 179
truncatum maximum, IV, 179
truncatum minus, IV, 178
truncatum morellianum, IV, 179
truncatum multiflorum, IV, 179
truncatum pallidum roseum, IV, 179
truncatum purpureum, IV, 178
truncatum purpurascens, IV, 179
truncatum purpureum, IV, 179
truncatum roseum, IV, 178
truncatum rubrum violaceum, IV, 179
truncatum ruckerianum, IV, 178, 185
truncatum russellianum, IV, 184, 185
truncatum salmonium, IV, 178
truncatum salmonium aurantiacum, IV, 179
truncatum salmonium brasiliense, IV, 179
truncatum salmonium flavum, IV, 179
truncatum salmonium marginatum, IV, 179
truncatum salmonium rubrum, IV, 179
truncatum snowi, IV, 179
truncatum spectabile, IV, 177, 178, 179
truncatum spectabile superbum, IV, 179
truncatum splendens, IV, 179
truncatum translucens, IV, 179
truncatum tricolor, IV, 178
truncatum vanhoutteanum, IV, 178
truncatum violaceum, IV, 177, 197
truncatum violaceum album, IV, 179
truncatum violaceum grandiflorum, IV, 179
truncatum violaceum superbum, IV, 178
vandesii, IV, 205
violaceum, IV, 179
violaceum elegans, IV, 200
violaceum grandiflorum, IV, 200
violaceum rubrum, IV, 200
violaceum speciosum, IV, 200
violaceum superbum, IV, 200
Epithelantha, III, 77, 92, 93, 208; IV, 65
micromeris, III, 93
Erdisia, II, 2, 104-107; III, 4
meyenii, II, 104, 105, 106
philippii, II, 104, 105
Erdisia—*continued*,
spiniflora, II, 104, 106
squarrosa, II, 104, 105, 107
Eriocerei, II, 154
Eriocereus, TI, 22, 147, 148
bonplandii, II, 157
cavendishii, II, 21
jusbertyi, II, 158
martianus, II, 220
martinii, II, 155
platygonus, II, 156
subrepandus, II, 151
tephracanthus, II, 136
tortuosus, II, 154
Eriosyce, II, 143; III, 78, 167, 186, 187
ceratistes, III, 79, 186
sandillon, II, 143; III, 186
Erythrina, I, 181
Erythrorhopsalis, IV, 208, 209
pilocarpa, IV, 209
Escobaria, IV, 3, 53-57
bella, IV, 54, 56
chaffeyi, IV, 54, 56
chihuahuensis, IV, 54, 55
dasyacantha, IV, 54, 55
lloydii, IV, 54, 57
runyonii, IV, 54, 55, 56
sneedii, IV, 54, 56
tuberculosa, IV, 54, 55, 57
Escontria, II, 2, 63, 65, 66
chiotilla, II, 65, 66, 76; IV, 270, 272
Espina, I, 76
Espina blanca, I, 41
Espinha de São Antonio, I, 19
Espostoa, II, 2, 60-63
lanata, II, 61, 62, 225; IV, 270
Etuberculatae, I, 72, 73
Eucereus, II, 3, 117
Euharrisia, II, 148
Eulychnia, II, 2, 66, 82-85, 181; III, 4
acida, II, 82, 83, 84
breviflora, II, 82, 83
castanea, II, 82, 84
clavata, II, 106
eburnea, II, 139
iquiquensis, II, 82, 83
spinibarbis, II, 82, 83; IV, 272
Eupereskia, I, 10
Euphorbia hystrix, II, 19
Facheiropreto, II, 177
preto da Serra de Cannabrava, II, 173
Facheiroa, II, 2, 173
pubiflora, II, 173
Feather ball, IV, 124
Ferocactus, III, 78, 123-147, 148, 177, 212
acanthodes, III, 124, 129, 130, 131, 147; IV, 287
alamosanus, III, 124, 137
chrysacanthus, III, 123, 127
covillei, III, 124, 132, 133, 134
crassihamatus, III, 124, 144
diguetii, III, 124, 131; IV, 287
echidne, III, 124, 136
flavovirens, III, 124, 138
fordii, III, 123, 126
glaucescens, III, 124, 137
hamatacanthus, III, 124, 144; IV, 287
horridus, III, 123, 128
johnsonii, III, 124, 141, 142
johnstonianus, IV, 287
latispinus, III, 124, 143; IV, 287
lecontei, III, 124, 129; IV, 287
macrodiscus, III, 124, 139, 140
melocactiformis, III, 124, 138, 139

Ferocactus—*continued*,
 nobilis, III, 123, 124, 141,
 142
 orcuttii, III, 124, 134, 141
 peninsulae, III, 124, 133, 134
 pringlei, III, 123, 125
 rectispinus, III, 124, 134, 135
 robustus, III, 124, 134, 135,
 136, 138
 rostii, III, 124, 146, 147
 santa-maria, III, 124, 131
 stainesii, III, 123, 124; IV,
 287
 townsendianus, III, 123, 126,
 127
 uncinatus, III, 124, 145, 146
 viridescens, III, 124, 135,
 140, 141; IV, 287
 wislizeni, III, 123, 125, 127,
 128, 129, 130, 132, 134
Ficindica, I, 42, 43
Ficoides, IV, 71, 125
Ficus-indicae, I, 45, 156, 166,
 177, 191
Floccosae, I, 44, 86; IV, 221, 234
Flor de baile, IV, 189
Flor de cera, I, 19
Flor de copa, III, 114
Flor de cuerno, II, 2, 18
Flor de la oración, III, 62
Formosi, II, 3, 13
Fox-tail cactus, IV, 7
Frailca, III, 78, 208-211
caespitosa, III, 209, 211
cataphracta, III, 209, 210,
 211
gracillima, III, 209
grahiana, III, 209
knippeliana, III, 209, 211
pumila, III, 209
pygmaea, III, 209, 210
schilinzkyana, III, 209, 210
Frutescentes, I, 73
Fulgidae, I, 44, 67
Fulvispinosae, I, 148
Furcraea, II, 92
Garranibullas, II, 179
Geotilla, II, 66
Giant cactus, II, 92, 164; IV,
 279
Giganton, II, 135
Gladiatores, III, 109
Glomeratae, I, 44, 87
Graciles, II, 159
Grandiflorae, IV, 221
Grandifoliae, I, 9, 11
Green-flowered petaya, III, 17
Grizzly bear cactus, I, 196
Gruonia, I, 24, 215
bradtiana, I, 215
crecifformis, I, 215
Guamacho, I, 17
Guasabara, IV, 266
Gymnocalycium, II, 1; III, 78,
 87, 152-166
anisitsii, III, 159, 160
brachyanthum, III, 153, 159;
 IV, 288
damsii, III, 153, 163
denudatum, III, 152, 155,
 156; IV, 288
gibbosum, III, 153, 157, 158,
 161; IV, 288
guerkeanum, III, 152, 154,
 155
hyptiacanthum, III, 152, 156
joossellianum, III, 153, 166
kurtzianum, III, 153, 162,
 163
lecanum, III, 152, 154, 156,
 157
megalothelos, III, 153, 162
melanocarpum, III, 153, 161
mihanovichii, III, 152, 153
monvillei, III, 153, 160, 161;
 IV, 288
mostii, III, 153, 158
multiflorum, III, 153, 159

Gymnocalycium—*continued*,
 netrellianum, III, 152, 153,
 154
 platense, III, 153, 163, 164,
 165
 reductum, III, 158, 159
 saglione, III, 152, 153, 157;
 IV, 288
 schickendantzii, III, 153,
 164, 165
 spegazzinii, III, 152, 155
 stuckertii, III, 153, 165
 uruguayense, III, 153, 161,
 162
 villosum, III, 103
Hamatocactus, III, 78, 104-
 106
setispinus, III, 104, 105; IV,
 287
Hariota, IV, 216, 219, 224, 240
alata, IV, 213
alternata, IV, 237
bambusoides, IV, 219
boliviana, IV, 240
cassutha, IV, 226
ceriformis, IV, 210
cereuscula, IV, 222
cinerea, IV, 212
clavata, IV, 224
conferta, IV, 227
coriacea, IV, 241
crenata, IV, 212
cribrata, IV, 225
crispata, IV, 245
crispata latior, IV, 245
cruciformis, IV, 215
cylindrica, IV, 231
fasciculata, IV, 229
floccosa, IV, 234
funalis, IV, 231
gracilis, IV, 217
grandiflora, IV, 231
horrida, IV, 229
houlettiana, IV, 238
knightii, IV, 211
knightii tenuispinis, IV, 215
lindbergiana, IV, 228
lumbricalis, IV, 230
macrocarpa, IV, 187
mesembrianthemoides, IV,
 222
micrantha, IV, 239
monacantha, IV, 212
pachyptera, IV, 243
paradoxa, IV, 237
parasitica, IV, 226
penduliflora, IV, 225
pentaptera, IV, 236
platycarpa, IV, 242
prismatica, IV, 222
ramosissima, IV, 216
ramulosa, IV, 240
rhombea, IV, 244
riedeliana, IV, 246
robusta, IV, 243
rugosa, IV, 234
saglione, IV, 222
salicornioides, IV, 217, 218
salicornioides bambusoides,
 IV, 218, 219
salicornioides gracilior, IV,
 217
salicornioides ramosior, IV,
 217
salicornioides schottmuelleri,
 IV, 218
salicornioides strictior, IV,
 217
sarmentacea, IV, 230, 231
spatulata, IV, 247
squamulosa, IV, 215
stricta, IV, 217
swartziana, IV, 213
teres, IV, 227
trigona, IV, 237
triquetra, IV, 243
tucumanensis, IV, 234
villigera, IV, 218

Harrisia, I, 24; II, 1, 2, 147-
 159; III, 60
aboriginum, II, 148, 154; IV,
 278
adscendens, II, 148, 155, 196
bonplandii, II, 148, 157; IV,
 278
brookii, II, 148, 151
earlei, II, 148, 154
eriophora, II, 148, 149; IV,
 278
fernowi, II, 148, 153
fragrans, II, 148, 149
gracilis, II, 148, 151, 152;
 IV, 278
guelichii, II, 148, 158
martinii, II, 148, 155; IV,
 278
nashii, II, 148, 150, 151;
 IV, 278
platygona, II, 148, 156
pomanensis, II, 148, 155, 156
portoricensis, II, 148, 150
simpsonii, II, 148, 152, 153
taylori, II, 153
tortuosa, II, 148, 154
undata, II, 151; IV, 278
Hatchet cactus, IV, 59
Hatiota, IV, 208, 216-219, 224
bambusoides, IV, 217, 218
cylindrica, IV, 217, 219
salicornioides, IV, 217
Hedgehog cactus, III, 177
Heliocereus, II, 2, 127-129,
 218, 225; IV, 198
amecamensis, II, 127, 129;
 IV, 277
cinnabarinus, II, 127, 129;
 IV, 277
coccineus, II, 127
elegantissimus, II, 127, 129;
 IV, 205
mallisoni, IV, 277
schrunkii, II, 127, 225
speciosus, II, 127, 128, 129,
 201, 210; III, 3; IV, 198,
 199, 276
Heliotropium, IV, 188
Hexagonae, II, 4
Hibiscus, I, 19
esculentus, I, 19
Hickenia, III, 78, 207, 208
microsperma, III, 207, 208;
 IV, 289
Homalocephala, III, 78, 181, 182
texasensis, III, 77, 181; IV, 288
Houlettianae, IV, 221
Hybocactus, III, 94, 152
Hydrochylus, IV, 88
Hylocereanae, II, 1, 183
Hylocereus, I, 24; II, 1, 63,
 126, 183-195, 210, 215,
 216; IV, 220, 282
antiguensis, II, 184, 193, 194
bronxensis, II, 183, 185, 226
calcaratus, II, 184, 193, 194
costaricensis, II, 183, 186
cubensis, II, 184, 188
extensus, II, 184, 190, 191
guatemalensis, II, 183, 184
lemairei, II, 184, 189, 194,
 226; IV, 282, 283
minutiflorus, II, 195, 196
monacanthus, II, 184, 190,
 226
napoleonis, II, 184, 191; IV,
 282
ocamponis, II, 126, 183, 184,
 185
polyrhizus, II, 183, 185, 226
purpusii, II, 183, 184
stenopterus, II, 184, 190
triangularis, II, 184, 185,
 191, 192, 193; IV, 282
tricotatus, II, 187; IV, 282
trigonus, II, 184, 192, 193;
 IV, 282

Hylocereus—*continued*,
 undatus, II, 184, 187, 188,
 189; IV, 281
venezuelensis, II, 183, 186,
 226
Imbricatae, I, 44, 60
Inamoena, I, 45, 125
Inarmatae, I, 208; IV, 180
Iniabanto, I, 19
Ipomoea, II, 92
Jaatoc, II, 88
Jaramataca, II, 111
Jasminocereus, II, 2, 146, 147
galapagensis, II, 146, 147
Joconostle, II, 93
Junco, II, 119
Junco espinoso, II, 119
Kadoesji, II, 18, 88
La bande de sud, II, 151
Lanuginosi, II, 59
Latispineae, III, 109
Lemaireocereus, I, 116; II, 2,
 25, 43, 69, 85-103, 114,
 135, 151; IV, 268, 273,
 274
aragonii, II, 8, 92, 93, 103
benckeii, IV, 273, 274
cartwrightianus, II, 8, 100
chende, II, 8, 90, 91; IV, 273
chichi, II, 85, 89
cumengei, II, 116
deficiens, II, 85, 94, 96
dumortieri, II, 85, 102
eichlamii, II, 85, 89, 90; IV,
 273
eruca, II, 115, 116
godingianus, II, 85, 91, 92,
 135
griseus, II, 20, 85, 87, 88, 89
 103, 225; IV, 273
gummosus, II, 116, 117
hollianus, II, 85, 86; IV, 273
humilis, II, 85, 100, 101
hystrix, II, 46, 85, 86, 87,
 225; IV, 273
laetus, II, 85, 99, 100
longispinus, II, 85, 89, 90
mixtecensis, II, 89, 91
montanus, II, 85, 97
pruinosis, II, 85, 88, 89
queretarocensis, II, 85, 96, 97
schumannii, II, 103
stellatus, II, 85, 92, 93, 94,
 169
thurberi, II, 85, 96, 97, 98;
 IV, 274
treleasei, II, 85, 93, 95, 224
weberi, II, 85, 95, 97, 164;
 IV, 273
Lemon vine, I, 10
Lengua de vaca, I, 164
Leocereus, II, 2, 108-110, 175,
 225
bahiensis, II, 108, 109
glaziovii, II, 108, 109
melannurus, II, 108, 109
Leon, I, 96
Leoncito, I, 96
Lepismium, IV, 208, 215, 216
alternatum, IV, 237
anceps, IV, 215
cavernosum, IV, 215, 216
cavernosum ensiforme, IV,
 215
cavernosum minus, IV, 216
commune, IV, 215, 216
cruciforme, IV, 215
dissimilis, IV, 236
duprei, IV, 216
fluminense, IV, 243
knightii, IV, 215, 216
laevigatum, IV, 216
mittleri, IV, 216
myosurus, IV, 215, 216
myosurus knightii, IV, 215
myosurus laevigatum, IV,
 215
paradoxum, IV, 237

Lepismium—continued,
 radicans, IV, 215, 216
 ramosissimum, IV, 216
 sarmentaceum, IV, 230
 tenue, IV, 213
Leptocaulis, I, 44, 46, 49
Leptocereus, II, 2, 77-82, 104,
 122; IV, 281
 arboreus, II, 77, 80
 assurgens, II, 77, 79, 80
 leonii, II, 77, 78, 79; IV, 272
 maxonii, II, 77, 80
 prostratus, II, 7, 79
 quadricostatus, II, 77, 81
 sylvestris, II, 77, 80, 81
 weingartianus, II, 77, 82
Leucorhaphes, IV, 221
Leucotrichae, I, 174
Leuchtenbergia, III, 78, 107-
 109
 principis, III, 107, 108
 Living rock, III, 83
Lobivia, III, 3, 49-60, 78, 179
 andalgalensis, III, 49, 55,
 56, 58
 boliviensis, III, 49, 52
 bruchii, III, 49, 50
 cachensis, III, 49, 52
 caespitosa, III, 49, 53
 chionanthus, III, 49, 58, 179
 cinnabarina, III, 40, 54
 corbula, III, 49, 56
 cumingii, III, 49, 59, 117, 177
 famatimensis, IV, 286
 ferox, III, 49, 50, 51, 52
 grandiflora, III, 49, 57
 grandis, III, 58
 haematantha, III, 49, 57
 lateritia, III, 49, 56
 longispina, III, 49, 51, 52
 pampana, III, 49, 56
 pentlandii, III, 49, 54, 55;
 IV, 285
 saltensis, III, 49, 53
 shaferei, III, 40, 52, 55
 thionanthus, III, 49, 57, 179
Lophocereus, II, 3, 170, 177,
 178, 179
 australis, II, 177; IV, 280
 sargentianus, II, 177
 schottii, II, 177, 178, 179;
 IV, 279, 280
Lophophora, III, 77, 83-85, 93
 lewinii, III, 84, 8
 williamsii, III, 84, 184; IV,
 286
 williamsii lewinii, III, 84
Loranthus aphyllus, I, 79
Lorentzianae, IV, 221
Machaerocereus, II, 2, 114-117
 eruca, II, 114, 115, 116; IV,
 276
 gummosus, II, 114, 116, 117;
 IV, 276
Macromeres, IV, 24
Maihuen, I, 40
Maihuenia, I, 8, 24, 40-42,
 45, 95
 brachydelphys, I, 41, 42; IV,
 175, 253
 patagonica, I, 41
 philippii, I, 41
 poeppigii, I, 41, 42; IV, 253
 tehuelches, I, 41, 42
 valentinii, I, 40, 41, 42
Malacocarpus, III, 78, 94, 96,
 138, 167, 177, 178, 187-
 207, 208, 216
 aciculatus, III, 198
 acutatus, III, 198
 apricus, III, 187, 192
 catamarcensis, III, 187, 197
 concinnus, III, 187, 192, 193;
 IV, 289
 corynodes, III, 198, 199
 corynodes erinaceus, III, 198,
 199
 courantii, III, 188

Malacocarpus—continued,
 curvispinus, III, 188, 203
 erinaceus, III, 187, 198, 199;
 IV, 289
 escayachensis, III, 188, 205
 graessneri, III, 188, 205
 grossei, III, 187, 190
 haselbergii, III, 188, 201,
 205; IV, 289
 heptacanthus, III, 218
 islayensis, III, 188, 201
 langsdorffii, III, 188, 199
 leninghausii, III, 188, 204;
 IV, 289
 linkii, III, 187, 195; IV, 289
 maassii, III, 188, 202
 mammillarioides, III, 188,
 203
 mammosus, III, 188, 200;
 IV, 289
 martinii, III, 188
 muricatus, III, 187, 194
 napinus, III, 187, 190, 191
 nigrispinus, III, 187, 190
 ottonis, III, 187, 195, 196;
 IV, 289
 patagonicus, III, 187, 197,
 198
 polyacanthus, III, 198, 200
 pulcherrimus, III, 187, 194
 reichei, III, 187, 191
 schumannianus, III, 187,
 189, 190
 scopa, III, 187, 193; IV, 289
 sellowianus, III, 188
 sellowianus tetracanthus, III,
 188
 sellowii, III, 188, 189
 sellowii tetracanthus, III,
 188
 strausianus, III, 188, 201
 tabularis, III, 187, 192, 193
 tephacanthus, III, 187, 188,
 189, 191 IV, 288
 tetracanthus, III, 188
 tuberisulcatus, III, 188, 202
Malus, I, 9
Mamillaria, IV, 6
Mamillopsis, IV, 3, 19, 20
 diguetii, IV, 19, 20
 senilis, IV, 19, 20
Mammariella, IV, 124
Mammillaria, IV, 6, 5
Mammillaria, I, 4; II, 3, 118;
 III, 80, 83, 84, 90, 91,
 92, 93, 108, 110, 208,
 210, 237; IV, 3, 10, 14,
 16, 19, 21, 24, 34, 48, 65,
 79, 107, 155, 163, 166,
 172, 175
 acanthophlegma, IV, 101,
 107, 108
 acanthophlegma abducta, IV,
 107
 acanthophlegma decandollii,
 IV, 107
 acanthophlegma elegans, IV,
 107
 acanthophlegma leucocephala,
 IV, 107
 acanthophlegma meisneri, IV,
 107
 acanthophlegma monacantha,
 IV, 107
 acanthostephes, IV, 41
 acanthostephes recta, IV, 41
 acicularis, IV, 171
 aciculata, IV, 88, 132
 actinoplea, IV, 172
 adunca, IV, 140
 aeruginosa, IV, 88
 affinis, IV, 88
 aggregata, III, 14; IV, 47
 albida, IV, 132, 133
 albiseta, IV, 171, 172
 aloidaea pulviligera, III, 81
 abides, III, 81
 alpina, IV, 63, 165

Mammillaria—continued,
 alversonii, IV, 46, 47
 amabilis, IV, 172
 ambigua, III, 98; IV, 175
 amoena, IV, 120, 167
 anancistraria, IV, 131
 ancistracantha, IV, 15
 ancistrata, IV, 150
 ancistria, IV, 131
 ancistrina, IV, 10
 ancistroides, IV, 149, 150
 ancistroides inuncinata, IV,
 132
 ancistroides major, IV, 150
 andreae, IV, 121
 anguinea, IV, 134
 angularis, IV, 88, 90, 92
 angularis Compressa, IV, 91,
 92
 angularis fulvescens, IV, 92
 angularis fulvispina, IV, 91
 angularis longiseta, IV, 91
 angularis rufispina, IV, 92
 angularis triacantha, IV, 91
 anisacantha, IV, 103, 105
 applanata, IV, 76
 areolesa, III, 80
 argentea, IV, 172
 arida, IV, 73
 arietina, IV, 41
 arietina spinosior, IV, 41
 arizonica, IV, 43, 46, 48
 armillata, IV, 157
 aselliformis, IV, 59
 asterias, IV, 31
 asteriflora, IV, 172
 atrata, III, 97, 98; IV, 122
 atrorubra, IV, 172
 atrosanguinea, IV, 172
 aulacantha, IV, 171
 aulacothele, IV, 30, 31
 aulacothele flavispina, IV, 30
 aulacothele multispina, IV,
 30
 aulacothele spinosior, IV, 30
 aulacothele sulcimamma, IV,
 30
 aurata, IV, 122
 aurea, IV, 123
 aureiceps, IV, 114
 auricoma, IV, 117
 aureora, IV, 118
 autumnalis, IV, 92
 badispina, IV, 172
 barbata, IV, 144, 145
 harlowii, IV, 172
 beguinii, IV, 16
 bellatula, IV, 165
 benckeii, IV, 171
 bergeana, IV, 150
 bergeni, IV, 172
 bergii, IV, 165
 besleri, III, 220
 bicolor, IV, 98, 99, 168
 bicolor cristata, IV, 98
 bicolor longispina, IV, 98
 bicolor nivea, IV, 98
 bicolor nobilis, IV, 98, 99
 bifurca, IV, 172
 biglandulosa, IV, 30
 bihamata, IV, 140
 binops, IV, 172
 bocasana, IV, 147
 bocasana cristata, IV, 147,
 148
 bocasana glochidiata, IV, 147
 bocasana kunzeana, IV, 145
 bocasana sericata, IV, 147
 bocasana splendens, IV, 147
 bocasiana, IV, 171
 bockii, IV, 77
 boederiana, IV, 154
 bombycina, IV, 50, 161
 borealis, IV, 45
 boucheana, IV, 79
 brachydelphys, IV, 175, 253
 brandegeei, IV, 65, 73, 74
 brandi, IV, 171

Mammillaria—continued,
 brandtii, IV, 172
 brevimmamma, IV, 31
 brevimmamma exsudans, IV,
 31
 breviseta, IV, 172
 brongniartii, IV, 109
 brownii, IV, 25, 33, 34
 bruennowii, IV, 172
 buchheimeana, IV, 75
 bumamma, IV, 33, 38,
 bussleri, IV, 26, 27
 caesia, IV, 118
 caespititia, IV, 126, 165
 caespitosa, III, 25, 26; IV,
 52, 134
 calcarata, IV, 48, 49
 calochlora, IV, 50
 camptotricha, IV, 61, 126,
 127
 candida, IV, 130
 candida rosea, IV, 130
 canescens, IV, 17, 133
 cantera, IV, 172
 caput-medusae, IV, 86, 87
 caput-medusae centrispina,
 IV, 86
 caput-medusae crassior, IV,
 86
 caput-medusae hexacantha,
 IV, 87
 caput-medusae tetracantha,
 IV, 86, 87; IV, 70
 caracassana, IV, 70
 caracassana, IV, 70, 71
 carnea, IV, 87, 88, 89, 102
 carnea aeruginosa, IV, 88
 carnea cirrosa, IV, 88
 carnea villifera, IV, 102
 carretii, IV, 149, 151, 160
 castaneoides, IV, 118
 cataphracta, IV, 88
 caudata, IV, 175
 celsiana, IV, 99, 112, 113
 celsiana guatemalensis, IV,
 115
 celsiana longispina, IV, 172
 centa, IV, 171
 centricirra, IV, 41, 77, 78,
 79, 80, 169
 centricirra amoena, IV, 79
 centricirra arietina, IV, 79
 centricirra bockii, IV, 78,
 79
 centricirra boucheana, IV,
 79
 centricirra ceratophora, IV,
 79
 centricirra cirrhosa, IV, 79
 centricirra conopsea, IV, 79
 centricirra cristata, IV, 79
 centricirra deflexispina, IV,
 79
 centricirra destorum, IV,
 79
 centricirra tampico, IV,
 79
 centricirra diacantha, IV,
 79
 centricirra diadema, IV, 79
 centricirra divaricata, IV,
 79
 centricirra divergens, IV,
 78, 79
 centricirra ehrenbergii, IV,
 79
 centricirra falcata, IV, 79
 centricirra flaviflora, IV, 63
 centricirra foersteri, IV, 79
 centricirra gebweileriiana,
 IV, 79
 centricirra gladiata, IV, 79
 centricirra glauca, IV, 79
 centricirra globosa, IV, 79
 centricirra grandidens, IV,
 79
 centricirra guileminiana,
 IV, 79

Mammillaria—*continued*,
centricirrho hopferiana, IV, 77
centricirrho hopferiana, IV, 79
centricirrho hystrix, IV, 79
centricirrho hystrix grandicornis, IV, 79
centricirrho hystrix longispina, IV, 79
centricirrho jorderi, IV, 79
centricirrho krameri, IV, 78, 79
centricirrho krameri longispina, IV, 79
centricirrho krausei, IV, 79
centricirrho lactescens, IV, 79
centricirrho lehmannii, IV, 79
centricirrho longispina, IV, 79
centricirrho macracantha, IV, 79, 80
centricirrho macrothele, IV, 77, 91
centricirrho magnimamma, IV, 78, 79
centricirrho megacantha, IV, 79
centricirrho microceras, IV, 79
centricirrho montsii, IV, 79
centricirrho moritziana, IV, 79
centricirrho neumanniana, IV, 79
centricirrho nordmannii, IV, 79
centricirrho obconella, IV, 79
centricirrho pachythele, IV, 179
centricirrho pazzanii, IV, 79
centricirrho pentacantha, IV, 79
centricirrho polygona, IV, 79
centricirrho posteriana, IV, 79
centricirrho pulchra, IV, 79
centricirrho recurva, IV, 78, 79
centricirrho schiedeana, IV, 79
centricirrho schmidtii, IV, 79
centricirrho spinosior, IV, 79
centricirrho subcurvata, IV, 79
centricirrho tetracantha, IV, 79
centricirrho uberimamma, IV, 79
centricirrho valida, IV, 79
centricirrho versicolor, IV, 79
centricirrho viridis, IV, 79
centricirrho zooderi, IV, 7
centricirrho zuccariniana, IV, 79
centrispina, IV, 95
cephalophora, IV, 41, 128
ceratites, IV, 16
ceratocentra, IV, 32
ceratophora, IV, 77
chapinensis, IV, 100
childsi, IV, 175
chinocephala, IV, 101, 107
chlorantha, IV, 43, 46
chrysacantha, IV, 121
chrysacantha fuscata, IV, 122
chrysantha, IV, 171
circumtexta, IV, 171
cirrhifera, IV, 90, 92, 94
cirrhifera albispina, IV, 91

Mammillaria—*continued*,
cirrhifera angulosior, IV, 90
cirrhifera fulvispina, IV, 91
cirrhifera longispina, IV, 92
cirrhifera major, IV, 91
cirrhosa, IV, 78
cirrifer longisetia, IV, 91
cirrosa, IV, 78
citrina, IV, 172
clava, IV, 30
clavata, IV, 15
closiana, IV, 172
coccinea, III, 79
collina, IV, 111
columbiana, IV, 127
columnaris, IV, 88, 107
columnaris minor, IV, 88
communis, III, 225
compacta, IV, 36, 109
compressa, IV, 42, 90
confinis, IV, 133
conica, IV, 70, 165
coniflora, IV, 133
conimamma, IV, 37, 38
conimamma major, IV, 38
conoidea, IV, 14, 17, 18
conopsea, IV, 77, 79
conopsea longispina, IV, 77
conspicua, IV, 108
contacta, IV, 172
convoluta, IV, 167
corbula, III, 56
cordigera, IV, 50, 161
corioides, IV, 175
cornifera, IV, 39
cornifera impexicoma, IV, 36, 37
cornifera mutica, IV, 39
cornimamma, IV, 37
cornuta, IV, 50
corollaria, IV, 172
coronaria, IV, 41, 141, 167, 169, 171, 176
coronaria minor, IV, 176
coronata, IV, 172
coryphides, IV, 172
crassispina, IV, 121, 122
crassispina gracilior, IV, 121
crassispina rufa, IV, 122
crebrispina, IV, 18
crebrispina nitida, IV, 172
criniformis, IV, 150, 151
criniformis albida, IV, 150
criniformis rosea, IV, 150
crinifera, IV, 172
crinita, IV, 150
crinita pauciseta, IV, 151
crocidata, IV, 87
crocidata quadrispina, IV, 87
crucigera, IV, 113
cubensis, IV, 48
cuneiflora, IV, 172
cunendstiana, IV, 171
curvata, IV, 31
curvispina, IV, 132, 172
curvispina parviflora, IV, 132
cylindraca, IV, 172
cylindraca, IV, 127
cylindrica flavispina, IV, 127
dactylithele, IV, 25
daedalea, IV, 98, 99, 168
daedalea viridis, IV, 86, 172
daimonoceras, IV, 36
dasyacantha, IV, 55, 57
dealbata, IV, 110
decholara, IV, 171
decipiens, IV, 131, 150
decipiens rosea, IV, 150
declivis, IV, 76
decora, IV, 173
decora obscura, IV, 173
dedalea viridis, IV, 86
deficiens, IV, 132
deficum, IV, 132
deflexispina, IV, 77
degrandii, IV, 172
delactiana, IV, 39, 40, 50

Mammillaria—*continued*,
deleuli, IV, 172
densa, IV, 134, 135
depressa, IV, 133, 140
deserti, IV, 46, 47
desertorum, IV, 172
destorum, IV, 79
de tampico, IV, 79
diacantha, IV, 86, 110
diacantha nigra, IV, 110
diacentra, IV, 166
diadema, IV, 77
diaphanacantha, IV, 17
dichotoma, IV, 175
difficilis, IV, 28
digitalis, IV, 167
dioica, IV, 158
dioica insularis, IV, 140
disciformis, III, 106
discolor, IV, 132, 133
discolor aciculata, IV, 132
discolor albida, IV, 132
discolor breviflora, IV, 133
discolor coniflora, IV, 133
discolor curvispina, IV, 132
discolor fulvescens, IV, 133
discolor monstrosa, IV, 132
discolor nigricans, IV, 174
discolor nitens, IV, 132
discolor prolifera, IV, 132, 133
discolor pulchella, IV, 132
discolor rhodacantha, IV, 133
divaricata, IV, 77, 172
divergens, IV, 77
dolichacantha, IV, 107
dolichocentra, IV, 106, 107
dolichocentra brevispina, IV, 106
dolichocentra galeottii, IV, 105
dolichocentra phaeacantha, IV, 105
dolichocentra picta, IV, 107
dolichocentra staminea, IV, 106
donatii, IV, 111
donkelaari, IV, 172
droegeana, IV, 10
dubia, IV, 172
dumetorum, IV, 128
durangensis, IV, 42
dyckiana, IV, 107, 108
eborina, IV, 173
eburnea, IV, 98
echinaria, IV, 136
echinata, IV, 136
echinata densa, IV, 134
echinata gracilior, IV, 136
echinata pallida, IV, 136
echinocactoides, IV, 17
echinoidea, IV, 30
echinops, IV, 102, 172
echinus, IV, 42
ehrenbergii, IV, 77
eichlamii, IV, 94, 95
elegans, IV, 101, 107, 108, 109, 113
elegans dealbata, IV, 110
elegans globosa, IV, 107
elegans klugii, IV, 107
elegans micrantha, IV, 107
elegans minor, IV, 107
elephantidens, IV, 32, 33, 38
elephantidens bumamma, IV, 33
elephantidens spinosissima, IV, 33
elongata, III, 80; IV, 134, 135, 170
elongata anguinea, IV, 134
elongata centrispina, IV, 136
elongata echinata, IV, 136
elongata intertexta, IV, 134
elongata minima, IV, 135
elongata rufescens, IV, 135
elongata rufocrocea, IV, 134

Mammillaria—*continued*,
elongata stella-aurata, IV, 134
elongata straminea, IV, 135
elongata subcrocea, IV, 134
elongata tennis, IV, 134
emskotteriana, IV, 57
emundtsiana, IV, 173
enneacantha, IV, 172
erecta, IV, 32
erectacantha, IV, 173
eriacantha, IV, 127
eriantha, IV, 81, 121, 122
eriantha, IV, 127
erythrosperma, IV, 151
erythrosperma similis, IV, 151, 152
euchlora, IV, 173
eugenia, IV, 169
evanescentis, IV, 32
evarescens, IV, 32
evarescens, IV, 32
evarescentis, IV, 32
eximia, IV, 118, 119
exsudans, IV, 31
falcata, IV, 78
farinosa, IV, 173
fasciculata, IV, 162, 163
fellneri, IV, 173
fennelii, IV, 159, 160
fertilis, IV, 131
fischeri, IV, 95
fissurata, III, 183; IV, 286
flava, IV, 173
flavescens, III, 212; IV, 166, 171, 172
flaviceps, IV, 122
flavicomma, IV, 166
flavovirens, IV, 85, 86
flavovirens cristata, IV, 86
floccigera, IV, 122
floccigera longispina, IV, 122
floribunda, III, 97, 98
foersteri, IV, 77
fordii, IV, 158
formosa, IV, 90
formosa discipula, IV, 90
formosa gracilispina, IV, 90
formosa laevior, IV, 90
formosa microthele, IV, 90
foveolata, IV, 83
fragilis, IV, 133, 134
fuliginosa, IV, 71
fulvispina, IV, 121, 122, 123, 176
fulvispina media, IV, 176
fulvispina minor, IV, 176
fulvispina pyrrhocentra, IV, 123
fulvispina rubescens, IV, 121
fulvolanata, IV, 172
funckii, IV, 92
furfuracea, III, 80
fuscata, IV, 119, 121
gabbii, IV, 73, 74
galeottii, IV, 105, 106, 107, 154
gebweilleriana, IV, 78
geminata, IV, 173
geminiflora, IV, 172
geminispina, IV, 98, 107, 108
geminispina monacantha, IV, 113
geminispina tetracantha, IV, 107
gibbosa, III, 98; IV, 173
gigantea, IV, 85
gigantothele, IV, 63
glabrata, IV, 173
glabrescens, IV, 172
gladiata, IV, 77
glanduligera, IV, 31
glauca, IV, 77
globosa, IV, 63
glochidiata, IV, 149, 150, 151
glochidiata alba, IV, 150
glochidiata albida, IV, 150
glochidiata aurea, IV, 143

Mammillaria—*continued*,
 glochidiata crinata, IV, 150
 glochidiata inuncinata, IV, 131
 glochidiata prolifera IV, 150
 glochidiata purpurea, IV, 149
 glochidiata rosea IV, 150
 glochidiata sericata, IV, 148, 149
 glomerata, IV, 124
 goeringii, IV, 172
 goerngi, IV, 49
 golziana, IV, 26, 27
 goodrichii, IV, 158
 goodridgei, IV, 156, 158, 159
 goodridgii, IV, 158
 gracilis, IV, 134, 136
 gracilis lactevirens, IV, 136
 gracilis pulchella, IV, 134, 136
 gracilis virens, IV, 136
 graessneriana, IV, 117
 grahamii, III, 43; IV, 60, 144, 154, 155, 156
 grahamii arizonica, IV, 155
 grahamii californica, IV, 156
 grandicornis, IV, 79, 173
 grandidens, IV, 79
 grandiflora, IV, 17, 18, 131
 grandis, IV, 172
 granulata, IV, 125
 grayhamii, IV, 156
 greggii, III, 93
 grisea, IV, 166
 grusonii, IV, 55
 grusonii similis, IV, 172
 guanajuatensis, IV, 85
 guebwilleriana, IV, 172
 guerkeana, IV, 29, 30
 guillemianiana, IV, 131
 gummifera, IV, 74
 haagiana, IV, 110
 haagiana validior, IV, 110
 haematocacta, IV, 173
 halei, IV, 21, 22
 hamata, IV, 140, 141
 hamata brevispina, IV, 141
 hamata longispina, IV, 141
 hamata principis, IV, 141
 hamuligera, IV, 151
 haseloffii, IV, 20, 118
 haynei, IV, 166
 haynii, IV, 166
 haynii minima, IV, 167
 haynii viridula, IV, 166
 heeseana, IV, 94
 heeseana brevispina, IV, 94
 heeseana longispina, IV, 94
 heinei, IV, 166
 helicteres, IV, 167
 hemisphaerica, IV, 75
 hepatica, IV, 117
 hermantiana, IV, 172
 hermannii, IV, 118
 hermannii flavicans, IV, 118
 heteracantha, IV, 168
 heteracentra, IV, 172
 heteromorpha, III, 83; IV, 25
 hevernickii, IV, 172
 hexacantha, IV, 167
 hexacentra, IV, 63
 heyderi, IV, 75, 76, 167
 heyderi applanata, IV, 76
 heyderi hemisphaerica, IV, 75, 76
 hidalgensis, IV, 88
 hirschtiana, IV, 43
 hirsuta, IV, 146
 hochderferi, IV, 171
 hoffmannseggii, III, 98
 hopferiana, IV, 77
 horripila, IV, 16
 humboldtii, IV, 130
 humilior, IV, 15
 hybrida, IV, 122
 hystrix, IV, 77, 79

Mammillaria—*continued*,
 imbricata, IV, 121
 impexicoma, IV, 36
 inclinis, IV, 96
 inconspicua, IV, 17
 incurva, IV, 173
 intertexta, IV, 134
 intertexta rufocrocea, IV, 135
 intricata, IV, 172
 inuncinata, IV, 132
 inuncta, IV, 121
 irregularis, IV, 167
 isabellina, IV, 118
 jalappensis, IV, 105
 joossensiana, IV, 167
 jorderi, IV, 79
 jucunda, IV, 173
 karstenii, IV, 71
 karwinskiana, IV, 95, 96, 97
 karwinskiana centrispina, IV, 95
 karwinskiana flavescens, IV, 95
 karwinskiana virens, IV, 95
 kewensis, IV, 104, 105
 kewensis albispina, IV, 105
 kleinii, IV, 173
 kleinschmidtiana, IV, 91
 klennerii, IV, 171
 klugii, IV, 107
 knippeliana, IV, 96
 krameri, IV, 77, 94
 krameri viridis, IV, 78
 krauseana, IV, 93
 kunthii, IV, 107
 kunzeana, IV, 145, 146
 lactescens, IV, 78
 laeta, IV, 63
 lamprochaeta, IV, 173
 lamuligera, IV, 151
 lanifera, IV, 113, 122, 123
 lapaixi, IV, 172
 lasiacantha, IV, 124, 128
 lasiacantha denudata, IV, 129
 lasiacantha minor, IV, 128
 lasiandra denudata, IV, 129
 lassauieri, IV, 167
 lassonneriei, IV, 167
 latimamma, IV, 41
 latispina, III, 143
 lehmannii, IV, 30, 31, 79
 lehmannii sulcimamma, IV, 30
 lenta, IV, 129, 130
 leona, IV, 136, 137, 138
 lesaunieri, IV, 167
 leucacantha, IV, 30, 31
 leucantha, IV, 31
 leucocarpa, IV, 93
 leucocentra, IV, 167
 leucocephala, IV, 108
 leucodasys, IV, 173
 leucodictia, IV, 173
 leucotricha, IV, 92
 lewinii, III, 84
 lindheimeri, IV, 76
 linkeana, IV, 118
 linkei, IV, 20
 littoralis, IV, 138
 livida, IV, 173
 longihamata, IV, 22
 longimamma, IV, 61, 62, 63
 longimamma compacta, IV, 63
 longimamma congesta, IV, 62, 63
 longimamma cristata, IV, 63
 longimamma gigantothele, IV, 62, 63
 longimamma globosa, IV, 63
 longimamma hexacentra, IV, 62
 longimamma laeta, IV, 63
 longimamma ludwigii, IV, 63

Mammillaria—*continued*,
 longimamma luteola, IV, 63
 longimamma malaena, IV, 63
 longimamma major, IV, 63
 longimamma melaleuca, IV, 63
 longimamma pseudo-melaleuca, IV, 63
 longimamma sphaerica, IV, 63
 longimamma spinosior, IV, 63
 longimamma uberiformis, IV, 63
 longisetia, IV, 90
 longispina, IV, 79, 107
 loricata, IV, 168
 ludwigii, IV, 76
 lutescens, IV, 172
 macdougallii, IV, 74
 macdowellii, IV, 85
 macracantha, IV, 79, 80
 macrantha, IV, 80
 macromeris, IV, 25, 26
 macromeris longispina, IV, 25
 macromeris nigrispina, IV, 25
 macrothele, IV, 30, 31
 macrothele biglandulosa, IV, 30
 macrothele lehmannii, IV, 30
 macrothele nigrispina, IV, 31
 maelenii, IV, 80
 magnimamma, IV, 41, 77, 128
 magnimamma arietina, IV, 41
 magnimamma lutescens, IV, 41
 magnimamma spinosior, IV, 41
 mainae, IV, 154
 mallettiana, IV, 99
 mammillaris, IV, 70
 martiana, IV, 30
 maschalacantha, IV, 92, 93
 maschalacantha dolichacantha, IV, 93
 maschalacantha leucotricha, IV, 92
 maschalacantha xantotricha, IV, 93
 mazatlanensis, IV, 138
 megacantha, IV, 77
 megacantha rigidior, IV, 77
 meiacantha, IV, 84
 meionacantha, IV, 84
 meisneri, IV, 107
 melaleuca, IV, 63
 meianacantha, IV, 173
 maianocentra, IV, 81
 meonacantha, IV, 84
 mercadensis, IV, 145
 meschalacantha, IV, 93
 micans, IV, 173
 micracantha, IV, 71, 173
 micrantha, IV, 71
 microcarpa, IV, 155, 156
 microceras, IV, 77
 microdasys, IV, 172
 micromeris, III, 92, 93; IV, 6, 109, 173
 micromeris fuligifera, III, 93
 micromeris greggii, III, 93
 microthele, IV, 71, 108, 109
 microthele brongniartii, IV, 109
 minima, IV, 134, 135
 miqueliana, IV, 172
 mirabilis, IV, 118
 missouriensis, IV, 51, 52, 53
 missouriensis caespitosa, IV, 52
 missouriensis nuttallii, IV, 53
 missouriensis robustior, IV, 52

Mammillaria—*continued*,
 missouriensis similis, IV, 52
 missouriensis viridescens, IV, 53
 mitis, IV, 175
 monancistraria, IV, 148
 monocentra, IV, 168
 monoclova, IV, 37
 monothele, IV, 172
 montana, IV, 45
 montsii, IV, 79
 morini, IV, 172
 moritziana, IV, 79
 mucronata, IV, 173
 muelhlenpfordtii, IV, 112
 multicolor, IV, 172
 multiceps, IV, 125, 126
 multiceps elongata, IV, 125
 multiceps grisea, IV, 125
 multiceps humilis, IV, 125
 multiceps perpusilla, IV, 125
 multiradiata, IV, 171
 multiseta, IV, 173
 multihamata, IV, 146
 mundtii, IV, 112
 mutabilis, IV, 92, 93, 94
 mutabilis autumnalis, IV, 93
 mutabilis laevis, IV, 93
 mutabilis longispina, IV, 94
 mutabilis xanthotricha, IV, 92
 mystax, IV, 92, 94
 napina, IV, 104
 neglecta, IV, 123
 neo-mexicana, IV, 45
 nervosa cristata, IV, 168
 nervosus cristatus, IV, 168
 neumanni glabrescens, IV, 78
 neumanniana, IV, 77
 nicholsonii, IV, 168
 nickelsae, IV, 35
 nickelsi, IV, 172
 nigerrima, IV, 172
 nigra, IV, 171
 nigricans, IV, 1, 6, 117
 nitens, IV, 132
 nitida, IV, 165
 nivea, IV, 98
 nivea cristata, IV, 99
 nivea daedalea, IV, 98
 nivea longispina, IV, 99
 nivea wendlei, IV, 99
 nivosa, IV, 71, 72
 nobilis, IV, 98, 99
 nogalensis, IV, 28
 nordmannii, IV, 79
 notesteinii, IV, 53
 nuda, IV, 168
 numina, IV, 172
 nuttallii, IV, 52, 53
 nuttallii borealis, IV, 53
 nuttallii caespitosa, IV, 52
 nuttallii robustior, IV, 52
 obconella, II, 102; IV, 79, 106, 107
 obconella galeottii, IV, 105
 obliqua, IV, 173
 obscura, IV, 87
 obscura galeottii, IV, 106
 obvallata, IV, 174
 ocamponis, IV, 145
 octacantha, IV, 30
 odieriana, IV, 121, 123
 odieriana aurea, IV, 123
 odieriana cristata, IV, 122
 odieriana rigidior, IV, 122
 odieriana rubra, IV, 122
 oettingenii, IV, 91
 olivacea, IV, 123
 oliviae, IV, 135
 olorina, IV, 174
 oothele, IV, 102, 174
 ottonis, IV, 26, 27
 ottonis tenuispina, IV, 26
 ovimamma, IV, 174
 ovimamma brevispina, IV, 174
 ovimamma oothele, IV, 174

Mammillaria—continued,
 pachytele, IV, 78
 painteri, IV, 151, 152
 pallescens, IV, 88
 palmeri, IV, 172
 papyracantha, III, 91
 parkinsonii, IV, 98
 parkinsonii rubra, IV, 98
 parkinsonii waltonii, IV, 98
 parmentieri, IV, 171
 parvimamma, IV, 70
 parvissima, IV, 126
 pazzanii, IV, 77
 peacockii, IV, 110
 pectinata, IV, 34, 35, 175
 pectinata cristata, IV, 34
 pectinifera, IV, 64
 pentacantha, IV, 77
 perbella, IV, 108, 111
 perote, IV, 110
 perpusilla, IV, 126
 perringii, IV, 112, 113
 persicina, IV, 174
 petrophila, IV, 73
 petterssonii, IV, 94
 pfeifferana, IV, 39
 pfeifferi, IV, 121, 123
 pfeifferi altissima, IV, 121
 pfeifferi dichotoma, IV, 121
 pfeifferi flaviceps, IV, 121
 pfeifferi fulvispina, IV, 121
 pfeifferi variabilis, IV, 121
 phaeacantha, IV, 116, 117, 122
 phaeacantha rigidior, IV, 117
 phaeotrica, IV, 174
 phellosperma, IV, 60, 65, 156
 phymatothele, IV, 76
 picta, IV, 169
 picturata, IV, 176
 pilispina, IV, 14
 plaschnicki, IV, 30
 plaschnickii straminea, IV, 30
 plecostigma, IV, 169
 plecostigma major, IV, 169
 plecostigma minor, IV, 169
 pleiocephala, IV, 74
 plinthimorpha, IV, 169
 plumosa, IV, 123, 124
 polia, IV, 172
 polyacantha, IV, 117
 polyactina, IV, 117
 polycentra, IV, 117
 polycephala, IV, 113
 polychlora, IV, 18
 polyedra, IV, 101, 102, 103
 polyedra anisacantha, IV, 103
 polyedra laevior, IV, 102, 103
 polyedra scleracantha, IV, 102, 103
 polyedra spinosior, IV, 102
 polygona, IV, 101, 105
 polymorpha, IV, 24, 30, 31, 174
 polythele, IV, 88, 168
 polythele aciculata, IV, 88, 132
 polythele columnaris, IV, 88
 polythele hexacantha, IV, 88
 polythele latimamma, IV, 88
 polythele quadrispina, IV, 88
 polythele setosa, IV, 88
 polytricha, IV, 102
 polytricha hexacantha, IV, 102
 polytricha tetracantha, IV, 102
 pomacea, IV, 117
 pondii, IV, 23
 porphyracantha, IV, 174
 poselgeri, IV, 22
 poselgeriana, IV, 118, 119

Mammillaria—continued,
 posteriana, IV, 79
 potosiana, IV, 15, 24, 50
 potosina, IV, 99
 potosina longispina, IV, 99
 pottsii, IV, 136, 137
 pottsii vera, IV, 137
 praelii, IV, 96
 pretiosa, IV, 117
 pretiosa cristata, IV, 118
 principis, IV, 141
 pringlei, IV, 115
 prismatica, III, 80, 81
 procera, IV, 174
 prolifera, IV, 70, 124, 125
 pruinosa, IV, 118
 pseudofusca, IV, 119, 120
 pseudomammillaris, IV, 132
 pseudoperbella, IV, 108, 109
 pseudoperbella rufispina, IV, 109
 pugionacantha, IV, 174
 pulchella, IV, 132
 pulchella nigricans, IV, 132
 pulcherrima, IV, 117, 172
 pulchra, IV, 121, 123, 169
 pulvilligera, III, 81
 punctata, IV, 174
 purpuracea, III, 80
 purpurascens, IV, 174
 purpurea, IV, 174
 purpusii, III, 90, 91
 pusilla, IV, 57, 70, 71, 124, 125
 pusilla caespititia, IV, 126
 pusilla cristata, IV, 125
 pusilla haitiensis, IV, 71, 124
 pusilla major, IV, 124
 pusilla mexicana, IV, 126
 pusilla minor, IV, 124
 pusilla multiceps, IV, 57
 pusilla texana, IV, 125, 126
 pycnacantha, IV, 34, 40, 41, 42
 pycnacantha scepontocentra, IV, 41
 pycnacantha spinosior, IV, 41
 pyramidalis, IV, 122
 pyrrhacantha, IV, 172
 pyrrhacantha pallida, IV, 172
 pyrrhocacantha, IV, 121
 pyrrhocentra, IV, 123
 pyrrhocentra gracilior, IV, 123
 pyrrhocentra, IV, 99, 100, 102
 pyrrhocentra donkelaeri, IV, 99
 pyrrhocentra, IV, 121
 quadrata, IV, 175
 quadrispina, IV, 88
 quadrispina major, IV, 88
 quehli, IV, 172
 radians, IV, 28, 34, 36, 37, 39, 42
 radians daemoneceras, IV, 36
 radians echinus, IV, 42
 radians globosa, IV, 36
 radians impexicoma, IV, 5, 36, 37
 radians sulcata, IV, 48, 49
 radicanissima, IV, 15, 16
 radiosa, IV, 43, 45
 radiosa alversonii, IV, 46, 47
 radiosa arizonica, IV, 45
 radiosa borealis, IV, 45
 radiosa chlorantha, IV, 43
 radiosa deserti, IV, 46
 radiosa neo-mexicana, IV, 45
 radiosa texana, IV, 45
 radlana, IV, 22
 radula, IV, 122
 ramossissima, IV, 51
 raphidacantha, IV, 50, 51
 raphidacantha ancistracantha, IV, 15

Mammillaria—continued,
 rebsamiana, IV, 82
 rebuti, IV, 172
 recta, IV, 108
 recurva, IV, 31, 77
 recurvata, IV, 27, 28
 recurvospina, IV, 27, 28, 47, 51, 123
 regia, IV, 174
 retusa, III, 81; IV, 38
 rhabdiantha, IV, 15, 16
 rhabdiantha humilior, IV, 15
 rhodacantha, IV, 133
 rhodacantha pallidior, IV, 133
 rhodantha, IV, 114, 117, 121, 122, 123, 176
 rhodantha andreae, IV, 121
 rhodantha aurea, IV, 123
 rhodantha aureiceps, IV, 114
 rhodantha callaena, IV, 122
 rhodantha celsii, IV, 123
 rhodantha centrispina, IV, 121
 rhodantha chrysacantha, IV, 122
 rhodantha crassispina, IV, 122
 rhodantha cristata, IV, 123
 rhodantha droegeana, IV, 10, 122
 rhodantha fulvispina, IV, 122, 123, 176
 rhodantha fuscata, IV, 122, 123
 rhodantha inuncta, IV, 123
 rhodantha major, IV, 121
 rhodantha neglecta, IV, 121, 123
 rhodantha odieriana, IV, 122
 rhodantha pfeifferi, IV, 122, 123
 rhodantha prolifera, IV, 121
 rhodantha pyramidalis, IV, 122
 rhodantha rubens, IV, 121
 rhodantha ruberrima, IV, 122
 rhodantha rubescens, IV, 122
 rhodantha rubra, IV, 122, 123
 rhodantha ruficeps, IV, 122, 123
 rhodantha schochiana, IV, 123
 rhodantha stenocephala, IV, 122
 rhodantha sulphurea, IV, 122
 rhodantha tentaculata, IV, 122
 rhodantha wendlandii, IV, 121
 rhodocentra, IV, 174
 rhodocentra gracilispina, IV, 174
 rigidispina, IV, 106, 107
 robusta, IV, 121
 robustispina, IV, 33, 34
 roematactina, IV, 171
 roessingii, IV, 172
 roii, IV, 172
 rosea, IV, 174
 roseana, IV, 22
 ruettii, IV, 115
 ruficeps, IV, 121
 rufidula, IV, 174
 rufocrocea, IV, 134, 174
 rungii, IV, 129
 ruschiana, IV, 174
 russea, IV, 122
 rutila, IV, 113, 169
 rutila octospina, IV, 169
 rutila pallidior, IV, 169
 salm-dyckiana, IV, 39, 40
 salm-dyckiana brunnea, IV, 39

Mammillaria—continued,
 salmiana, IV, 172
 saluciana, IV, 171
 sanguinea, IV, 118, 119
 sartorii, IV, 82
 sartorii brevispina, IV, 82
 sartorii longispina, IV, 82
 saxatilis, IV, 169
 scepontocentra, IV, 41
 schaeferi, IV, 112
 schaeferi longispina, IV, 112
 scheeri, IV, 17, 28, 29, 40, 51, 55
 scheeri valida, IV, 28, 29
 scheidweileriana, IV, 148
 schelhasei, IV, 149
 schelhasei lanuginosior, IV, 149
 schelhasei rosea, IV, 149
 schelhasei sericata, IV, 49
 schelhasei truncinata, IV, 149
 schiedeana, IV, 128
 schlechtendalii, IV, 30
 schlechtendalii laevior, IV, 30
 schmerwitzii, IV, 169
 schmidtii, IV, 78
 schochiana, IV, 123
 schumanniana, IV, 59
 schumannii, IV, 57, 58, 59
 scleracantha, IV, 103
 scolymoides, IV, 18, 39
 scolymoides longiseta, IV, 39
 scolymoides nigricans, IV, 39
 scolymoides raphidacantha, IV, 15
 seegeri, IV, 118
 seegeri gracilispina, IV, 118
 seegeri mirabilis, IV, 118
 seegeri pruinosa, IV, 118
 seemannii, IV, 170
 seideliana, IV, 144
 seidelii, IV, 174
 seitziana, IV, 83
 semilonia, IV, 172
 sempervivi, IV, 86, 87
 sempervivi lacteviridis, IV, 86
 sempervivi tetracantha, IV, 86
 senckiana, IV, 83
 senckei, IV, 83, 99
 senilis, IV, 19, 20
 senilis diguetii, IV, 20
 senilis haseloffii, IV, 20
 senilis linkei, IV, 20
 senkii, IV, 99
 sericata, IV, 128
 setispina, IV, 22, 23
 setosa, IV, 88
 severini, IV, 174
 similis, IV, 49, 52
 similis caespitosa, IV, 52
 similis robustior, IV, 52
 simonis, IV, 172
 simplex, IV, 53, 61, 70, 71
 simplex affinis, IV, 71
 simplex flavescens, IV, 166
 simplex parvimamma, IV, 70
 simpsonii, III, 90
 solitaria, IV, 175
 sororia, IV, 170
 spaethiana, III, 91
 speciosa, IV, 51, 174, 175
 spectabilis, IV, 105, 174
 sphacelata, IV, 138, 170
 sphaerica, IV, 61, 62, 158
 sphaerotracha, IV, 130
 sphaerotracha rosea, IV, 130
 spinaurea, IV, 170
 spinii, IV, 133
 spinosa, IV, 175
 spinosior, IV, 79
 spinosissima, III, 179; IV, 117, 118, 119, 164
 spinosissima auricoma, IV, 118, 119

- Mammillaria*—*continued*,
spinosissima aurea, IV, 118
spinosissima brunnea, IV, 118, 119
spinosissima eximia, IV, 118
spinosissima flavida, IV, 118
spinosissima haseloffii, IV, 118
spinosissima hepatica, IV, 118
spinosissima herrmannii, IV, 118
spinosissima isabellina, IV, 118
spinosissima linkeana, IV, 118
spinosissima mirabilis, IV, 118
spinosissima pruinosa, IV, 118
spinosissima pulcherrima, IV, 118
spinosissima rubens, IV, 118
spinosissima sanguinea, IV, 118
spinosissima seegeri, IV, 118
spinosissima vulpina, IV, 118
splendens, IV, 107
squarrosa, IV, 91
staurotypa, IV, 87
stella-aurata, IV, 134, 135
stella-aurata gracilispina, IV, 134
stella-aurata minima, IV, 135
stella de tacubaya, IV, 164
stellaris, IV, 124
stellata, IV, 124
stenocephala, IV, 121
stephani, IV, 172
stipitata, IV, 15
straminea, IV, 166
strobiliformis, IV, 17, 48, 54, 55
strobiliformis caespititia, IV, 54
strobiliformis durispina, IV, 54
strobiliformis pubescens, IV, 54
strobiliformis rufispina, IV, 54
stueberi, IV, 121, 123
suaveolens, IV, 170
subangularis, IV, 90, 91
subcirrhifera, IV, 91
subcrocea, IV, 134, 135
subcrocea anguinea, IV, 135
subcrocea intertexta, IV, 134
subcrocea rufescens, IV, 134
subcrocea rutila, IV, 135
subcurvata, IV, 77
subchinata, IV, 134
suberecta, IV, 172
subpolyedra, IV, 101, 105
subtetragona, IV, 88
subulata, IV, 175
subulifera, IV, 174
sulcata, III, 82; IV, 48, 49
sulcimamma, IV, 30
sulco-glandulifera, IV, 15
sulcolanata, IV, 23, 37, 38, 51
sulcolanata macracantha, IV, 38
sulphurea, IV, 116, 121
supertexta, IV, 107, 108
supertexta caespitosa, IV, 108
supertexta compacta, IV, 108
supertexta dichotoma, IV, 113
supertexta rufa, IV, 134
supertexta tetraantha, IV, 107, 108
tacubayensis, IV, 164
tecta, IV, 174
tellii, IV, 172
- Mammillaria*—*continued*,
tentaculata, IV, 121, 123, 169
tentaculata conothele, IV, 123
tentaculata fulvispina, IV, 123
tentaculata picta, IV, 122
tentaculata rubra, IV, 123
tentaculata ruficeps, IV, 121, 123
tenuis, IV, 134, 135
tenuis arrecta, IV, 134
tenuis coerulescens, IV, 134
tenuis derubescens, IV, 134
tenuis media, IV, 134
tenuis minima, IV, 134
tetracantha, IV, 79, 106, 107
tetracentra, IV, 77
tetrancistra, IV, 60
texana, IV, 125
texensis, IV, 76
thelocampitos, IV, 30
thornberi, IV, 162
toaldoae, IV, 98
tomentosa, IV, 175
tomentosa flava, IV, 173
tortolensis, IV, 72
triachantha, IV, 90
trichacantha, IV, 151
trigona, III, 81
trohartii, IV, 170
tuberculosa, IV, 53, 54, 55, 137
turbinata, III, 106, 107
uberiformis, IV, 63, 64
uberiformis gracilior, IV, 63
uberiformis hexacentra, IV, 62
uberiformis major, IV, 63
uberiformis variegata, IV, 63
uberimamma, IV, 78
uhdeana, IV, 118
umbrina, IV, 164, 166
uncinata, IV, 140
uncinata biuncinata, IV, 140
uncinata rhodacantha, IV, 140
uncinata spinosior, IV, 140
uniseta, IV, 170
urbaniana, IV, 48
utahensis, IV, 43
valida, IV, 28, 30, 34, 8, 8
varimamma, IV, 175
varimamma, IV, 172
venusta, IV, 58
verhaertiana, IV, 164
versicolor, IV, 77
vetula, IV, 130, 131
vetula major, IV, 131
villa-lerdo, IV, 172
villifera, IV, 89, 102
villifera aeruginosa, IV, 88
villifera carnea, IV, 88
villifera cirrhosa, IV, 88
villifera cirrosa, IV, 88
villosa, IV, 172
viperina, IV, 170, 171
virens, IV, 95
viridis, IV, 79, 96, 97
viridis praelii, IV, 96
viridula, IV, 166
vivipara, IV, 43, 44, 45, 48
vivipara neo-mexicana, IV, 48
vivipara radiosa, IV, 43, 45
vivipara radiosa neo-mexicana, IV, 45
vivipara vera, IV, 43, 44
voburnensis, IV, 100
vulpina, IV, 118
waltonii, IV, 98
webbiana, IV, 87
wegeneri, IV, 175
wegeneri cristata, IV, 172
wilcoxii, IV, 153, 154
wildiana, IV, 143
- Mammillaria*—*continued*,
wildiana compacta, IV, 143
wildiana cristata, IV, 143
wildiana major, IV, 143
wildiana monstrosa, IV, 143
wildiana rosea, IV, 148
wildiana spinosior, IV, 143
wildii, IV, 143
wildii compacta, IV, 143
wildii cristata, IV, 143
williamsii, III, 84
winkleri, IV, 41
wissmannii, IV, 52
woburnensis, IV, 100
wrightii, IV, 152, 153, 154
xanthispina, IV, 172
xanthotricha laevior, IV, 93
xanthotricha, IV, 92
zegschwartzii, IV, 175
zephyranthiflora, IV, 160
zephyranthoides, IV, 159, 160
zepnickii, IV, 175
zeyeriana, IV, 171
zooderi, IV, 79
zuccariniana, IV, 89, 90
zuccarinii, IV, 80
- Mammillaria*, IV, 23
Manca caballo, III, 175
Mandacaru, II, 9
Mandacaru de boi, II, 9
Mandacaru de penacho, II, 30
Martin's tail-cereus, II, 224
Matéare, I, 13
Matucana, III, 78, 102-104
haynei, III, 102, 103, 104
Mediocactus, II, 183, 210-212
coccineus, II, 211, 212
megalanthus, II, 211, 212, 213
Melocactus, II, 170; III, 91, 140, 166, 175, 186, 216, 220, 221, 226, 227, 229, 231, 234, 236, 237; IV, 41, 124, 281
aciculosus, III, 223
aciculosus adauctus, III, 223
acuatus, III, 188, 189
acutatus, III, 188
albispinus, III, 223
ambiguus, III, 72, 98
amoenus, III, 221, 232
angusticostatus, III, 222
appropinquatus, III, 223
approximatus, III, 222
arcuatus, III, 223, 224
argenteus, III, 222
argenteus tenuispinus, III, 222
armatus, III, 223
atrosanguineus, III, 230
atrovirens, III, 238
baarsianus, III, 223
bargei, III, 223, 224
besleri, III, 219, 220
besleri affinis, III, 142
bradleyi, III, 230
bradypus, II, 27
brongniartii, III, 225, 227
brongniartii, III, 225
buysianus, III, 223
caesius, III, 221, 233, 234
caesius griseus, III, 233
capillaris, III, 223
cephalenoplus, III, 221, 233
columna-trajani, II, 76
communiformis, III, 224, 225, 238
communis, III, 221, 224, 225, 226, 231, 232, 238
communis acicularis, III, 224
communis atrosanguineus, III, 230
communis bradleyi, III, 230
communis conicus, III, 226
communis croceus, III, 230
- Mammillaria*—*continued*,
communis eustachianus, III, 230
communis grengeli, III, 230
communis havannensis, III, 230
communis hookeri, III, 230
communis joerdensii, III, 232
communis laniferus, III, 224
communis macrocephalus, III, 226
communis magnisulcatus, III, 224
communis oblongus, III, 226
communis ovatus, III, 230, 231
communis spinosior, III, 224
communis viridis, III, 230
compactus, III, 223
contortus, III, 223
cordatus, III, 223
cornutus, III, 222, 224
coronatus, III, 238
crassicosatus, III, 232
crassispinus, III, 226
croceus, III, 230, 231
curvispinus, III, 221, 229, 237
cylindricus, III, 223
delessertianus, III, 221, 229, 237
depressus, III, 176, 235
dichroacanthus, III, 224, 230
dilatatus, III, 223
eburneus, III, 223
elegans, III, 72
elmeetti, IV, 289
elongatus, III, 223
ernesti, III, 227
euryacanthus, III, 223
eustachianus, III, 230, 231
evertszianus, III, 222
excavatus, III, 237
exsertus, III, 223, 224
extensus, III, 223
ferox, III, 178, 221, 222, 238
ferus, III, 223
firmus, III, 223
flammerus, III, 223
flavispinus, III, 223
flexilis, III, 223
flexus, III, 222
fluminensis, III, 238
gardenerianus, III, 235
gilliesii, III, 75
gilvispinus, III, 223
gilvispinus planispinus, III, 223
gladiatus, III, 119, 120
goniacanthus, III, 235
goniodacanthus, III, 235
gracilis, III, 223
grandis, III, 223
grandispinus, III, 223
grengeli, III, 230
griseus, III, 221, 233
grollianus, III, 223, 224
guatemalensis, III, 227
harlowii, III, 232
havannensis, III, 221, 230, 232
hexacanthus, III, 222
hispanicolicus, III, 226
hookeri, III, 230
hookerianus, III, 238
humilis, III, 233, 234
hystrix, III, 237
incurvus, III, 222
inflatus, III, 223
ingens, III, 169
intermedius, III, 222
intermedius laticostatus, III, 223
intermedius rotundatus, III, 223
intermedius tenuispinus, III, 223

Melocactus—*continued*,
 intortus, III, 230, 231
 intricatus, III, 206
 inversus, III, 223
 koolijckianus, III, 222
 koolijckianus adustus, III, 222
 laciniatus, III, 182
 lamarkii, III, 224
 langsdorfii, III, 199
 latispinus, III, 143
 lehmanni, III, 221, 222, 224
 lemarii, III, 226
 leopoldii, III, 238
 leucacanthus, III, 223
 leucaster, III, 238
 limis, III, 222
 linkii, III, 230, 231
 linkii trispinus, III, 231
 lobelii, III, 238
 lutescens, III, 223
 macracanthoides, III, 230, 231
 macracanthus, III, 222, 224, 226
 macrocanthus, III, 230
 mamillariaeformis, III, 221; IV, 41
 mamillariiformis IV, 41
 martialis, III, 223
 maxonii, III, 227, 228
 melocactoides, III, 235
 melocactus, III, 224
 meonacanthus, III, 224, 225
 microcarpus, III, 223
 microcephalus, III, 222, 224
 microcephalus olivascens, III, 223
 miqueli, III, 230, 231
 monoclonos, III, 41
 monvilleanus, III, 237
 nanus, III, 222
 negyi, III, 236
 neryi, III, 236
 nigro-tomentosus, III, 238
 obliquus, III, 222
 obovatus, III, 223
 obtusipetalus, III, 221, 232
 obtusipetalus crassicosatus, III, 232
 octogonus, III, 238
 oreas, III, 227
 orthacanthus, III, 138
 ovatus, III, 223
 pachycentrus, IV, 289
 parthoni, III, 235
 parvispinus, III, 222, 224, 237
 patens, III, 222, 224
 pentacanthus, III, 223
 pentacentrus, III, 235
 peruvianus, III, 234
 pinguis, III, 223
 pinguis areolatus, III, 223
 pinguis laticostatus, III, 223
 pinguis planispinus, III, 223
 pinguis tenuissimus, III, 223
 placentifformis, III, 219
 platyacanthus, III, 172
 poliacanthus, III, 198, 199
 portoricensis, III, 230, 231
 pulvinosus, III, 223
 pusillus, III, 222, 224, 225
 pyramidalis, III, 222, 224
 pyramidalis carneus, III, 222
 pyramidalis compressus, III, 223
 pyramidalis costis-angustroebus, III, 223
 pyramidalis pumilus, III, 223
 radiatus, III, 223
 radiatus contortus, III, 223
 retiusculus, III, 222
 retiusculus angusticostatus, III, 222

Melocactus—*continued*,
 recurvus, III, 141, 143
 repens, III, 238
 reticulatus, III, 223
 reversus, III, 222
 roseus, III, 222
 rotatus, III, 223
 rotifer, III, 223
 rotifer angustior, III, 223
 rotula, III, 224
 rotula angusticostatus, III, 223
 rotula validispinus, III, 223
 rubellus, III, 222
 rubellus ferox, III, 222
 rubellus hexacanthus, III, 222
 rubens, III, 221, 224, 225
 rudis, III, 222
 ruestii, III, 227
 rufispinus, III, 225
 salmianus, III, 222, 224
 salmianus aciculosus, III, 223
 salmianus adauctus, III, 223
 salmianus contractus, III, 223
 salmianus quadrispinus, III, 224
 salmianus spectabilis, III, 223
 salmianus trispinus, III, 224
 salvador, III, 221, 228, 229
 salvatoris, III, 229
 san salvador, III, 228
 schlumbergerianus, III, 230
 sellowii, III, 188, 189
 sordidus, III, 223
 spatanginus, III, 222, 224
 spatangus, III, 222
 spina christi, III, 178, 238
 stellatus, III, 223
 stellatus dilatatus, III, 223
 stellatus flavispinus, III, 223
 stellatus inflatus, III, 223
 stellatus sordidus, III, 223
 stramineus, III, 222
 stramineus trichacanthus, III, 222
 tenuispinus, III, 196
 tenuissimus, III, 223
 tephracanthus, III, 188, 189
 trachycephalus, III, 223
 trichacanthus, III, 222
 trigonaster, III, 223
 trigonus, III, 223
 tuberculatus, III, 170
 uncinatus, III, 223, 224
 violaceus, III, 235
 viridescens, III, 140
 wendlandii, III, 230
 xanthacanthus, III, 230
 zuccarinii, III, 222, 224
Melocardus echinatus, III, 225
 Melon cactus, III, 228
 Melon thistle, IV, 71
 Melones, III, 221
 Mescal button, III, 84; IV, 286
 Mesembryanthemoides, IV, 220
 Mesembryanthemum, IV, 222
 Microgoni, III, 202
 Mila, III, 77, 78, 211, 212
 caespitosa, III, 211
 Miqueliana, I, 44, 78
 Mission cactus, I, 186
 Monster cactus, III, 170, 171
 Monvillea, II, 1, 21-25
 amazonica, II, 21, 24
 cavendishii, II, 21, 22, 118, 224; IV, 268
 diffusa, II, 21, 24
 insularis, II, 21, 22, 23
 maritima, II, 21, 24, 25
 phatnosperma, II, 21, 24
 spegazzinii, II, 21, 23; IV, 268

Muyusa, II, 160
 Myrtillocactus, II, 1, 178-181
 cochal, TI, 179, 180; IV, 280
 eichlamii, II, 179, 180, 181
 geometrizans, II, 179, 181; IV, 280
 schenckii, II, 179, 180
 Neoabbottia, IV, 280-282
 paniculata, IV, 247, 280, 281, 282
 Najá de Culebra, I, 19
 Neobesseyia, IV, 3, 51-53
 missouriensis, IV, 51, 53
 notesteinii, IV, 51, 53
 similis, IV, 51, 52
 wissmannii, IV, 51, 52
 Neolloydia, IV, 3, 14-18
 beguinii, IV, 14, 16
 ceratites, IV, 14, 16, 17
 clavata, IV, 14, 15
 conoidea, IV, 14, 15, 17, 131
 horripila, IV, 14, 16
 pilispina, IV, 14
 texensis, IV, 14, 18
 Neomammillaria, IV, 3, 65-176, 250
 albicans, IV, 68, 138, 139
 amoena, IV, 68, 120, 121
 applanata, IV, 66, 76
 arida, IV, 66, 73, 74
 armillata, IV, 69, 157
 aureiceps, IV, 67, 113, 114
 barbata, IV, 69, 144, 145
 bocasana, IV, 69, 147
 boedeckeriana, IV, 69, 154
 bombycina, IV, 69, 160, 161
 brandegeei, IV, 66, 73, 74
 camptotricha, IV, 68, 126, 127
 candida, IV, 68, 130
 carnea, IV, 66, 88, 89
 carretii, IV, 69, 157, 160
 celsiana, IV, 67, 112
 cerralboa, IV, 68, 116
 chinocephala, IV, 67, 100, 101
 collina, IV, 67, 111
 collinsii, IV, 67, 99, 101, 104
 compressa, IV, 67, 90, 91, 92
 confusa, IV, 67, 102
 konzattii, IV, 67, 103, 105
 crocidata, IV, 66, 87
 dealbata, IV, 67, 102, 110
 decipiens, IV, 68, 131
 densispina, IV, 68, 119
 denudata, IV, 68, 129
 dioica, IV, 69, 158, 159
 discolor, IV, 68, 131, 132
 donatii, IV, 67, 111
 echinaria, IV, 68, 136
 eichlamii, IV, 67, 94, 95
 elegans, IV, 67, 107, 108, 165
 elongata, IV, 68, 133, 134
 eriacantha, IV, 68, 127, 128
 evermannia, IV, 67, 97
 fasciculata, IV, 69, 162
 fertilis, IV, 68, 131
 flavovirens, IV, 66, 85
 formosa, IV, 66, 90
 fragilis, IV, 68, 133
 fraileana, IV, 69, 157
 galeottii, IV, 67, 105
 gaumeri, IV, 66, 72
 geminispina, IV, 67, 98, 99
 gigantea, IV, 66, 85
 glochidata, IV, 69, 149, 150
 goodridgei, IV, 69, 158
 graessneriana, IV, 68, 116, 117
 gummifera, IV, 66, 74
 haageana, IV, 67, 110
 hamata, IV, 68, 140, 141
 hemisphaerica, IV, 66, 73, 75, 76
 heyderi, IV, 66, 73, 75, 76
 hirsuta, IV, 69, 146
 jaliscana, IV, 69, 160

Neomammillaria—*continued*,
 johnstonii, IV, 66, 80, 81
 karwinskiana, IV, 67, 95
 kewensis, IV, 67, 104, 106
 kunzeana, IV, 69, 145, 147, 152
 lanata, IV, 67, 104, 105
 lasiacantha, IV, 68, 128, 129
 lenta, IV, 68, 128, 129
 lloydii, IV, 66, 89
 longicoma, IV, 69, 146, 147, 150
 longiflora, IV, 69, 162, 163
 mammillaris, IV, 66, 70, 74, 75
 maddockii, IV, 66, 70, 74, 75
 macracantha, IV, 66, 78, 79, 80
 magnimamma, IV, 66, 77, 78, 94
 mainae, IV, 69, 153, 154
 mammillaris, IV, 66, 70, 71, 166
 mazatlanensis, IV, 68, 137, 138
 meiacantha, IV, 66, 84, 85
 melanocentra, IV, 66, 81, 82
 mercadensis, IV, 69, 145, 146
 microcarpa, IV, 69, 152, 155, 157
 milleri, IV, 69, 156, 176
 multiceps, IV, 68, 125, 126
 multifloris, IV, 69, 148
 multihamata, IV, 69, 146, 152
 mundtii, IV, 67, 110, 112
 mystax, IV, 67, 92, 93, 95
 napina, IV, 67, 104
 nelsonii, IV, 69, 163
 nivosa, IV, 66, 71, 72
 nunezii, IV, 68, 119, 120
 obscura, IV, 66, 87
 occidentalis, IV, 69, 160, 161
 oliviae, IV, 68, 135
 ortegae, IV, 66, 83, 84, 250
 painteri, IV, 69, 151, 152
 palmeri, IV, 68, 140
 parkinsonii, IV, 67, 96, 98
 peninsularis, IV, 66, 85
 perbella, IV, 67, 111
 petrophila, IV, 66, 73
 petterssonii, IV, 67, 93, 94
 phaeacantha, IV, 68, 116
 phymatothele, IV, 66, 75, 76, 77
 plumosa, IV, 68, 121, 123
 polyedra, IV, 67, 102, 103
 polygona, IV, 67, 101
 polythele, IV, 66, 86, 88
 pottii, IV, 68, 136, 137, 250
 praelii, IV, 67, 96
 pringlei, IV, 68, 115
 prolifera, IV, 68, 124, 125, 166
 pseudoperbella, IV, 67, 109
 pygmaea, IV, 69, 142
 pyrrhocephala, IV, 67, 99, 103
 rekoii, IV, 68, 136, 141, 142
 rhodantha, IV, 68, 120, 121
 ruestii, IV, 67, 114, 115
 runyonii, IV, 66, 81
 safordii, IV, 69, 149, 151, 160
 sartorii, IV, 66, 82, 83
 schiedweiliana, IV, 69, 148
 schiedeana, IV, 68, 127, 128
 schelhaesi, IV, 69, 149
 scrippsiana, IV, 66, 84, 85
 seideliana, IV, 69, 144
 seitziana, IV, 66, 82, 83
 sempervivi, IV, 66, 86
 sheldoni, IV, 69, 156, 157
 slevinii, IV, 68, 139
 solisii, IV, 68, 142, 143
 sphacelata, IV, 68, 138
 spinosissima, IV, 68, 117
 standleyi, IV, 67, 96, 97

- Neomammillaria*—*continued*,
 subpolyedra, IV, 67, 105, 106
 swinglei, IV, 69, 158
 tacubayensis, IV, 69, 164
 tenampensis, IV, 67, 101, 104
 tetracantha, IV, 67, 106, 108
 trichacantha, IV, 69, 151
 umbrina, IV, 69, 164
 uncinata, IV, 68, 140
 vetula, IV, 68, 130, 131
 verhaertiana, IV, 69, 164
 villifera, IV, 67, 102
 viridiflora, IV, 69, 153
 wilcoxei, IV, 69, 153
 wildii, IV, 69, 141, 143
 woburnensis, IV, 67, 96, 100
 wrightii, IV, 69, 152, 153
 xanthina, IV, 69, 164
 yucatanensis, IV, 67, 114
 zephyranthoides, IV, 69, 159, 164
 zuccariniana, IV, 66, 89, 90
Neoporteria, III, 59, 77, 94-100; IV, 122
 chilensis, III, 94, 97, 99
 fusca, III, 94, 96, 99
 jussieui, III, 94, 96, 97
 nidus, III, 94, 95
 nigricans, III, 94, 95, 96
 occulta, III, 94, 95
 subgibbosa, III, 94, 97, 203
Neoraimondia, II, 3, 181-183; IV, 281
 macrostibas, II, 181, 182
 Night-blooming cereus, II, 113, 187, 188
Noble leuchtenbergia, III, 108
Nocturna, IV, 186
Nopal, I, 34
Nopal cardón, I, 184
Nopal de Castilla, I, 224
 silvestre, IV, 263
Nopalea, I, 8, 24, 33-39, 43, 155, 216; IV, 253
 angustifrons, IV, 253
 auberi, I, 34, 37, 38, 39; IV, 253
 coccifera, IV, 253
 cochenillifera, I, 34, 181; IV, 253
 dejecta, I, 34, 36, 37; IV, 253
 gaumeri, I, 34, 37, 216
 guatemalensis, I, 33, 34, 35
 inaperta, I, 33, 34, 37, 38
 karwinskiana, I, 34, 37, 38, 39
 lutea, I, 33, 34, 35
 moniliformis, I, 33, 206
Nopalea, I, 26
Nopalnochetzli, I, 5
Nopalxochia, IV, 177, 204-205
 phyllanthoides, IV, 205
Notocactus, III, 94, 187
Nyctocereus, II, 1, 2, 108, 117-120, 121
 guatemalensis, II, 117, 118, 119, 120; IV, 276
 hirschtianus, II, 118, 119
 neumannii, II, 118, 119
 oaxacensis, II, 118, 120
 serpentinus, II, 20, 118; IV, 276
Obtusae, II, 4
Ohulago, I, 103
Olago, I, 103
Old man cactus, II, 27
Oleosi, III, 23
Oligogoni, II, 3, 77
Olneya tesota, IV, 275
Ophiorhizalis, IV, 208
Opuntia, I, 8, 14, 24, 25, 30, 32, 33, 34, 38, 39, 40, 42-215, 217-225; II, 3, 92, 118, 156, 164; IV, 42, 186, 209, 214, 217, 220, 249, 250, 252, 253, 260, 264, 266
 abjecta, IV, 257
Opuntia—*continued*,
 acanthocarpa, I, 56, 57; IV, 254
 aciculata, I, 160, 165
 acracantha, I, 91
 aequatorialis, I, 110, 116, 219, 220
 affinis, I, 169, 170
 airampo, I, 161
 albicans, I, 191; IV, 264
 albicans laevior, I, 191
 albiflora, I, 73, 74
 albisetosa, I, 134
 alcahes, I, 58, 67, 69, 70
 alexanderi, IV, 256
 alifayucca, I, 185
 alfayucca, I, 185
 allairei, I, 126
 alpicola, I, 214
 alpina, I, 33
 alta, I, 165, 166
 amarilla, IV, 264
 americana, I, 214
 ammophila, I, 211; IV, 265
 amyclaea, I, 112, 177, 181, 185
 amyclaea ficus-indica, I, 177
 anacantha, I, 107, 109, 110
 anahuacensis, I, 169
 andicola, I, 89, 90 IV, 255
 andicola elongata, I, 89
 andicola fulvispina, I, 89
 andicola major, I, 89
 andicola minor, I, 90
 angusta, I, 101
 angustata, I, 124, 129, 140, 142, 149
 angustata comonduensis, I, 124
 antillana, I, 110, 115, 163; IV, 259
 aoracantha, I, 90, 91; IV, 255
 aquosa, I, 29
 arborea, I, 209
 arborescens, I, 43, 63, 64, 65; IV, 254
 arborescens spinosior, IV, 254
 arborescens versicolor, I, 62
 arbuscula, I, 47, 50, 51
 arechavaletai, I, 156, 158; IV, 262
 arenaria, I, 193, 194, 195; IV, 264
 argentina, I, 209, 211, 212
 arizonica, I, 147, 148
 arkansana, I, 128
 articulata, I, 89
 assumptionis, I, 159
 atacamensis, I, 90, 94
 atrispina, I, 140, 142
 atropes, I, 169, 170
 attulica, I, 214
 auberi, I, 37; IV, 253
 aulacothecae, I, 95
 aurantiaca, I, 101, 107
 aurantiaca extensa, I, 107
 australis, I, 87, 88
 austrina, I, 45, 126, 130; IV, 265
 ayrampo, IV, 261
 azurea, I, 140, 143
 bahamama, I, 202, 203, 204
 bahaiensis, I, 209, 210, 211
 balearica, I, 161
 ballii, I, 137
 barbata, I, 214
 barbata gracillima, I, 214
 bartramii, I, 181; IV, 265
 basilaris, I, 118, 119, 120, 136; IV, 259
 basilaris albiflora, I, 120; IV, 259
 basilaris coerulea, I, 120
 basilaris cordata, I, 120; IV, 259
Opuntia—*continued*,
 basilaris cristata, I, 120
 basilaris minima, IV, 259
 basilaris nana, I, 120
 basilaris nevadensis, I, 120
 basilaris pfersdorffii, I, 120
 basilaris ramosa, I, 119, 120
 basilaris treleasei, I, 119
 beckeriana, I, 168; IV, 263
 bella, I, 110, 111, 112
 bentonii, I, 161, 163
 bergeriana, I, 149, 152, 169; IV, 262
 bernardina, I, 57, 81
 bernardina cristata, I, 57
 bernhardinii, I, 214
 betancourt, I, 214
 bicolor, I, 214; II, 106
 bigelovii, I, 58, 59; IV, 254
 blakeana, I, 144, 145
 boldinghii, I, 149, 155; IV, 262
 boliviana, I, 71, 97, 98; IV, 256
 bonaerensis, I, 156, 158; IV, 262
 bonplandii, I, 160, 168, 223, 224
 borinquensis, I, 102, 103, 104
 brachyarthra, I, 193, 194; IV, 264
 brachyclada, I, 120
 brachydelphis, I, 42; IV, 253
 bradtiana, I, 215
 brandegeei, I, 25, 28
 brasiliensis, I, 209, 210, 211; IV, 265
 brasiliensis gracilior, IV, 265
 brasiliensis minor, I, 210; IV, 265
 brasiliensis schomburgkii, I, 210
 brasiliensis spinosior, I, 210
 brasiliensis tenuifolia, I, 210
 brasiliensis tenuior, I, 210
 brunnescens, I, 149, 150
 bulbispina, I, 79, 83; IV, 255
 bulbosa, I, 131
 burrageana, I, 67, 70
 cacanapa, I, 165, 166; IV, 263
 caeruleascens, I, 51
 caesia, I, 144
 caespitosa, I, 127
 calacantha, I, 214
 calacantha rubra, I, 214
 calantha, I, 136
 californica, I, 58; IV, 253
 calmalliana, I, 60, 61
 calva, I, 89
 camanchica, I, 144, 145; IV, 262
 camanchica albispina, I, 144
 camanchica luteo-staminea, I, 144
 camanchica orbicularis, I, 144
 camanchica rubra, I, 144
 camanchica salmonea, I, 144
 campestris, I, 90, 99
 camuessa, I, 191
 canada, I, 160, 167, 168
 candelabriformis, I, 182
 candelabriformis rigidior, I, 182
 canina, I, 107, 108
 cantabrigiensis, I, 121, 160, 167; IV, 263
 canterai, I, 159
 caracassana, I, 110, 116, 219
 cardenche, I, 64
 cardiosperma, I, 156, 157, 158
 cardona, I, 184
 caribaea, I, 47, 48, 49; IV, 253
 carolina, I, 214
 carrizalensis, I, 79
Opuntia—*continued*,
 castillae, I, 185, 186
 catacantha, I, 43, 208; IV, 264
 cathocantha, IV, 264
 cereiformis, I, 79, 215
 cervicornis, I, 194
 chaetocarpa, I, 182
 chaffeyi, I, 30, 212, 213
 chakensis, I, 158; IV, 262
 chapistle, I, 27
 chata, IV, 264
 chavena, I, 184
 chella, IV, 254
 chihuahuensis, I, 144, 145
 chlorotica, I, 142, 159, 160, 161; IV, 262
 chlorotica santa-rita, I, 142
 cholla, I, 60, 61, 62; IV, 254
 chrysacantha, I, 167
 ciliosa, I, 214
 ciribe, I, 8, 59, 60
 claviroides, I, 72, 73; IV, 254
 claviroides cristata, I, 7; IV, 254
 claviroides fasciata, I, 73
 claviroides fastigiata, I, 73
 claviroides monstrosa, I, 73
 clavata, I, 79, 81, 215; II, 106, 107; IV, 255
 clavellina, I, 52, 54; IV, 253
 coccifera, I, 35
 coccinea, I, 114
 cochenillifera, I, 33, 34, 35
 cochinelifera, I, 34
 cochineria, I, 192
 coerulea, I, 134
 coindetti, I, 184
 columbiana, I, 193
 comonduensis, I, 118, 124
 compressa, IV, 259
 confusa, I, 147
 congesta, I, 50
 consoleana, I, 214
 consolei, I, 214
 convexa, I, 165
 cordobensis, I, 188, 189
 cornigata, I, 95
 corotilla, I, 96
 corrugata, I, 90, 95; III, 179; IV, 256
 corrugata monvillei, I, 95
 costigera, I, 65
 covillei, I, 140, 145, 146
 crassa, I, 178, 179; IV, 262
 crassa major, I, 178
 cretochaeta, I, 183
 crinifera, I, 159, 176; IV, 263
 crinifera lanigera, I, 176
 cristata, I, 64
 cristata tenuior, I, 64
 cruciata, I, 207
 crystalenia, I, 193
 cubensis, I, 163
 cucumiformis, I, 97
 cuija, I, 149, 167
 cumingii, I, 77
 curassavica, I, 102, 103, 112; IV, 257
 curassavica elongata, I, 102
 curassavica longa, I, 102
 curassavica major, I, 102
 curassavica media, I, 102
 curassavica minima, I, 102
 curassavica minor, I, 102
 curassavica taylori, I, 103
 curvospina, I, 160
 cyanea, IV, 264
 cyanella, I, 165, 166
 cyclodes, I, 147, 148
 cycloidea, I, 128
 cylindrica, I, 71, 75, 77; IV, 254
 cylindrica cristata, I, 78; IV, 254

Opuntia—continued,
 cylindrica cristata minor, I, 78
 cylindrica monstrosa, I, 78
 cylindrica robustior, I, 78
 cymochila, I, 131, 132
 cymochila montana, IV, 260
 dactylifera, II, 97, 98
 darrahiana, II, 102, 106
 darwinii, I, 88, 90, 93, 94, 97; IV, 256
 davisii, I, 52, 54, 55
 deamii, I, 181, 187
 decipiens, I, 63, 65; IV, 254
 decipiens major, I, 64
 decipiens minor, I, 65
 decumana, I, 157, 180, 181, 186
 decumbens, I, 111, 116, 117, 121, 218; IV, 259
 decumbens irrorata, I, 117
 decumbens longispina, I, 117
 deflexa, I, 157
 dejecta, I, 37
 delaetiana, I, 149, 152
 delicata, I, 126, 132, 133
 deltica, I, 165
 demissa, I, 146, 147
 demorenia, I, 214
 demoriana, I, 214
 depauperata, I, 100, 101, 216, 217, 219
 deppel, I, 214
 depressa, I, 111, 117, 118
 deserta, I, 57, 58
 diademata, I, 89, 90; IV, 255
 diademata calva, I, 89
 diademata inermis, I, 89
 diademata oligacantha, I, 89
 diademata polyacantha, I, 89
 dichotoma, I, 214
 diffusa, I, 37
 digitalis, I, 72
 diguetii, I, 26, 29
 dillei, I, 147, 148
 dillenii, I, 106, 114, 115, 116, 159, 162, 163, 181, 222, 223; IV, 262, 263
 dillenii minor, I, 163
 dillenii orbiculata, I, 163
 dimorpha, I, 96
 diplacantha, I, 184
 discata, I, 140, 149; IV, 262
 discolor, I, 107, 109, 218, 219
 distans, I, 149, 155
 diversispina, IV, 262
 dobbieana, I, 181, 187, 224, 225
 dolabriformis, I, 207
 domingensis, IV, 259
 dorffii, IV, 259
 drummondii, I, 102, 104, 106; IV, 257
 dulcis, I, 165, 166
 durangensis, I, 169
 eborina, I, 214
 eburnea, I, 95
 eburnispina, IV, 260
 echinocarpa, I, 56, 57, 81; IV, 254
 echinocarpa major, I, 57
 echinocarpa nuda, I, 57, 58
 echinocarpa parkeri, I, 57, 58
 echinocarpa robustior, I, 57
 effulgia, IV, 264
 eichlamii, I, 181, 187, 188
 elata, I, 110, 156, 157
 elata delaetiana, I, 152
 elatior, I, 149, 153, 219; IV, 262
 elatior deflexa, I, 157
 ellemeetiana, I, 75
 ellisiana, I, 165; IV, 263
 elongata, I, 179, 181
 elongata laevior, I, 180

Opuntia—continued,
 emoryi, I, 80; IV, 255
 engelmannii, I, 140, 147, 148, 149, 164, 165, 167; IV, 262
 engelmannii cristata, IV, 262
 engelmannii cuija, I, 167
 engelmannii cyclodes, I, 147, 148
 engelmannii discata, IV, 262
 engelmannii dulcis, I, 165
 engelmannii littoralis, I, 165
 engelmannii monstrosa, I, 148
 engelmannii occidentalis, I, 146
 eocarpa, I, 144
 erecta, I, 214
 erinacea, I, 193, 195, 196; IV, 264
 erythrocentron, I, 174
 exaltata, I, 71, 75, 76, 77
 expansa, I, 147, 148
 extensa, I, 107
 exuviata, I, 63, 65; IV, 254
 exuviata angustior, I, 63
 exuviata major, I, 64
 exuviata spinosior, I, 63
 exuviata stellata, I, 63
 exuviata viridior, I, 63
 ferox, I, 206, 207
 ferruginispina, I, 165
 festiva, I, 214
 ficus-barbica, I, 177
 ficus-indica, I, 43, 156, 157, 177, 178, 181, 185, 189, 191; IV, 263
 ficus-indica albispina, I, 214
 ficus-indica amyclaea, I, 185
 ficus-indica decumana, I, 180
 ficus-indica gymnocarpa, I, 180
 filipendula, I, 137, 138; IV, 262
 flavicans, I, 191
 flavispina, I, 214
 flexibilis, I, 114
 flexospina, I, 165
 floccosa, I, 71, 86, 87; III, 102
 floccosa denudata, I, 86, 87
 floribunda, I, 74
 foliosa, I, 105, 106
 folio minori, I, 129
 formidabilis, I, 91, 92
 fragilis, I, 193, 194; IV, 264
 fragilis brachyarthra, I, 193; IV, 264
 fragilis caespitosa, I, 193
 fragilis tuberiformis, I, 193
 fragilis frutescens, I, 47
 frustulenta, I, 104, 105
 frutescens, I, 47, 48
 frutescens brevispina, I, 47, 48
 frutescens longispina, I, 47, 48
 fulgida, I, 67; IV, 254
 fulgida mamillata, I, 67
 fuliginosa, I, 149, 155
 fulvispina, I, 174
 fulvispina badia, I, 175
 fulvispina laevior, I, 175
 furiosa, I, 66
 fuscoatra, I, 126, 133
 fuscaulis, I, 191, 192
 fusiformis, I, 130, 131
 galapageia, I, 45, 149, 150, 151, 152
 galeottii, I, 6
 geissei, I, 78
 gilliesii, I, 91
 gilva, I, 163
 gilvescens, I, 149
 gilvoalba, I, 165, 166
 glaberrima, I, 178
 glauca, I, 178
 glaucescens, I, 200, 201
 glaucophylla, I, 214

Opuntia—continued,
 glaucophylla laevior, I, 214
 glomerata, I, 87, 89, 90, 94, 96; II, 138; IV, 255
 glomerata albispina, I, 90
 glomerata flavispina, I, 90
 glomerata minor, I, 90
 golziana, I, 26
 gomei, I, 165, 166
 gorda, I, 191, 192
 gosseliniana, I, 140, 141, 151
 gracilior, IV, 262
 gracilis, I, 47
 gracilis subpatens, I, 48
 grahamii, I, 7, 83, 84
 grandiflora, I, 126, 127, 129
 grandis, I, 200, 201; IV, 264
 grata, I, 92, 94, 95
 grata leoniana, IV, 256
 greggii, IV, 255
 greenii, I, 131
 greggiana, I, 147, 148
 griffithsiana, I, 165
 grosseiana, I, 107, 110
 guanicana, I, 208
 guatemalensis, I, 110, 113, 218, 219
 guerrana, I, 191, 192
 guilanchi, I, 173, 174
 gymnocarpa, I, 180
 haematocarpa, I, 166
 haenquiana, IV, 261
 haitiensis, I, 206
 hanburyana, I, 149, 153, 154
 hattoniana, I, 103
 helleri, I, 150, 152
 hempeliana, I, 86, 87; III, 101
 hernandezii, I, 181; IV, 263
 heteromorpha, I, 79
 hevernickii, I, 214
 hickenii, I, 90, 92, 93; IV, 255
 hieronymi, I, 211
 hispanica, IV, 264
 hitchenii, I, 214
 hochderfferi, I, 198
 horizontalis, I, 37, 90
 horrida, I, 162; IV, 263
 humifusa, I, 127, 128, 132; IV, 259
 humifusa cymochila, I, 128
 humifusa greenii, I, 128
 humifusa macrorrhiza, I, 128
 humifusa microsperma, I, 127
 humifusa oplocarpa, I, 128
 humifusa parva, I, 127
 humifusa stenochila, I, 128
 humifusa vaseyi, I, 146
 humilis, I, 113, 114; IV, 259
 humistrata, I, 120
 hypsophila, I, 71, 72
 hyptiacantha, I, 176, 181, 183, 184
 hystricina, I, 193, 197
 hystrix, I, 65, 66
 icterica, I, 173
 ignescens, I, 90, 98
 ignota, I, 90, 99
 imbricata, I, 48, 55, 60, 63, 64, 65, 66; IV, 254
 imbricata crassior, I, 63
 imbricata ramosior, I, 64
 imbricata tenuior, I, 64
 impedita, IV, 257
 inaequalis, I, 128
 inaequilateralis, I, 181, 187, 188, 189
 inamoena, I, 121, 125
 incarnadilla, I, 185, 186
 inermis, I, 161, 162, 163
 insularis, I, 150, 152
 intermedia, I, 127, 128
 intermedia prostrata, I, 128
 intricata, I, 119
 invicta, I, 79

Opuntia—continued,
 involuta, I, 87
 irrorata, I, 116, 117
 italica, I, 214; IV, 266
 ithypetala, I, 190
 jamaicensis, I, 110, 113
 joconostle, I, 214
 juniperina, I, 193, 197
 jussieuui, I, 214
 karwinskiana, I, 38
 keyensis, I, 159, 162, 163, 222, 223
 kiska-loro, I, 107, 108
 kleiniae, I, 47, 51, 64, 65
 kleiniae cristata, I, 51
 kleiniae laetevirens, I, 51
 kunzei, I, 80; IV, 255
 labouretiana, I, 180, 189; IV, 263
 labouretiana macrocarpa, I, 181
 laevis, I, 159, 161, 168; IV, 262
 lagopus, I, 86, 87, 88
 lanceolata, I, 177, 179, 203
 lanigera, I, 176; IV, 263
 larreyi, I, 191
 lasiacantha, I, 181, 182, 183, 184
 lata, I, 126, 220
 laxiflora, I, 165
 ledienii, I, 153
 lemaireana, I, 156
 leonina, I, 96; IV, 256
 leptarthra, I, 101
 leptocarpa, I, 166, 167
 leptocaulis, I, 47, 48, 49, 73, 213; III, 183; IV, 253
 leptocaulis brevispina, I, 47
 leptocaulis laetevirens, I, 47
 leptocaulis longispina, I, 47
 leptocaulis major, I, 48
 leptocaulis stipata, I, 47; IV, 253
 leptocaulis vaginata, I, 47
 leucacantha, I, 174, 175; IV, 253, 263
 leucacantha laevior, I, 175
 leucacantha subferox, I, 175
 leucacantha, I, 175
 leucophaea, I, 96
 leucostata, I, 214
 leucosticta, I, 174
 leucotricha, I, 151, 174, 175; IV, 263
 leucotricha fulvispina, I, 174
 ligostica, I, 128
 lindheimeri, I, 44, 148, 160, 164, 165, 166; IV, 263
 lindheimeri cyclodes, I, 147
 lindheimeri dulcis, I, 165
 lindheimeri littoralis, I, 165
 lindheimeri occidentalis, I, 146
 linguiformis, I, 160, 163
 littoralis, I, 147, 160, 164, 165
 lloydii, I, 60, 63
 longiclada, I, 161
 longiglochria, I, 214
 longispina, I, 95
 lubrica, I, 118, 119
 lucayana, I, 106, 163
 lucens, I, 149
 lucida, I, 214
 lurida, I, 173; IV, 263
 macateei, I, 126, 133, 220, 221
 macbridei, IV, 261
 maddockiana, I, 169, 170, 171
 mackensenii, I, 137, 138, 139
 macracantha, I, 187, 202, 203, 204, 205
 macrarthra, I, 126, 129
 macrocalyx, I, 118, 122
 macrocentra, I, 140, 141, 176; IV, 265

Opuntia—*continued*,
macrophylla, I, 214
macrorhiza, I, 126, 127, 128, 130, 131, 139, 166; IV, 260
maculacantha, I, 134; IV, 260
maelenii, I, 134
magenta, I, 142, 146
magna, I, 64
magnarenensis, I, 147
magnifolia, I, 34
maidenii, IV, 259
maiuen, I, 41
maldonadensis, I, 75
mamillata, I, 67, 68; IV, 254
maritima, I, 162; IV, 262
maxillare, I, 77
maxima, I, 177, 180, 181; IV, 263
media, I, 199, 200
mediterranea, I, 128
megacantha, I, 178, 181, 182, 185, 186; IV, 264
megacantha lasiacantha, I, 182
megacantha tenuispina, I, 184
megacantha trichacantha, I, 186
megacarpa, I, 145
megalantha, I, 169
megalarthra, I, 192
megarhiza, I, 137
mendocienses, I, 65
mesacantha, I, 127, 132; IV, 260
mesacantha cymochila, I, 131
mesacantha grandiflora, I, 129
mesacantha greenei, I, 131
mesacantha macrorhiza, I, 130
mesacantha microsperma, I, 127
mesacantha oplocarpa, I, 131-144
mesacantha parva, I, 127, 128
mesacantha sphaerocarpa, I, 144
mesacantha stenochila, I, 132
mesacantha vaseyi, I, 146
metternichii, I, 176
mexicana, I, 34
micrarthra, I, 171
microcarpa, I, 144, 206
microdasys, I, 118, 120, 121, 122, 123, 136; IV, 259
microdasys laevior, I, 120
microdasys minor, I, 120
microdasys monstrosa, IV, 259
microdasys rufida, I, 122
microdisca, I, 133, 135, 136
microthele, I, 73
mieckleyi, I, 156, 158
militaris, I, 102, 104, 163
millspaughii, I, 202, 204, 206
minima americana, I, 102
minor, I, 139
minor catilescens, I, 102
micuclii, I, 78
missouriensis, I, 199, 200; IV, 264
missouriensis albispina, I, 199, 200
missouriensis elongata, I, 214
missouriensis erythrostemma, I, 214
missouriensis microsperma, I, 199, 200
missouriensis platycarpa, I, 199, 200
missouriensis rufispina, I, 199, 200

Opuntia—*continued*
missouriensis salmonea, I, 214
missouriensis subinermis, I, 199
missouriensis trichophora, I, 195
missouriensis watsonii, IV, 264
modesta, I, 67
mojavensis, I, 140, 145
molesta, I, 54, 60, 62, 66, 67
monacantha, I, 127, 156, 157; IV, 262
monacantha deflexa, I, 156
monacantha gracilior, I, 156, 157; IV, 262
monacantha variegata, 157; IV, 262
moniliformis, I, 33, 202, 206, 207, 208; IV, 264
montana, I, 214
montevicensis, I, 107, 109
monticola, I, 95
morenoi, I, 214
morisii, I, 128
mortolensis, I, 47
multiflora, I, 113, 114
myriacantha, I, 150, 152, 214
nana, I, 127, 217
napolea, IV, 265
nashii, I, 106, 163, 202, 203; IV, 264
nelsonii, I, 172
nemoralis, I, 102, 104
neorbuscula, I, 50
nigricans, I, 152, 153
nigrispina, I, 90, 97
nigrita, I, 183, 184
nopalilla, I, 38
oblongata, I, 173
obovata, IV, 264
occidentalis, I, 140, 146, 147, 153, 163, 165
ochrocentra, IV, 262
oligacantha, I, 89, 182
oplocarpa, I, 132; IV, 260
opuntia, I, 43, 126, 127, 128, 217; IV, 259
orbiculata, I, 176; IV, 263
orbiculata metternichii, I, 176
ottonis, I, 214, 215; III, 195
ovallei, I, 95, 214
ovata, I, 90, 95, 96; IV, 256
ovata leonina, IV, 256
ovoides, I, 95
pachona, I, 184
pachyarthra flava, I, 214
pachyclada rosea, I, 214
pachyclada spaethiana, I, 214
pachypus, I, 75, 77
pallida, I, 60, 65, 66
palmdora, I, 202
palmeri, I, 161
pampeana, I, 134
papyracantha, I, 89, 90; IV, 255
paraguayensis, I, 158
parishii, I, 57, 79, 81, 82
parkeri, I, 58
parmentieri, I, 95
parote, I, 214
parryi, I, 56, 57, 58, 81; IV, 254
parva, I, 161
parvispina, I, 157
parvula, I, 178
pascoensis, I, 100, 101
patagonica, I, 41
pelaguensis, I, 90
penicilligera, I, 135; IV, 261
pennellii, I, 110, 115, 219
pentlandii, I, 71, 72, 77, 90, 97, 98; IV, 256
perita, I, 65, 66

Opuntia—*continued*,
pes-corvi, I, 104
pestifer, I, 102, 103, 217, 218, 220; II, 19
phaeacantha, I, 139, 140, 142, 144; IV, 262
phaeacantha brunnea, I, 144
phaeacantha major, I, 144; IV, 262
phaeacantha nigricans, I, 144
philippii, I, 41
phyllacantha, I, 96
phyllanthus, I, 215; IV, 187
picardae, IV, 264
piccolomini, I, 214
piccolominiana, I, 191
pilifera, I, 176, 177, 184
pintadera, I, 176
pisciformis, IV, 258
pittache, I, 29
pittieri, I, 181, 188, 189
platyacantha, I, 33, 89; IV, 255
platyacantha deflexispina, I, 89, 90
platyacantha gracilior, I, 89
platyacantha monvillei, I, 89
platyclada, I, 214
platynoda, I, 109
plumbea, I, 126, 131
plumosa nivea, I, 89, 90
poepigii, I, 40, 41
pollardii, I, 126, 221
polyacantha, I, 193, 195, 196, 197, 199, 200; IV, 264
polyacantha albispina, I, 199
polyacantha borealis, I, 199
polyacantha microsperma, I, 200
polyacantha platycarpa, I, 199
polyacantha rufispina, I, 200
polyacantha trichophora, I, 195
polyacantha watsonii, I, 199
polyantha, I, 113, 114, 115; IV, 259
polymorpha, I, 89; II, 138
porterii, I, 28
pottsii, I, 137, 138; IV, 262
praecox, I, 214
prate, I, 191
procumbens, I, 160, 167
prolifera, I, 61, 67, 69, 70; IV, 254
prostrata, I, 128
prostrata spinosior, I, 214; IV, 263
protracta, I, 214
protracta elongata, I, 214
pruinosa, I, 191
pseudococcinellifer, I, 153
pseudotuna, I, 184, 214; III, 238
pseudotuna elongata, I, 214
pseudotuna spinosior, I, 214
puberula, I, 116, 117, 121, 122; IV, 259
pubescens, I, 100, 101, 173; IV, 257, 259
pulchella, I, 79, 82; IV, 255
pulverata, I, 214
pulverulenta, I, 78
pulverulenta miquelii, I, 78
pulvinata, I, 120
pumila, I, 100, 101; IV, 256
purpurea, I, 97
pusilla, I, 102, 105, 106; IV, 255, 256
pycnacantha, IV, 259
pycnantha, I, 118, 123; IV, 259
pycnantha margaritana, I, 123
pyriformis, I, 160, 168
pyrocarpa, I, 166

Opuntia—*continued*,
pyrrhacantha, I, 97
quimilo, I, 181, 190, 191
quipa, I, 125
quitensis, I, 149, 154, 224
rafinesquei, I, 127, 128, 129; IV, 259, 260
rafinesquei arkansana, I, 127, 129; IV, 259
rafinesquei cymochila, I, 131
rafinesquei cymochila montana, I, 131
rafinesquei fusiformis, I, 130
rafinesquei grandiflora, I, 129
rafinesquei greenei, I, 132
rafinesquei macrorhiza, I, 131
rafinesquei microsperma, I, 127, 199
rafinesquei minor, I, 127, 128
rafinesquei parva, I, 128
rafinesquei stenochila, I, 132
rafinesquei vaseyi, I, 146
rafinesquiiana, I, 127, 129; IV, 259
rafinesquiiana arkansana, I, 129; IV, 259
rahmeri, I, 94
ramosissima, I, 46, 48, 52; IV, 253
ramosissima cristata, IV, 253
ramosissima denudata, IV, 253
ramulifera, I, 47
rastera, I, 140, 149
raupiana, I, 90, 92
recedens, I, 128
recondita, I, 52, 53
recurvospina, I, 144
reflexa, I, 165
repens, I, 102, 103, 104, 115, 116
reptans, I, 214
reticulata, IV, 264
retroscina, I, 107, 109, 218
retrospinosa, I, 95
rhodantha, I, 193, 197, 198; IV, 264
rhodantha brevispina, I, 198
rhodantha flavispina, I, 198
rhodantha pisciformis, I, 198
rhodantha schumanniana, I, 198
riparia, I, 149
robusta, I, 182, 191, 192; IV, 264
robusta viridior, I, 191
rosea, I, 17, 63, 65, 78; IV, 254
roseana, I, 130
rositlora, I, 78
rotundifolia, I, 27; IV, 251
roxburghiana, I, 156
rubescens, I, 163, 202, 208, 209; IV, 264
rubiclora, I, 133, 146
rubrifolia, I, 144
rufescens, I, 175
rufida, I, 118, 119, 122
rugosa, I, 145
russellii, I, 90, 94
ruthei, I, 64
rutilla, I, 196; IV, 264
sabinii, I, 194
sacharosa, I, 14
salicornioides, I, 215; IV, 217
salmiana, I, 73, 74; IV, 254
salmii, I, 214
sanguinocula, I, 131
santa-rita, I, 140, 142
scheeri, I, 159, 176; IV, 262
schickendantzii, I, 74, 107
schomburgkii, I, 214
schottii, I, 79, 80, 81; IV, 255
schottii greggii, I, 80; IV, 255
schumannii, I, 90, 149, 155
schweriniana, I, 199, 200; IV, 264

- Opuntia*—continued,
 segethii, I, 75, 76, 79
 seguina, I, 130
 semispinosa, I, 147
 senilis, I, 86, 159, 176
 sericea, I, 134
 sericea coerulea, I, 134
 sericea longispina, I, 134
 sericea muhlenii, I, 134
 serpentina, I, 56, 57, 58, 69, 81; II, 224
 setispina, I, 45, 137, 138
 shaferei, I, 71, 72
 shreveana, I, 142
 sinclairii, I, 165, 166
 skottsbergii, I, 90, 96, 97
 soederstromiana, I, 149, 154, 221
 soehrensii, I, 133, 134, 135; IV, 261
 spathulata, I, 28, 29
 spathulata aquosa, I, 30
 speciosa, I, 214; IV, 205
 spegazzinii, I, 73, 74
 sphaerica, I, 90, 96; IV, 256
 sphaerocarpa, I, 193, 198, 199
 sphaerocarpa utahensis, I, 199
 spinalba, I, 130
 spinaurea, I, 214
 spiniflora, I, 215; II, 106
 spinosior, I, 52, 67, 68; IV, 254
 spinosior neomexicana, I, 68
 spinosissima, I, 103, 175, 202, 204, 205; IV, 264
 spinotecta, I, 64
 spinulifera, I, 178, 181, 182
 spinuliflora, I, 214
 spinulosa, I, 214
 spirocentra, IV, 265
 splendens, I, 199
 squarrosa, I, 165
 stanlyi, I, 79, 80; IV, 255
 stapelia, I, 66
 stapeliaceae, I, 65, 66; IV, 254
 stellata, I, 64
 stenarthra, I, 158
 stenochila, I, 126, 132
 stenopetala, I, 200, 201
 stipata, IV, 253
 straminea, I, 214
 streptacantha, I, 181, 184, 185; IV, 263
 stricta, I, 161, 178, 223
 stricta spinulescens, I, 214
 strigil, I, 136
 subarmata, I, 165, 166; IV, 263
 subferox, I, 175
 subinermis, I, 214
 subterranea, I, 90, 92
 subulata, I, 71, 75, 76, 77, 79; III, 179; IV, 254
 sulphurea, I, 133, 134, 150; IV, 260
 sulphurea laevior, I, 134
 sulphurea major, I, 134
 sulphurea minor, I, 134
 sulphurea pallidior, I, 134
 superbospina, I, 144
 syringacantha, I, 89
 taponia, I, 124, 160, 164
 tarapacana, I, 90, 94
 tardospina, I, 140, 141
 taylori, I, 102, 103
 tenajo, I, 49
 tenuispina, I, 137, 139
 teres, I, 71
 tesajo, I, 7, 48, 49
 tessellata, I, 46
 tessellata cristata, I, 46; IV, 253
 tessellata denudata, IV, 253
 testudinis-crus, I, 206
 tetracantha, I, 52, 53, 54
- Opuntia*—continued,
 texana, I, 165, 166
 thurberi, I, 52, 53, 54
 tidballii, I, 160
 todari, IV, 265
 tomentella, I, 173, 174
 tomentosa, I, 173, 174; IV, 263
 tortisperma, I, 131, 132
 tortispina, I, 126, 128, 131, 194; IV, 260
 toumeyii, I, 144; IV, 262
 tracyi, I, 102, 105; IV, 257
 treleasei, I, 118, 119
 treleasei kernii, I, 119
 triacantha, I, 110, 112, 113, 115, 204
 tribuloides, I, 86
 tricolor, I, 165, 166
 trichophora, I, 193, 195
 tuberculata, I, 214
 tuberiformis, I, 92
 tuberosa, I, 32, 33
 tuberosa albispina, IV, 212
 tuberosa spinosa, I, 89
 tuna, I, 210, 113, 114, 116, 149, 157, 163; IV, 259, 262, 263
 tuna humilior, IV, 259
 tuna humilis, I, 114
 tuna laevior, I, 114
 tuna orbiculata, I, 114
 tunicata, I, 60, 65, 66, 83, 121; IV, 254, 266
 tunicata laevior, I, 66
 tunoidea, I, 116, 162
 tunoides, I, 116
 turgida, IV, 265
 turpinii, I, 89
 turpinii polymorpha, I, 89
 tweediei, I, 134
 umbrella, I, 156
 undosa, I, 177, 279
 undulata, I, 65, 177, 179
 ursina, I, 174, 195, 196, 197; IV, 264
 ursus horribilis, IV, 264
 urumbella, I, 157
 utahensis, I, 298
 utkilio, I, 107, 109, 110
 vaginata, I, 47, 48; IV, 253
 valida, I, 147
 vaseyi, I, 140, 142, 145, 146
 velutina, I, 169, 172
 verschaffeltii, I, 44, 71, 72
 verschaffeltii digitalis, I, 71, 72
 versicolor, I, 44, 52, 54, 60, 62; IV, 254
 vestita, I, 72, 72, 87; IV, 254
 vexans, I, 64, 65
 vilis, I, 79, 82, 83
 violacea, I, 244
 virgata, I, 47
 viridiflora, I, 52, 55
 vivipara, I, 52
 vulgaris, I, 127, 128, 156, 157, 163, 177; IV, 259, 262
 vulgaris balearica, I, 161
 vulgaris major, I, 127
 vulgaris media, I, 127
 vulgaris minor, I, 127
 vulgaris nana, I, 127, 129
 vulgaris rafinesquei, I, 127
 vulgo, I, 157
 vulpina, I, 134
 wagneri, I, 74
 weberi, I, 84, 85
 wentiana, I, 110, 116
 wetmorei, IV, 255
 whipplei, I, 32, 43, 52, 55, 68; IV, 253
 whipplei laevior, I, 55
 whipplei spinosior, I, 68
 wilcoxii, I, 169, 172
 winteriana, I, 166
 wootonii, I, 147, 148
 wrightii, I, 51
- Opuntia*—continued,
 xanthoglochla, I, 130, 131
 xanthostemma, I, 198; IV, 264
 xanthostemma elegans, I, 198
 xanthostemma fulgens, I, 198
 xanthostemma gracilis, I, 198
 xanthostemma orbicularis, I, 198
 xanthostemma rosea, I, 198
 xerocarpa, I, 198
 youngii, I, 130
 zebrina, I, 149, 155, 222
 zacuapanensis, I, 183
 zuniensis, I, 144
 Opuntiacel, I, 217
 Opuntiae, IV, 252
 Opuntiacel, I, 8, 24; IV, 252
 Orbiculatae, I, 45, 176
 Orchids, II, 42
 Oreocereus, II, 2, 60, 108, 171-173; III, 77; IV, 274
 celsianus, II, 171, 172, 173, 226; IV, 279
 celsianus bruennowii, II, 171
 lanatus, II, 61
 Organito de vibora, II, 111
 Organo, II, 48, 74; IV, 273
 Oroya, III, 77, 102
 peruviana, III, 102
 Pachycereus, II, 2, 28, 68-76, 96
 calvus, II, 69; IV, 271
 chrysomallus, II, 69, 72, 73, 74; IV, 271
 columna-trajani, II, 69, 76
 gaumeri, II, 69, 71; IV, 271
 grandis, II, 69, 72
 lepidanthus, II, 69, 76; IV, 272
 marginatus, II, 69, 74, 75, 92, 102 IV, 271
 ocuttii, II, 69, 70
 pecten-aboriginum, II, 69, 70, 71, 72, 76; IV, 271
 pringlei, II, 69, 70 IV, 270, 271
 queretaroensis, II, 96
 ruficeps, II, 69, 75
 titan, II, 69; IV, 271
 Palmadora, I, 202
 Palmadorae, I, 45, 201
 Palmatoria, I, 202
 Paradoxae, IV, 221
 Pasacana, II, 133
 Passiflora, II, 92
 Pataquisca, I, 76
 Pediocactus, III, 77, 90, 91, 109
 simpsonii, III, 88, 90, 92; IV, 68, 286
 Peirescia, I, 9
 Peireskia, I, 9
 Pelecypora, IV, 3, 59, 60, 64
 aselliformis, IV, 8, 59
 aselliformis concolor, IV, 59
 aselliformis cristata, IV, 64
 aselliformis grandiflora, IV, 59
 aselliformis pectinata, IV, 64, 6
 aselliformis pectinifera IV, 64
 fimbriata, IV, 59
 micromerus, III, 93
 pectinata, IV, 64, 65
 pectinata cristata, IV, 65
 pectinifera, IV
 Pellote, III, 84, 107
 Pence, IV, 260
 Pencea-chica, IV, 260
 Peniocereus, II, 2, 112, 113
 greggii, II, 112; IV, 275
 johnstonii, IV, 275
 Penquilla, IV, 260
- Pentapterae, IV, 221
 Pentlandianae, I, 44, 90
 Peote, IV, 59
 Peotillo, IV, 59
 Perescia, I, 9
 Pereskia, I, 8-24, 25, 26, 40, 75; IV, 251, 252, 290
 acardia, I, 10
 aculeata, I, 10, 11, 14; IV, 251
 aculeata lanceolata, I, 10
 aculeata latifolia, I, 10
 aculeata longispina, I, 10
 aculeata rotunda, I, 10
 aculeata rotundifolia, I, 10
 aculeata rubescens, I, 10
 affinis, I, 24
 amapola, I, 14
 amapola argentina, IV, 251
 argentina, I, 14
 autumnalis, I, 9, 11, 12; IV, 251
 bahiensis, I, 9, 19, 20; IV, 252
 bleo, I, 4, 9, 17, 18, 20; IV, 251
 brasiliensis, I, 10
 calandriniaefolia, I, 29, IV, 252
 colombiana, I, 9, 17
 conzattii, I, 9, 24
 crassicaulis, I, 29
 cruenta, I, 24
 cubensis, I, 9, 22
 foetens, I, 10
 fragrans, I, 10
 galeottiana, IV, 251
 glomerata, I, 94
 godseffiana, I, 10, 11; IV, 251
 grandiflora, I, 24
 grandifolia, I, 9, 18, 19, 20, 21; IV, 251
 grandispina, I, 24
 guamacho, I, 9, 15, 16; IV, 251
 guatemalensis, IV, 251
 haageana, I, 24
 horrida, I, 9, 21, IV, 255
 humboldtii, IV, 251
 lanceolata, I, 10
 longispina, I, 10
 longispina rotundifolia, IV, 251
 longispina rubescens, IV, 251
 lychnidiflora, I, 9, 12, 13
 moorei, I, 9, 15
 nicoyana, I, 9, 13
 ochracarpa, I, 19
 opuntiaeflora, I, 26, 27
 panamensis, I, 17, 18
 pereskia, I, 7, 9, 10; IV, 251
 pflanzii, IV, 251
 philippii, I, 41
 pititache, I, 29
 plantaginea, I, 24
 poeppigii, I, 41; IV, 253
 portulacifolia, I, 9, 22, 23, 24
 recurvifolia, IV, 251
 rosea, I, 17
 rotundifolia, I, 27 IV, 251
 sacharosa, I, 9, 10, 14, 15; IV, 251
 spathulata, I, 28, 29; IV, 251
 subulata, I, 75, 76; IV, 175, 254
 tampicana, I, 9, 17
 undulata, I, 10
 verticillata, IV, 251
 weberiana, I, 9, 15
 zehntneri, I, 9, 13, 14; IV, 252
 zinniaeflora, I, 9, 17, 20, 21; IV, 251
 Pereskiae, I, 8, 24; IV, 252
 Pereskioopsis, I, 8, 14, 17, 24, 25-30, 43; IV, 252

Pereskopsis—*continued*,
aquosa, I, 25, 29,
autumnalis, I, 11, 12; IV, 251
brandegeei, I, 28
chapistle, I, 25, 27; IV, 251
diguetii, I, 25, 26, 27
kellermanii, I, 25, 30
opuntiaeflora, I, 25, 26, 27
pittitache, I, 25, 29; IV, 252
porteri, I, 25, 28; IV, 251
recurvifolia, IV, 251
rotundifolia, I, 25, 27, 28;
 IV, 251
scandens, IV, 252
spathulata, I, 25, 28; IV, 251
velutina, I, 25, 26
Pereskio-puntia, I, 25
Pertinato, III, 232
Peruviana, II, 4
Pest pear, I, 161, 163
Petaya, III, 17
Peyote, III, 84, 107, 184; IV, 59
Peyotillo, IV, 59
Pfeiffera, IV, 208, 209, 210, 211
cereiformis, IV, 210, 211
ianthothele, IV, 208, 210
rhaphaloides, IV, 209
Phaeacanthae, I, 45, 136, 139,
 141, 148
Phellosperma, IV, 3, 60, 65, 156
tetrancistra, IV, 60, 61
Phyllanthos, IV, 205
Phyllanthus, IV, 186
Phyllanthus, IV, 186
Phyllocactus, II, 3, 221, 222;
 IV, 185, 186, 198, 200
ackermannii, IV, 198
ackermannii hybridus, IV, 200
ackermannii major, IV, 200
acuminatus, IV, 188, 189
acutifrons, IV, 200
agatha, IV, 200
alatus, IV, 200
alatus major, IV, 200
albus grandiflorus, IV, 199
albus grandissimus, IV, 200
albus perfectus, IV, 200
albus superbiens, IV, 200
albus superbissimus, IV, 200
albus superbus, IV, 199
alexandrinae, IV, 200
amabilis, IV, 200
amabilis perfectus, IV, 200
amarantinus, IV, 200
angularis, IV, 192
anguliger, IV, 191, 192
arnoldi, IV, 200
atrosanguineus, IV, 200
aurantiacus, IV, 200
aurantiacus superbus, IV, 200
belgicus, IV, 200
bergei, IV, 200
bergeri, IV, 200
biformis, IV, 202
billiardieri, IV, 200
binderi, IV, 200
blindlii, IV, 200
blindlii, IV, 200
boehmii, IV, 200
boliviensis, IV, 200
bolivillanensis, IV, 200
bothii, IV, 200
bradei, IV, 204
brongnartii, IV, 200
buestii, IV, 200
burmeisteri, IV, 200
campannii, IV, 200
caparti, IV, 200
capelleanus, IV, 200
carolus magnus, IV, 200
cartagensis, IV, 197
cartagensis refractus, IV, 197
cartagensis robustus, IV, 197
castneri, IV, 200
caudatus, IV, 190
caulorrhizus, IV, 192
chiapensis, IV, 203, 204

Phyllocactus—*continued*,
chico, IV, 200
coccineus, IV, 200
colmariensis, IV, 200
colombiensis, IV, 200
cooperi, IV, 199
costaricensis, IV, 193
courantii, IV, 200
crassuliefolius, IV, 200
crenato grandiflorus, IV, 199
crenatus, IV, 192, 193
crenatus amarantinus, IV,
 192
crenatus amarantinus, IV,
 200
crenatus caulorrhizus, IV, 200
crenatus elegans, IV, 192
crenatus erleri, IV, 192
crenatus haageanus, IV, 192
crenatus hirsutis, IV, 200
crenatus lateralis, IV, 200
crenatus lateritius, IV, 192
crenatus latifolius, IV, 200
crenatus luteus, IV, 200
crenatus roseus, IV, 192
crenatus ruber, IV, 200
crenatus splendens, IV, 192
crenatus superbus, IV, 292
crenatus vogelii, IV, 192
crispielsi, IV, 200
curtisii, IV, 200
dangeli, IV, 200
darrahii, IV, 190, 191
decumbens, IV, 200
delicatus, IV, 179
demouline, IV, 200
deveauxi, IV, 200
dieffenbachianus, IV, 200
dolores, IV, 200
dumouline, IV, 200
edwardsii, IV, 200
eichlamii, IV, 203
elegans, IV, 200
epirus, IV, 200
erebus, IV, 199
erectus perfectus, IV, 200
erectus superbus, IV, 200
ernesti, IV, 200
erubescens, IV, 200
fastuosus, IV, 200
feastii, IV, 200
felonis, IV, 200
feltoni, IV, 200
floribundus, IV, 200
formosus, IV, 200
franzii, IV, 200
fuentii, IV, 200
funkii, IV, 200
gaertneri, IV, 183, 184
gaillardae, IV, 187
germanus, IV, 200
gloriosus, IV, 200
gordonianus, IV, 200
grandidissimus, IV, 200
grandiflorus, IV, 200
grandiflorus albiflorus, IV,
 200
grandiflorus ruber, IV, 200
grandilobus, IV, 192
grandis, IV, 188, 189
guatemalensis, IV, 195
guebwillerianus, IV, 200
guedeneyi, IV, 200
guentneri, IV, 200
guyanensis, IV, 188
haagei, IV, 199
hamburgiensis, IV, 200
hansii, IV, 200
hauffii, IV, 200
havermansii, IV, 200
helenus, IV, 200
hempeli, IV, 200
hibridus, IV, 200
hibridus gordonianus, IV, 199
hibridus wrayi, IV, 199
hildmanni, IV, 199
hiapensis, IV, 200

Phyllocactus—*continued*,
hookeri, IV, 197
ignescens, IV, 200
incomparabilis miniatus, IV,
 200
jenkinsonii, IV, 200
jenkinsonii superbus, IV, 200
johnsonii, IV, 200
jordanis, IV, 200
kampmannii, IV, 200
keithii, IV, 200
kermesinus magnus, IV, 200
kiardi, IV, 200
kranzii, IV, 200
krausei, IV, 200
laarsenii, IV, 200
laetingii, IV, 200
laloyi, IV, 200
latifrons, IV, 188, 189
laudowi, IV, 200
leopoldii, IV, 200
lepidocarpus, IV, 194
londonii, IV, 189
longipes, IV, 189
lorentzii, IV, 200
lorenzii, IV, 200
luthii, IV, 189
lodmani, IV, 200
ludwigi, IV, 200
lunus, IV, 200
macquianus, IV, 189
macrocarpus, IV, 193
macrolobus, IV, 192
macropteris, IV, 193
maclenii, IV, 289
magnificus, IV, 200
maigretii, IV, 200
makoyi, IV, 200
marginatus, IV, 297
marsus, IV, 199
mauritanus, IV, 189
mayanus, IV, 200
mexicanus, IV, 189, 200
meyerianus, IV, 200
muehlenpfordtii, IV, 200
muelhousianus, IV, 200
multiflorus, IV, 200
nelsonii, IV, 203
neubertii, IV, 200
niedtii, IV, 200
niger, IV, 200
nitens, IV, 200
nymphorea beata, IV, 200
oxypetalus, IV, 188, 189
paraguayensis, IV, 200
pentneri, IV, 200
persdorfii, IV, 199
phyllanthoides, IV, 200, 205
phyllanthoides albiflorus, IV,
 205
phyllanthoides striatus, IV,
 205
phyllanthoides striatus multi-
florus, IV, 205
phyllanthus, IV, 187, 188
phyllanthus boliviensis, IV,
 187
phyllanthus columbiensis, IV,
 187
phyllanthus paraguayensis,
 IV, 187
phyllantoides, IV, 200
phyllantoides crenatus, IV,
 200
pittieri, IV, 194
potstachianus, IV, 200
poulettianus, IV, 200
pressleri, IV, 200
pulcherrimus, IV, 201
pumilus, IV, 185
purpureus, IV, 200
purpusii, IV, 188
quiliardetti, IV, 201
raveaudii, IV, 201
rebuti, IV, 201
reichei, IV, 201
reinckii, IV, 207

Phyllocactus—*continued*,
roseus albus, IV, 189
roseus carmineus, IV, 201
roseus carneus, IV, 201
roseus floribundus, IV, 201
roseus grandiflorus, IV, 199
roseus grandissimus, IV, 199
roseus miniatus, IV, 201
roseus splendidus, IV, 201
roseus superbus, IV, 189
roydii, IV, 201
ruber, IV, 201
ruber perfectus, IV, 201
ruber violaceus, IV, 201
ruelcheri, IV, 200
ruetii, IV, 199
russellianus, IV, 184, 185
sarniensis, IV, 201
schaffieri, IV, 201
schallerianus, IV, 201
schlimmi, IV, 201
schmidtii, IV, 201
selloi, IV, 189
sellowii, IV, 201
serratus, IV, 191
smithii, IV, 189
smoli, IV, 189
specillimus, IV, 201
speciosissimus, IV, 201
speciosissimus feltonii, IV,
 200
speciosissimus grandiflorus,
 IV, 201
speciosus roseus, IV, 201
splendens, IV, 201
splendidus, IV, 201
stenesii, IV, 201
stenopetalus, IV, 196, 197
strictus, IV, 196
superbissimus, IV, 201
superbus, IV, 200
tettanii, IV, 201
thomasianus, IV, 193
tonduzii, IV, 200
tricolor, IV, 201
triumphans, IV, 200
tuna, IV, 200
ulbrechtii, IV, 200
undiflorus, IV, 201
vandesii, IV, 201
victoria-regia, IV, 200
vitellimus, IV, 201
vogelii, IV, 200
vonhoffii, IV, 201
warszewiczii, IV, 201
weingartii, IV, 200
wippenmannii, IV, 200
wittmackianus, IV, 201
wrayi, IV, 200
zarka, IV, 200
Phyllocereus, II, 221; IV, 185
Pilocereus, II, 3, 13, 25, 44,
 51, 52, 58, 59, 178, 181;
 III, 60, 78; IV, 185, 186,
 268, 271
acranthus, II, 169
albisetosus, II, 58; IV, 270
albispinus, II, 59
albispinus crenatus, II, 5
alensis, II, 55
andryanus, II, 59
angulosus, IV, 268
arrabidae, II, 42
auratus, IV, 268
barbatus, II, 50
bruennowii, II, 171, 172; IV,
 279
celsianus, II, 171, 172, 226;
 IV, 279
celsianus bruennowii, II, 171,
 172, 226
celsianus fossilatus, II, 272
celsianus gracilior, II, 171
celsianus lanuginosior, II,
 171, 172
celsianus williamsii, II, 171
chrysacanthus, II, 48

Pilocereus—continued,
chrysomallus, II, 59, 72; IV, 271
coeruleus, II, 5
columna, II, 59, 76
columna-trajani, II, 76
cometes, II, 52; IV, 269
consolei, II, 44
crenulatus, II, 49
curtisii, II, 44; IV, 269
dautwitzii, II, 61, 63, 225; IV, 270
dautwitzii cristatus, II, 63
divaricatus, II, 151
engelmannii, II, 164
erythrocephalus, III, 7
euphorbioides, II, 33
exerens, II, 42, 43; IV, 268
fimbriatus, II, 151
flavicomus, II, 52
flavispinus, II, 60
floccosus, II, 50
foersteri, II, 52
fossulatus, II, 171, 172; IV, 279
fossulatus gracilis, II, 171
fossulatus pilosior, II, 171
fouachianus, II, 50
foveolatus, II, 172
fulviceps, II, 72, 73; IV, 271
fulvispinosus, II, 50
ghiesbreghtii, II, 60
giganteus, II, 164, 167; IV, 279
glaucescens, II, 59
gounellei, II, 34
grandispinus, II, 87
haageanus, II, 62
haagei, II, 61
hagendorpi, II, 27
haworthii, II, 44
hermentianus, II, 58; IV, 270
hoogendorpii, II, 225
hoppenstedtii, II, 27, 28; IV, 268
houlettianus, II, 3; IV, 269
houlettianus leucocephalus, IV, 269
houlettianus niger, IV, 269
houlettii, II, 52, 53; IV, 269
jubatus, II, 52
kanzleri, II, 171; IV, 279
kranzleri, IV, 279
lanatus, II, 61, 63; IV, 270
lanatus cristatus, II, 63; IV, 270
lanatus haagei, II, 61, 63
lanuginosus, II, 49
lanuginosus virens, IV, 269
lateralis, II, 27, 28
lateribarbatulus, II, 76
leninghausii, III, 204, 205
leucocephalus, IT, 52
lutescens, II, 44
macrocephalus, II, 31
macrostibas, II, 18
marschalleckianus, II, 53
melocactus, II, 29, 30
militaris, II, 73
monacanthus, II, 155
moritzianus, II, 41
niger, II, 4; IV, 269
niger aureus, IV, 269
nigricans, II, 44
nobilis, II, 44
oligogonus, IV, 268
oligogonus houlettianus, IV, 268
oligogonus sublanatus, IV, 268
pascana, II, 133
pentaedrophorus, II, 31
pfeifferi, II, 224
plumieri, II, 47
polyedrophorus, II, 31
polygonus, II, 47
polylophus, II, 32

Pilocereus—continued,
polyptychus, IV, 269
pringlei, II, 69
repandus, II, 17, 18
rhodacanthus, III, 7
robinii, II, 39
royenii, II, 50, 31
royenii armatus, II, 50
ruficeps, II, 75
russellianus, II, 33, 34
sargentianus, II, 177, 178; IV, 280
schlumbergeri, II, 47
schottii, II, 177, 178; IV, 280
scoparius, II, 41
senilis, II, 27; IV, 268
senilis cristatus, II, 27
senilis flavispinus, II, 27
senilis longispinus, II, 27
setosus, II, 34, 35
sterkmannii, II, 41
straussii, II, 171, 172, 226
strictus, II, 44, 60
strictus consolei, II, 44
strictus fouachianus, II, 50
sublanatus, IV, 268
swartzii, II, 47
terscheckii, II, 140
tetetzoo, II, 76
tetetzoo cristatus, IV, 272
thurberi, II, 97
tilophorus, IV, 268
trichacanthus, II, 44
ulci, II, 52
urbanianus, II, 43
vellozoi, II, 29
verheinei, II, 58
virens, II, 43
williamsii, II, 172; IV, 279
Pin pillow, I, 102
Piptanthocereus, II, 3, 22
azureus, II, 15
beneckeii, II, 18; IV, 273
chalybaeus, II, 16
forbesii, II, 7
hankeanus, II, 7
labouretianus, II, 7
jamacaru, II, 8
jamacaru caesius, II, 15
jamacaru cyaneus, II, 8
jamacaru glaucus, II, 8
peruvianus, IV, 266
peruvianus monstrosus, IV, 266
spiegazzinii, IV, 268
validus, II
Pisciformes, IV, 258
Piscos colorado, II, 62
Pitahaya, II, 96, 98, 122, 165; III, 40; IV, 190
agre, II, 117, 122
agria, II, 117, 122
de San Juan, II, 122
dulce, II, 98, 122
Pitahayita, II, 111, 122
Pitajaya, II, 122
Pitajua, II, 122
Pitalla, II, 122
Pitaya, II, 122
Pitayita, II, III, 122
Pithaya, II, 122
Platanillo de monte, IV, 197
Platyopuntia, I, 45, 73, 84, 92, 99, 100, 135, 211; IV, 180
Polar bear cactus, I, 87
Polyacanthae, I, 45, 193
Portulaca, I, 9
Prickly pear, I, 43, 212; IV, 250
Prismaticae, IV, 220
Pseudorhipsalis, IV, 208, 213-214, 243
alata, IV, 213, 214
himantoclada, IV, 213, 214
Pterocactus, I, 24, 30-33
decipiens, I, 32, 33
fischeri, I, 31
hickenii, I, 31

Pterocactus—continued,
kuntzei, I, 30, 32, 33; IV, 252
kurtzei, I, 32
pumilus, I, 31, 32
tuberosus, I, 31, 32, 33; IV, 252
valentinii, I, 88
Pubescentes, I, 141
Puipute, I, 13
Pumilae, I, 45, 100; IV, 256
Queen of the night, IV, 283
Quiabentia, IV, 252
Quiabentia zehntneri, IV, 252
Quiabento, I, 14; IV, 252
Quimilo, I, 190
Quipa, I, 125
Raho de raposa, II, 109
Rainbow cactus, III, 19, 27
Ramosissimae, I, 44, 46
Rat tail cactus, II, 218
Rathbunia, II, 2, 139, 169, 170
alamosensis, II, 117, 169, 170
kerberi, II, 169, 170
sonorensis, II, 169
Rebutia, III, 3, 45-48, 60, 176; IV, 285
deminuta, IV, 285
fielbrigii, III, 45, 46, 205
minuscule, III, 45, 46, 180; IV, 285
pseudominuscula, III, 45, 46, 47
pygmaea, III, 45, 47
steinmannii, III, 45, 47
Recurvatae, IV, 24, 51
Repandae, II, 4
Rhipsalidanae, II, 1; IV, 177, 206, 208
Rhipsalidopsis, IV, 208, 209
rosea, IV, 209, 247
Rhipsalis, I, 8; II, 3, 124, 209, 214, 222; III, 4; IV, 180, 208, 209, 210, 211, 212, 213, 214, 216, 219-247, 250
aculeata, IV, 221, 231
aethiopica, IV, 226
alata, IV, 213, 214, 241, 242, 243
alternata, IV, 237
anceps, IV, 215, 216
angustissima, IV, 241, 242
bambusoides, IV, 218
biolleyi, II, 215
boliviana, IV, 221, 240
brachiata, IV, 222
brevibarbis, IV, 215
bucheni, IV, 247
burchellii, IV, 220, 225
calmaiformis, IV, 231
campos-portoana, IV, 220, 224
capilliformis, IV, 220, 224
caripensis, IV, 227
carnosa, IV, 247
cassutha, IV, 219, 220, 225, 226, 233
cassutha rhodocarpa, IV, 226
cassutha, IV, 225
cassutha dichotoma, IV, 226
cassutha hookeriana, IV, 226
cassutha major, IV, 227, 234
cassutha mauritiana, IV, 226
cassutha mociniana, IV, 226, 227
cassutha pendula, IV, 226, 227
cassutha pilosiuscula, IV, 227, 234
cassutha rhodocarpa, IV, 226
cassutha swartziana, IV, 226
cassutha tenuior, IV, 227
cassythoides, IV, 227
cassythoides, IV, 226
cavernosa, IV, 215, 216
cereiformis, IV, 210
cereuscula, IV, 220, 222
chloroptera, IV, 243
chrysantha, IV, 236, 247
chrysocarpa, IV, 236

Rhipsalis—continued,
clavata, IV, 220, 223
clavata delicatula, IV, 223, 224
comorensis, IV, 226
conferta, IV, 227, 228
coriacea, IV, 221, 241, 242
crassa, IV, 243
cribrata, IV, 220, 225
cribrata filiformis, IV, 225, 247
crispa, IV, 243
crispa latior, IV, 245
crispa major, IV, 245
crispata, IV, 221, 245
crispata latior, IV, 245
crispimarginata, IV, 221, 244, 245
cuneata, IV, 221, 246
cylindrica, IV, 231
densiareolata, IV, 228
dichotoma, IV, 226
dissimilis, IV, 221, 236, 247
dissimilis setulosa, IV, 236
echinata, IV, 222
elliptica, IV, 220, 221, 242, 243
elliptica helioides, IV, 213
ensiformis, IV, 215
erythrocarpa, IV, 228
erythrolepis, IV, 228, 247
fasciculata, IV, 220, 227, 229
filiformis, IV, 247
floccosa, IV, 221, 227, 234
floribunda, IV, 228
foveolata, IV, 236
frondosa, IV, 247
funalis, IV, 231, 234
funalis gracilior, IV, 231
funalis gracilis, IV, 235
funalis minor, IV, 234
gibberula, IV, 221, 235
gonocarpa, IV, 221, 238
gracilis, IV, 217, 224
grandiflora, IV, 221, 231, 232, 233
grandiflora minor, IV, 234
hadrosoma, IV, 231
harrisii, IV, 2
heteroclada, IV, 220, 224
himantoclada, IV, 213
hookeriana, IV, 226
horrida, IV, 229
houlettiana, IV, 221, 238
houlettii, IV, 238
ianthothole, IV, 210
itatiaiae, IV, 247
jamaicensis, IV, 221, 242
kegnelii, IV, 238
knightii, IV, 215
lagenaria, IV, 247
larmentacea, IV, 230
leicophloea, IV, 241
leucorhaphis, IV, 221, 232
lindbergiana, IV, 220, 228
linearis, IV, 221, 239
loefgrenii, IV, 221, 232, 233
lorentziana, IV, 221, 240
lumbicoides, IV, 220, 230
macahensis, IV, 247
macrocarpa, IV, 187
macropogon, IV, 215, 216
madagascariensis, IV, 229
madagascariensis dasycerca, IV, 229
megalantha, IV, 221, 232
mesembrianthoides, IV, 222
mesembryanthemoides, IV, 220, 222
micrantha, IV, 221, 237, 239, 247
microcarpa, IV, 247
minutiflora, IV, 226, 227
miquelii, IV, 247
mittleri, IV, 216
monacantha, IV, 212, 213
myosurus, IV, 215, 216

- Rhipsalis*—*continued*,
nevasesii, IV, 232
neves-armondii, IV, 221, 233
novasesii, IV, 232
oblonga, IV, 221, 246
oligosperma, IV, 247
pacheco-leonii, IV, 236
pachyptera, IV, 221, 243, 247
pachyptera crassior, IV, 243
pachyptera purpurea, IV, 243
paradoxa, IV, 221, 237
parasitica, IV, 226, 227, 229
pendula, IV, 225, 227
penduliflora, IV, 225
penduliflora laxa, IV, 225
pentagona, IV, 236
pentaptera, IV, 221, 236
peruviana, IV, 212
phyllanthus, IV, 187
pilocarpa, IV, 208, 209
pilosa, IV, 229
pittieri, IV, 221, 233
platycarpa, IV, 212, 221, 242
prismatica, IV, 220, 222
pterocharpa, IV, 239
pterocharpis, IV, 237
pulcherrima, IV, 230
pulchra, IV, 220, 229
pulvinigera, IV, 221, 227, 233, 234
puniceo-discus, IV, 221, 235, 247
purpureus, IV, 221, 241
radicans, IV, 215, 216
radicans anceps, IV, 215
radicans ensiformis, IV, 215
radicans rosea, IV, 215
ramosissima, IV, 216
ramulosa, IV, 214, 221, 240, 241
regnelliana, IV, 238
regnellii, IV, 238
rhombea, IV, 241, 244, 245, 247
rhombea crispata, IV, 245
riedeliana, IV, 247
rigida, IV, 233
robusta, IV, 231, 243
rosea, IV, 209, 210, 247
roseana, IV, 221, 246
rugulosa, IV, 234
russelli, IV, 221, 242
saglionis, IV, 222, 223, 247
saglionis rubrodiscus, IV, 222
salicorne, IV, 218
salicornioides, IV, 216, 217, 218, 222, 235
salicornioides bambusoides, IV, 218, 219
salicornioides gracilior, IV, 217
salicornioides gracilis, IV, 217
salicornioides ramosior, IV, 217
salicornioides stricta, IV, 217
salicornioides strictior, IV, 217
salicornioides viligera, IV, 218
salicornioides, IV, 222
sansibarica, IV, 226, 227
sarmentacea, IV, 230
sarmentosa, IV, 230
schottmuelleri, IV, 218
setulosa, IV, 236
shaferei, IV, 220, 228, 229
simmleri, IV, 220, 223
spatulata, IV, 247
squamulosa, IV, 215, 216
stricta, IV, 217
suarensis, IV, 222
suareziana, IV, 222, 245
- Rhipsalis*—*continued*,
sulcata, IV, 221, 235, 236, 239
swartziana, IV, 213, 244
taglionis, IV, 247
teres, IV, 220, 227
tetragona, IV, 222
tonduzii, IV, 221, 239, 240
trigona, IV, 221, 237, 247
tucumanensis, IV, 221, 234
turpinii, IV, 247
undulata, IV, 226, 227
virgata, IV, 220, 227
warmingiana, IV, 221, 238
wercklei, IV, 240, 247
wettsteinii, IV, 247
zanzibarica, IV, 226
- Ringent-flowered cactus, IV, 178
Robustae, I, 45, 191
Sacacil, I, 214
Sacamatraca, II, 111
Sacasil, II, 111
Sacharosa, I, 10, 14
Sacred mushroom, III, 84
Saguaro, II, 105
Sahuaro, II, 165, 166
Salmianae, I, 44, 73, 75
Saman, II, 223
Sandillon, III, 186
Saramatraca, II, 111
Scheerianae, I, 45, 159
Schlumbergera, IV, 177, 182-185
epiphyllodes, IV, 184
gaertneri, IV, 182, 183, 184
russelliana, IV, 182, 183, 184
Sclerocactus, III, 78, 212-215
polycancistrus, III, 212, 213, 214
whipplei, III, 212, 213
Sehucan, II, 81
Selenicereus, II, 1, 58, 183, 195, 196-210, 221; IV, 249
boeckmannii, II, 197, 202; IV, 283
brevispinus, II, 197, 201, 202
coniflorus, II, 197, 198, 199; IV, 283
donkelaari, II, 197, 200
grandiflorus, II, 128, 196, 197, 198, 201, 210, 218; IV, 199, 283
hamatus, II, 196, 197, 203, 204; IV, 283
hondurensis, II, 197, 199
inermis, II, 197, 207, 208, 209
kunthianus, II, 197, 201
macdonaldiae, II, 196, 197, 202, 203; IV, 283
maxonii, II, 198; IV, 283
miravallensis, II, 213
murrillii, II, 197, 206
nelsonii, IV, 283, 284
pringlei, II, 199
pteranthus, II, 111, 196, 197, 200; IV, 283
spinulosus, II, 197, 207
urbanianus, II, 197, 198, 212; IV, 283
vagans, II, 197, 205, 206, 207
wercklei, II, 197, 208, 209
- Sempervivum tomentosum*, I, 41
Setispinae, I, 45, 136
Sinita, II, 177
Snake cactus, III, 90
Snowy cactus, III, 232; IV, 7
Solisia, IV, 3, 64, 65, 250
pectinata, IV, 64
Soroco, II, 62
Sowesa, IV, 270
Spanish dildos, II, 87
Spear-shaped opuntia, I, 179
Spineless opuntia, IV, 253
- Spinosissimae*, I, 43, 45, 201, 202, 203, 208; IV, 264
Stenocactus, III, 109
Stenocereus, II, 69, 85
stellatus, II, 92
stellatus tenellianus, II, 92
stellatus tonellianus, II, 92
Stenogoni, III, 118, 123
Stenopetalae, I, 45, 200
Stenopuntia, I, 200
Stetsonia, II, 2, 64, 65
coryne, II, 64, 65; IV, 270
Strawberry cactus, III, 37
Streptacanthus, I, 45, 112, 156, 177, 181, 192, 225
Strigiles, I, 45, 136
Stromatocactus, III, 80
kotschoubeyi, III, 82
Strombocactus, III, 78, 106, 107
disciformis, III, 106; IV, 287
Strophocactus, II, 183, 221, 222
wittii, II, 221, 222
Suaharo, II, 165
Subulatae, I, 44, 71, 75
Sucker, I, 43, 103
Suguro, II, 165
Sulcatae, IV, 221
Sulcolanatae, IV, 24
Sulphureae, I, 45, 133; IV, 261
Suwarro, II, 165
Suwarrow, II, 165
Tacinga, I, 24, 39, 40
funalis, I, 38, 39; IV, 266
Tail of the fox, II, 109
Tapuna pear, I, 192
Tasajillo, I, 26, 30
Tasajo, I, 43
macho, I, 63
Tenuiores, II, 22
Teonanactl, III, 84
Tephrocactus, I, 42, 43, 44, 71, 72, 79, 84, 85, 90, 95, 97, 106, IV, 256
andicolus, I, 89
aoracanthus, I, 91
calvus, I, 89
diadematus, I, 43, 89
platycanthus, I, 89; IV, 255
polycanthus, IV, 255
pusillus, I, 106
retrospinosus, I, 95; IV, 256
retrospinosus, IV, 256
turpinii, I, 89
Teretes, I, 71
Thelocactus, III, 148; IV, 3, 6-13
buckii, IV, 6, 8, 9
bicolor, IV, 6, 11
fossulatus, IV, 6, 10
hexadrophorus, IV, 6, 10, 11
leucacanthus, IV, 6, 8, 9
lloydii, IV, 6, 11
lophothele, IV, 6, 7, 8
nidulans, IV, 6, 9, 10
phymatothele, IV, 6, 7, 8
pottsii, IV, 6, 12
rinconensis, IV, 6, 7, 8
tulensis, IV, 6, 11
Theloides, IV, 13
Thurberianae, I, 44, 52
Tomentosae, I, 45, 172
Torch thistle, III, 237
Tortispinae, I, 45, 104, 126, 130, 133, 136, 193; IV, 265
Tortuosii, II, 77
Toumeya, III, 77, 91, 92
papyracantha, III, 91
Trichocereus, II, 2, 92, 130-146; III, 3, 60, 75; IV, 268
bridgesii, II, 130, 134, 136
candicans, II, 130, 134, 142; IV, 277
chiloensis, II, 130, 137, 138; IV, 266
- Trichocereus*—*continued*,
coquimbatus, II, 130, 138, 139; IV, 277
cuzcoensis, II, 130, 136
fascicularis, II, 64, 130, 141
huascha, II, 130, 141, 142; III, 57
lamprochlorus, II, 130, 132, 133
macrogonus, II, 130, 136; IV, 277
pachanoi, II, 130, 134, 135
pasacana, II, 130, 132, 133, 134, 140, 145, 225; IV, 277
peruvianus, II, 130, 136
schickendantzii, II, 130, 144; IV, 278
shaferei, II, 130, 144
spachianus, II, 130, 131, 132; IV, 277
strigosus, II, 130, 143, 144
terscheckii, II, 130, 140; IV, 277
thelegonoides, II, 130, 131
thelegonus, II, 130, 131
Trigonae, IV, 221
Tuna, I, 5, 43, 114, 181, 186; II, 66, 93; IV, 257
cardona, I, 184
colorado, II, 101
de agua, I, 30
de cohado, II, 84
de Espana, I, 224
elator, I, 153
major, I, 163
Tunae, I, 45, 110, 116, 148
Tunilla, IV, 261
Turk's cap, III, 221, 231
Turk's head, III, 221, 231
Utahia, III, 78, 215
sileri, III, 215
Vanilla, IV, 219
Vanilla claviculata, IV, 291
Vestitae, I, 44, 71
Visnaga, III, 131, 134
Visnager, III, 170
Weberianae, I, 44, 84
Weberocereus, II, 183, 214-216
biolleyi, II, 214, 215
panamensis, II, 214, 215
tunilla, II, 214
Werckleocereus, II, 183, 216, 217
glaber, II, 216
tonduzii, II, 216, 217; IV, 284
West Indian gooseberry, I, 10
Wilcoxia, I, 6; II, 2, 22, 110-112; III, 4
papillosa, II, 110, 112
poselgeri, II, 110, 111; IV, 274
striata, II, 110, 111; IV, 275
viperina, II, 110
Wilmattea, II, 183, 195, 196
minutiflora, II, 195, 196, 216
Wittia, IV, 177, 206, 207
amazonica, IV, 206
costancensis, IV, 213
panamensis, IV, 206, 207
Woolly nipple-cactus, IV, 71
Zacoub, II, 187
Zamia pumila, I, 181
Zanthoxylum, II, 92
Zehntnerella, II, 2, 176, 177
squamulosa, II, 176, 177
Zuwarro, II, 165
Zygocactus, I, 9; II, 22; IV, 177-180, 182, 183, 185, 200, 209
altensteinii, IV, 177, 178
candidus, IV, 182
delicatus, IV, 177, 180
obtusangulus, IV, 181
opuntioides, IV, 180, 181
truncatus, IV, 177, 178, 179, 183, 220

